



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



HARVARD UNIVERSITY  
DEPARTMENT OF  
GEOLOGY AND GEOGRAPHY



From the Library of  
**JAY BACKUS WOODWORTH**  
Class of 1894  
TEACHER OF GEOLOGY AT HARVARD  
FROM 1894 TO 1925

The Gift of  
G. S. HOLDEN      R. W. SAYLES  
R. A. F. PENROSE      E. WIGGLESWORTH

1926

Transferred to  
CABOT SCIENCE LIBRARY  
June 2005

B-297



**GUIDE TO THE MINERAL COLLECTIONS IN THE  
ILLINOIS STATE MUSEUM**



**STATE OF ILLINOIS**

**FRANK O. LOWDEN, GOVERNOR**

**DEPARTMENT OF REGISTRATION AND EDUCATION**

**FRANCIS W. SHEPARDSON, PH.D., LL.D., DIRECTOR**

---

**STATE MUSEUM DIVISION**

**A. R. CROOK, PH.D., CHIEF**

**SPRINGFIELD**

---

**BOARD OF MUSEUM ADVISERS**

**Administration and Fine Arts:**

**CHARLES L. HUTCHINSON, A.M.,** Corn Exchange National  
Bank, Chicago

**Botany:**

**CHARLES F. MILLSPAUGH, M.D.,** Field Museum, Chicago

**Ethnology:**

**CHARLES L. OWEN, A.B.,** Field Museum, Chicago

**Manufacturing and Business:**

**EDWARD W. PAYNE,** State Bank, Springfield

**Zoölogy:**

**HENRY B. WARD, PH.D.,** University of Illinois, Urbana

**A FREE INSTITUTION FOR THE PEOPLE :**

**TO PROMOTE SCIENCE**

**TO PROMOTE ART**

**TO PROMOTE BUSINESS**

***Hours:***

**Week Days: 8 A.M. till 12 noon, 1:30 P.M. till 5 P.M.**

**Saturdays: 8 A.M. till 3 P.M.**





*LETTER OF TRANSMITTAL*

STATE MUSEUM, SPRINGFIELD

August 31, 1919

*Francis W. Shepardson, LL.D.*

*Director, Department of Registration and Education*

DEAR SIR:

For the purpose of increasing the usefulness of the collections in the State Museum a series of guidebooks was planned a few years ago. The first to be printed was the *General Guide*, which appeared in 1914 and which is now exhausted. Herewith is submitted a *Guide to the Mineral Collections*, upon which the chief has been working in moments which could be spared from other work for the past several years.

Hoping that it may prove of service to students of mineralogy and also to those visitors whose interest is of a general character, I am

Yours very respectfully,

A. R. CROOK,

*Chief, State Museum Division*



QE  
386  
I5

GUIDE TO THE  
**MINERAL COLLECTIONS**  
IN THE  
ILLINOIS STATE MUSEUM

*By*

**A. R. CROOK, PH.D.**

*Chief, State Museum Division, Department of  
Registration and Education*

---

SPRINGFIELD, ILLINOIS  
1920

**Published October 1920**

**Composed and Printed By  
The University of Chicago Press  
Chicago, Illinois, U.S.A.**

## PREFACE

In the following pages an attempt has been made to so describe the minerals constituting our constantly growing collections as to emphasize the most important ones and at the same time to present to the reader a good idea of the science of mineralogy. The average visitor approaches the subject as a child would and just as the human race has done. When early man wandered up stream courses and found a gold nugget he doubtless was attracted by its yellow color, in time noticed the weight, softness, etc., and learned to use it as an ornament. The child does the same, using the senses of sight, feel, taste, and smell in making the acquaintance of any strange substance. Hence the physical characteristics—form, color, hardness, and weight—of various minerals are described, and then their chemical constituents, geological and geographical relations, and use are given. By becoming acquainted with various minerals the visitor obtains a knowledge of the science.

In preparation of this work the writer has used chiefly Dana's, Mier's, Lacroix's, and Tschermak's mineralogies and Tutton's and Groth's crystallographies, in addition to individual articles in U.S.G.S. reports, scientific journals, etc.

Especial thanks are due Professor O. C. Farrington for painstaking revision of the manuscript, Professor W. S. Bayly for careful reading of the proof, and the University of Chicago Press for the thorough manner in which their part of the work has been done.

Professor Farrington supplied photographs for Plates IV, XVIII, XIX*a*, and XXVII*a*. Plate VI*b* is after W. M. Foote, and Plate Ia is reproduced by permission of B. F. Buck & Co. The other illustrations are by the writer.

A. R. CROOK

August, 1919



## TABLE OF CONTENTS

---

	PAGE
LIST OF ILLUSTRATIONS . . . . .	xiii
ABBREVIATIONS . . . . .	xxi
INTRODUCTION . . . . .	I
CLASS I. ELEMENTS . . . . .	5
LIST OF ELEMENTS AND THEIR ATOMIC WEIGHTS . . . . .	45
CLASS II. SULPHIDES . . . . .	46
CLASS III. SULPHANTIMONITES, SULPHARSENITES . . . . .	68
CLASS IV. HALOIDS . . . . .	76
CLASS V. OXIDES . . . . .	82
CLASS VI. CARBONATES . . . . .	112
CLASS VII. SILICATES . . . . .	132
CLASS VIII. NIOBATES, TANTALATES . . . . .	175
CLASS IX. PHOSPHATES, ETC. . . . .	175
CLASS X. BORATES, URANATES . . . . .	177
CLASS XI. SULPHATES, CHROMATES, TELLURATES . . . . .	179
CLASS XII. TUNGSTATES, MOLYBDATES . . . . .	185
CLASS XIII. ORGANIC ACID SALTS . . . . .	187
CLASS XIV. HYDROCARBONS . . . . .	187
SUMMARY . . . . .	194
NAMES OF MINERALS . . . . .	196
THE USES OF MINERALS . . . . .	197
HISTORY OF THE STUDY OF MINERALS . . . . .	198
LIST OF MINERALS . . . . .	202
INDEX . . . . .	279





## LIST OF ILLUSTRATIONS

	PAGE
FIG. 1. MODEL OF OCTAHEDRON; PREVAILING OUTLINE OF DIAMOND . . . . .	5
FIG. 2. AXES . . . . .	6
FIG. 3. METHOD OF CONSTRUCTING CRYSTALLOGRAPHIC AXES . . . . .	7
FIG. 4. CONSTRUCTION OF AN OCTAHEDRAL PLANE . . . . .	7
FIG. 5. COMPLETED OCTAHEDRON . . . . .	8
FIG. 6. MODEL OF TRAPEZOHEDRON . . . . .	9
FIG. 7. COMPLETED TRAPEZOHEDRON . . . . .	9
FIG. 8. MODEL OF TRISOCTAHEDRON . . . . .	10
FIG. 9. TRISOCTAHEDRON COMPLETED . . . . .	10
FIG. 10. MODEL OF HEXOCTAHEDRON . . . . .	12
FIG. 11. HEXOCTAHEDRON COMPLETED . . . . .	12
FIG. 12. GROWTH OF UNSHADED PLANES OF THE WOODEN OCTAHEDRON PRODUCE THE GLASS TETRAHEDRON COVERING IT . . . . .	12
FIG. 13. RIGHT-HANDED TETRAHEDRON . . . . .	12
FIG. 14. WOODEN OCTAHEDRON INCLOSED BY GLASS TETRAHEDRON . . . . .	13
FIG. 15. LEFT-HANDED TETRAHEDRON . . . . .	13
FIG. 16. INTERPENETRATING SUPPLEMENTARY TETRAHEDRONS (TETRAHEDRAL TWINS) . . . . .	13
FIG. 17. TETRAHEDRAL TWIN WITH CORNERS TRUNCATED . . . . .	14
FIG. 18. MODEL OF RIGHT-HANDED HEXATETRAHEDRON . . . . .	15
FIG. 19. CONSTRUCTION OF HEXATETRAHEDRON . . . . .	15
FIG. 20. INTERPENETRATING TRUNCATED TETRAHEDRONS BEVELED BY HEXATETRAHEDRONS . . . . .	16
FIG. 21. MODEL OF "SPINEL TWIN" . . . . .	17
FIG. 22. DRAWING OF SPINEL TWIN . . . . .	17
FIG. 23. TWO KINDS OF AXES OF SYMMETRY . . . . .	18
FIG. 24. DOTTED LINES SHOW DIRECTION OF PLANES OF SYMMETRY IN CUBE . . . . .	19
FIG. 25. ANGLES OF INCIDENCE AND REFRACTION . . . . .	19
FIG. 26. MODEL OF ORTHORHOMBIC BIPYRAMID . . . . .	25
FIG. 27. ORTHORHOMBIC PYRAMIDAL PLANE AND AXES . . . . .	25
FIG. 28. OBTUSE BIPYRAMID CHARACTERISTIC OF SULPHUR . . . . .	26
FIG. 29. MODEL OF BRACHYDOMES AND MACROPINACOIDS . . . . .	26
FIG. 30. UPPER BRACHYDOME PLANES . . . . .	26
FIG. 31. MODEL OF MACRODOMES AND BRACHYPINACOIDS . . . . .	27
FIG. 32. UPPER MACRODOMES . . . . .	27

	PAGE
FIG. 33. PRISM AND BASAL PLANES . . . . .	27
FIG. 34. BASE, MACROPINACOID, AND BRACHYPINACOID . . . . .	28
FIG. 35. MODEL OF BASE, MACRO- AND BRACHYPINACOID . . . . .	28
FIG. 36. MODEL OF PRISM, BRACHYPINACOID, AND BRACHYDOME . . . . .	28
FIG. 37. RIGHT-HANDED SPHENOID . . . . .	28
FIG. 38. SULPHUR, USUAL HABIT . . . . .	29
FIG. 39. SULPHUR, SPHENOIDAL HABIT . . . . .	29
FIG. 40. AXES OF MONOCLINIC CRYSTAL . . . . .	30
FIG. 41. CUBE . . . . .	33
FIG. 42. TETRAHEXAHEDRON MODEL . . . . .	35
FIG. 43. CONSTRUCTION OF TETRAHEXAHEDRON . . . . .	35
FIG. 44. MODEL OF A DODECAHEDRON . . . . .	39
FIG. 45. CONSTRUCTION OF A DODECAHEDRON . . . . .	40
FIG. 46. STIBNITE CRYSTAL . . . . .	47
FIG. 47. CUBE TRUNCATED BY OCTAHEDRON . . . . .	49
FIG. 48. MODEL OF CUBE TRUNCATED BY OCTAHEDRON . . . . .	49
FIG. 49. GALENA, JO DAVIESS COUNTY, ILLINOIS . . . . .	50
FIG. 50. MODEL OF PLANES APPEARING ON GALENA . . . . .	50
FIG. 51. TWIN LAMELLAE IN GALENA . . . . .	51
FIG. 52. SPHALERITE . . . . .	53
FIG. 53. MODEL OF THREE-FACED TETRAHEDRON, A TRISTETRAHEDRON . . . . .	54
FIG. 54. CONSTRUCTION OF TRIGONAL TRISTETRAHEDRON . . . . .	54
FIG. 55. SPHALERITE . . . . .	54
FIG. 56. ACUTE PRIMARY BIPYRAMID . . . . .	57
FIG. 57. OBTUSE PRIMARY BIPYRAMID . . . . .	57
FIG. 58. MODEL OF SECONDARY BIPYRAMID . . . . .	58
FIG. 59. DITETRAGONAL BIPYRAMID . . . . .	58
FIG. 60. MODEL OF DITETRAGONAL BIPYRAMID . . . . .	58
FIG. 61. PRIMARY PRISM . . . . .	59
FIG. 62. SECONDARY PRISM . . . . .	59
FIG. 63. MODEL OF DITETRAGONAL PRISM . . . . .	59
FIG. 64. COMBINATION OF PRIMARY PRISM, SECONDARY PRISM, SEC- ONDARY BIPYRAMID, AND DITETRAGONAL BIPYRAMID . . . . .	59
FIG. 65. MODEL OF BISPHENOID . . . . .	60
FIG. 66. CHALCOPYRITE . . . . .	60
FIG. 67. CHALCOPYRITE, FRENCH CREEK, PENNSYLVANIA . . . . .	60
FIG. 68. CHALCOPYRITE, NEUDORF . . . . .	61
FIG. 69. PYRITOHEDRON DERIVED BY DISAPPEARANCE OF TETRA- HEXAHEDRAL PLANES DARKENED, AND GROWTH OF THE OTHER PLANES . . . . .	62
FIG. 70. MODEL OF A DIPLOID . . . . .	62
FIG. 71. MODEL OF COMBINATION OF PYRITOHEDRON AND CUBE . . . . .	63

# LIST OF ILLUSTRATIONS

XV

	PAGE
FIG. 72. MARCASITE . . . . .	64
FIG. 73. MARCASITE . . . . .	64
FIG. 74. "SPEARHEAD PYRITES" . . . . .	65
FIG. 75. "COCKSCOMB PYRITES" . . . . .	65
FIG. 76. ARSENOPYRITE . . . . .	67
FIG. 77. ARSENOPYRITE . . . . .	67
FIG. 78. SYMMETRY PLANES OF A DITRIGONAL POLAR CRYSTAL . . . . .	68
FIG. 79. AXES OF HEXAGONAL SYSTEM . . . . .	68
FIG. 80. MODEL OF PRIMARY PYRAMID . . . . .	69
FIG. 81. BASAL SECTION SHOWING RELATION OF PRIMARY, SECONDARY, AND DIHEXAGONAL PYRAMIDS AND PRISMS . . . . .	69
FIG. 82. MODEL OF DIHEXAGONAL BIPYRAMID . . . . .	69
FIG. 83. MODEL OF PRIMARY HEXAGONAL PRISM . . . . .	70
FIG. 84. SECONDARY HEXAGONAL PRISM . . . . .	70
FIG. 85. MODEL OF DIHEXAGONAL PRISM . . . . .	70
FIG. 86. RHOMBOHEDRON RESULTING FROM DISAPPEARANCE OF DARKENED PLANES OF THE INTERIOR FIGURE—HEXAGONAL BIPYRAMID . . . . .	71
FIG. 87. MODEL OF POSITIVE SCALENOHEDRON . . . . .	71
FIG. 88. MODEL OF SCALENOHEDRON TRUNCATED BY RHOMBOHEDRON . . . . .	71
FIG. 89. MODEL OF TRUNCATED PRISM . . . . .	71
FIG. 90. PYRARGYRITE, CRYSTAL FORM . . . . .	72
FIG. 91. PREVAILING FORM OF TETRAHEDRITE . . . . .	73
FIG. 92. CHARACTERISTIC FORM OF TETRAHEDRITE . . . . .	73
FIG. 93. TENNANTITE (SCHWATZITE) . . . . .	74
FIG. 94. MODEL OF SANDBERGERITE . . . . .	74
FIG. 95. HALITE CUBE FROM SALT BRINE . . . . .	77
FIG. 96. FLUORITE . . . . .	78
FIG. 97. FLUORITE . . . . .	78
FIG. 98. MODEL OF FLUORITE TWIN . . . . .	79
FIG. 99. QUARTZ; PRISM AND RHOMBOHEDRON . . . . .	83
FIG. 100. QUARTZ . . . . .	83
FIG. 101. A FORM OF QUARTZ COMMON AT ALSTON MOOR, ENGLAND . . . . .	83
FIG. 102. QUARTZ . . . . .	83
FIG. 103. QUARTZ; POSITIVE LEFT TRIGONAL TRAPEZOHEDRON . . . . .	84
FIG. 104. QUARTZ; POSITIVE RIGHT TRIGONAL TRAPEZOHEDRON . . . . .	84
FIG. 105. TWO RIGHT-HANDED QUARTZ CRYSTALS TWINNED . . . . .	85
FIG. 106. BRAZIL TWIN; RIGHT- AND LEFT-HANDED QUARTZ CRYSTALS INTERPENETRATING . . . . .	85
FIG. 107. BRAZIL TWIN. TWO RIGHT-HANDED CRYSTALS JUXTAPOSED . . . . .	86
FIG. 108. QUARTZ; BOURG DE OISANS TWIN . . . . .	86

	PAGE
FIG. 109. QUARTZ ETCHED WITH HYDROFLUORIC ACID; A, LEFT-HANDED; B, RIGHT-HANDED . . . . .	86
FIG. 110. AIRY'S SPIRAL IN RIGHT-HANDED CRYSTAL . . . . .	88
FIG. 111. CUPRITE . . . . .	92
FIG. 112. ZINCITE . . . . .	93
FIG. 113. CORUNDUM . . . . .	94
FIG. 114. CORUNDUM . . . . .	95
FIG. 115. CORUNDUM WITH TWINNING LAMELLAE PARALLEL TO R . . . . .	95
FIG. 116. CROSS-SECTION OF DICHRSCOPE . . . . .	96
FIG. 117. DIRECTION OF VIBRATION OF TWO RAYS OF LIGHT PASSING THROUGH CALCITE PRISM . . . . .	96
FIG. 118. RUBY . . . . .	97
FIG. 119. SAPPHIRE . . . . .	97
FIG. 120. MODEL OF RHOMBOHEDRON . . . . .	98
FIG. 121. HEMATITE . . . . .	99
FIG. 122. TABULAR HEMATITE CRYSTAL TWINNED PARALLEL TO THE PRISM . . . . .	99
FIG. 123. MANGANITE . . . . .	100
FIG. 124. MANGANITE . . . . .	100
FIG. 125. GOETHITE . . . . .	102
FIG. 126. GOETHITE . . . . .	102
FIG. 127. SPINEL TWIN . . . . .	105
FIG. 128. SPINEL . . . . .	105
FIG. 129. MAGNETITE . . . . .	106
FIG. 130. CASSITERITE . . . . .	108
FIG. 131. CASSITERITE . . . . .	108
FIG. 132. RUTILE . . . . .	109
FIG. 133. RUTILE TRIPLET . . . . .	110
FIG. 134. RUTILE OCTET . . . . .	110
FIG. 135. CALCITE. POSITIVE RHOMBOHEDRON; CLEAVAGE RHOMBOHEDRON . . . . .	113
FIG. 136. CALCITE. NEGATIVE RHOMBOHEDRON . . . . .	113
FIG. 137. CALCITE. NEGATIVE ACUTE RHOMBOHEDRON . . . . .	114
FIG. 138. CALCITE, SCALENOHEDRON . . . . .	114
FIG. 139. CALCITE, PRISM AND NEGATIVE OBTUSE RHOMBOHEDRON . . . . .	114
FIG. 140. CALCITE, SHOWING PRISM, NEGATIVE OBTUSE RHOMBOHEDRON, AND BASE . . . . .	114
FIG. 141. CALCITE; COMBINATION OF SCALENOHEDRON AND RHOMBOHEDRON . . . . .	114
FIG. 142. CALCITE RHOMBOHEDRON . . . . .	114
FIG. 143. CALCITE SCALENOHEDRON . . . . .	115
FIG. 144. CALCITE SCALENOHEDRON . . . . .	115

# LIST OF ILLUSTRATIONS

xvii

	PAGE
FIG. 145. CALCITE PRISM . . . . .	115
FIG. 146. CALCITE SCALENOHEDRON . . . . .	116
FIG. 147. CALCITE SCALENOHEDRON . . . . .	116
FIG. 148. CALCITE ETCHED WITH DILUTE HYDROCHLORIC ACID . . . . .	118
FIG. 149. DOLOMITE ETCHED WITH DILUTE HYDROCHLORIC ACID . . . . .	118
FIG. 150. ELASTICITY COEFFICIENT OF CALCITE . . . . .	119
FIG. 151. ELASTICITY COEFFICIENT OF DOLOMITE . . . . .	119
FIG. 152. GLIDE PLANES IN CALCITE . . . . .	119
FIG. 153. ARAGONITE . . . . .	123
FIG. 154. BASAL SECTION OF ARAGONITE TRIPLET . . . . .	124
FIG. 155. BASAL SECTION OF ARAGONITE; INTERPENETRANT TRIPLET . . . . .	124
FIG. 156. WITHERITE . . . . .	125
FIG. 157. CROSS-SECTION OF WITHERITE . . . . .	125
FIG. 158. CERUSSITE . . . . .	127
FIG. 159. CERUSSITE . . . . .	127
FIG. 160. THREE CERUSSITE CRYSTALS INTERPENETRATING PARALLEL TO PRISM PLANES . . . . .	127
FIG. 161. AXES OF MONOCLINIC CRYSTAL . . . . .	128
FIG. 162. MONOCLINIC BIPYRAMID . . . . .	129
FIG. 163. ORTHODOMES AND CLINODOMES . . . . .	129
FIG. 164. MODEL OF PRISM AND BASAL PLANE . . . . .	130
FIG. 165. MALACHITE SECTION . . . . .	130
FIG. 166. AZURITE CRYSTAL, SHOWING ALSO POSITION OF OPTIC AXES AND AXIAL PLANE . . . . .	131
FIG. 167. MODEL OF AN ORTHOCLASE CRYSTAL . . . . .	134
FIG. 168. ORTHOCLASE . . . . .	134
FIG. 169. ADULARIA ORTHOCLASE . . . . .	134
FIG. 170. MODEL OF CARLSBAD TWIN, INTERPENETRATING . . . . .	134
FIG. 171. BAVENO TWIN, COMPOSITION FACE . . . . .	134
FIG. 172. MANEBACH TWIN, COMPOSITION FACE . . . . .	134
FIG. 173. ORTHOCLASE SECTION; AXIAL PLANE, ANGLE OF EXTINCTION . . . . .	135
FIG. 174. ORTHOCLASE (ADULARIA); AXIAL PLANE . . . . .	136
FIG. 175. ORTHOCLASE; POSITIVE AND NEGATIVE DIRECTION OF EXTINCTION . . . . .	136
FIG. 176. TRICLINIC AXES OF COPPER SULPHATE; NO AXES AT RIGHT ANGLES . . . . .	139
FIG. 177. ALBITE . . . . .	140
FIG. 178. ALBITE, SHOWING ALBITE LAW . . . . .	140
FIG. 179. ALBITE, PERICLINE TWIN . . . . .	141
FIG. 180. ALBITE; AXIAL PLANE . . . . .	141
FIG. 181. ALBITE SECTION . . . . .	141

	PAGE
FIG. 182. ALBITE SECTION . . . . .	141
FIG. 183. RHOMBIC SECTION OF ANORTHITE . . . . .	142
FIG. 184. ANORTHITE, SHOWING POSITION OF AXIAL PLANE AND BISECTRIX . . . . .	142
FIG. 185. MICROSCOPIC SECTION OF LEUCITE BETWEEN CROSSED NICOLS . . . . .	145
FIG. 186 <i>a</i> . DIOPSIDE . . . . .	147
FIG. 186 <i>b</i> . PHOTOGRAPH OF DIOPSIDE FROM CANTLEY, QUEBEC, CANADA . . . . .	147
FIG. 187. DIOPSIDE . . . . .	148
FIG. 188. DIOPSIDE, SHOWING OPTIC AXES, ACUTE BISECTRIX, AXES OF ELASTICITY . . . . .	148
FIG. 189. AUGITE . . . . .	148
FIG. 190. AUGITE TWIN . . . . .	148
FIG. 191. AUGITE CROSS-SECTION PERPENDICULAR TO PRISM PLANES	149
FIG. 192. ENSTATITE, SHOWING PARALLEL EXTINCTION . . . . .	149
FIG. 193. DIOPSIDE, SHOWING OBLIQUE EXTINCTION ANGLE . . . . .	149
FIG. 194. RHODONITE . . . . .	151
FIG. 195. ANTHOPHYLLITE, AXIAL PLANE AND OPTIC AXES . . . . .	152
FIG. 196. HORNBLLENDE . . . . .	154
FIG. 197. HORNBLLENDE . . . . .	154
FIG. 198. HORNBLLENDE SECTION PERPENDICULAR TO PRISM . . . . .	154
FIG. 199. HORNBLLENDE . . . . .	154
FIG. 200. ANTHOPHYLLITE; PARALLEL EXTINCTION . . . . .	155
FIG. 201. TREMOLITE; EXTINCTION ANGLE . . . . .	155
FIG. 202. HORNBLLENDE; EXTINCTION ANGLE . . . . .	155
FIG. 203 <i>a</i> . PHOTOGRAPH OF BERYL FROM BRAZIL . . . . .	157
FIG. 203 <i>b</i> . BERYL . . . . .	157
FIG. 204. GARNET . . . . .	158
FIG. 205. GARNET . . . . .	159
FIG. 206. GARNET . . . . .	159
FIG. 207. ZIRCON . . . . .	161
FIG. 208. ZIRCON . . . . .	161
FIG. 209. TOPAZ . . . . .	162
FIG. 210. TOPAZ, SHOWING OPTIC AXES AND AXIAL PLANE . . . . .	162
FIG. 211. TOURMALINE . . . . .	164
FIG. 212. TOURMALINE . . . . .	164
FIG. 213. TOURMALINE, ANALOGOUS END . . . . .	165
FIG. 214. FOUR-TWINNED CRYSTALS OF STILBITE . . . . .	166
FIG. 215. STILBITE SHEAF . . . . .	166
FIG. 216. LEUCITE . . . . .	167
FIG. 217. NATROLITE . . . . .	168

# LIST OF ILLUSTRATIONS

xix

	PAGE
FIG. 218. MUSCOVITE . . . . .	169
FIG. 219. MUSCOVITE: PRESSURE FIGURE, PERCUSSION FIGURE, OPTIC AXES . . . . .	169
FIG. 220. BIOTITE; AXIAL PLANE . . . . .	171
FIG. 221. BASAL SECTION OF BIOTITE, SHOWING POSITION OF AXIAL PLANE AND PERCUSSION FIGURE . . . . .	171
FIG. 222. APATITE . . . . .	175
FIG. 223. BARITE . . . . .	179
FIG. 224. BARITE . . . . .	179
FIG. 225. BARITE . . . . .	179
FIG. 226. CELESTITE . . . . .	180
FIG. 227. ANGLESITE . . . . .	181
FIG. 228. ANGLESITE . . . . .	181
FIG. 229. GYPSUM . . . . .	182
FIG. 230. GYPSUM . . . . .	182
FIG. 231. GYPSUM . . . . .	182
FIG. 232. GYPSUM TWINNED BY JUXTAPOSITION . . . . .	183
FIG. 233. GYPSUM, TWINNED BY INTERPENETRATION . . . . .	183
FIG. 234. WOLFRAMITE . . . . .	185
FIG. 235. WULFENITE . . . . .	186
FIG. 236. WULFENITE . . . . .	186

PLATE I. *a*, PREVAILING FORMS OF THE DIAMOND. G. F. WILLIAMS COLLECTION; *b*, GLASS MODEL OF CULLINAN DIAMOND, THE LARGEST DIAMOND EVER FOUND.

PLATE II. CONSTRUCTION OF RIGHT-HAND UPPER OCTANT OF TRISOCTAHEDRON ABOVE AND OF TRAPEZOHEDRON BELOW.

PLATE III. CONSTRUCTION OF ONE OCTANT OF HEXOCTAHEDRON.

PLATE IV. DENDRITIC COPPER FROM CALUMET AND HECLA MINING REGION, MICHIGAN.

PLATE V. MUKEROP METEORITE, ONE-SIXTH NATURAL SIZE. FELL IN AMALIA-GOAMUS, WEST AFRICA.

PLATE VI. *a*, STIBNITE, JAPAN; *b*, MOLYBDENITE FROM ALDFIELD, PONTIAC COUNTY, QUEBEC, CANADA.

PLATE VII. *a*, GROUP OF PYRITE CUBES, SHOWING STRIATIONS, CENTRAL CITY, COLORADO; *b*, PYRITE. A PYRITOHEDRON AND CUBES, COLORADO.

PLATE VIII. MARCASITE, JO DAVIESS COUNTY, ILLINOIS.

PLATE IX. MARCASITE DISKS, GULF MINE, SPARTA, RANDOLPH COUNTY, ILLINOIS.



- PLATE X. MARCASITE, SHOWING RADIATED INTERNAL STRUCTURE.
- PLATE XI. MARCASITE COATING GALENA, MARSDEN MINE, JO DAVIES COUNTY, ILLINOIS.
- PLATE XII. FLUORITE GROUP FROM ROSICLARE, HARDIN COUNTY, ILLINOIS.
- PLATE XIII. *a*, FLUORITE CUBES, ROSICLARE, ILLINOIS; *b*, OCTAHEDRONS CLEAVED OUT BY TEN-YEAR-OLD BOY, SHOWING EASE AND REGULARITY OF CLEAVAGE.
- PLATE XIV. *a*, SMOKY QUARTZ, "CAIRNGORM," FROM MONTANA; *b*, QUARTZ, MONTGOMERY COUNTY, ARKANSAS.
- PLATE XV. QUARTZ GROUP, MONTGOMERY COUNTY, ARKANSAS.
- PLATE XVI. QUARTZ, "BOURG DE OISANS" TWIN, HOT SPRINGS, ARKANSAS.
- PLATE XVII. AMETHYST, THUNDER BAY, LAKE SUPERIOR.
- PLATE XVIII. MOSS AGATE, INDIA.
- PLATE XIX. *a*, BOTRYOIDAL HEMATITE, CUMBERLAND, ENGLAND; *b*, LIMONITE, HARDIN COUNTY, ILLINOIS.
- PLATE XX. CALCITE, WEBB CITY, MISSOURI.
- PLATE XXI. *a*, CALCITE, "DOG-TOOTH SPAR," JOPLIN, MISSOURI; *b*, CALCITE, "ICELAND SPAR," SHOWING DOUBLE REFRACTION.
- PLATE XXII. CALCITE, JOPLIN, MISSOURI.
- PLATE XXIII. CALCITE SCALENOHEDRON, ROSSIE, ST. LAWRENCE COUNTY, NEW YORK.
- PLATE XXIV. *a*, CALCITE, JOPLIN, MISSOURI; *b*, QUARTZ GEODE WITH LARGE FLAT RHOMBOHEDRAL CRYSTALS, ST. FRANCISVILLE, MISSOURI.
- PLATE XXV. *a*, ARAGONITE CRYSTALS FOUR INCHES IN DIAMETER, CIANCIANA, SICILY; *b*, STALACTITES, BISBEE, ARIZONA.
- PLATE XXVI. *a*, MICROCLINE, "AMAZON STONE," PIKE'S PEAK, COLORADO; *b*, MICROCLINE, PIKE'S PEAK, COLORADO.
- PLATE XXVII. *a*, GARNETS; DODECAHEDRON FROM SALIDA, COLORADO, TRAPEZOHEDRON FROM NORTH CAROLINA, AND COMBINATION FROM FORT WRANGEL, ALASKA; *b*, A DODECAHEDRON NEARLY FOUR INCHES IN DIAMETER FROM SALIDA, COLORADO.
- PLATE XXVIII. *a*, TOURMALINE DOUBLY TERMINATED; VARIOUSLY COLORED CRYSTAL FROM MESA GRANDE, CALIFORNIA; *b*, BLACK, WELL-CRYSTALLIZED SPECIMEN FROM HADDAM, CONNECTICUT.
- PLATE XXIX. *a*, APATITE, RENFREW, CANADA; *b*, BARITE, ALSTON MOOR, ENGLAND.
- PLATE XXX. GYPSUM, SHOWING FISHTAIL TWIN AND CURLED FORM.
- PLATE XXXI. GYPSUM, "SELENITE," WAYNE COUNTY, UTAH.

## ABBREVIATIONS

- $a, b, c$  = Crystallographic axes  
 $a, b, c$  = Direction of greatest, medium, and least elasticity  
 $\alpha, \beta, \gamma$  = Greatest, medium, and least index of refraction  
 $\alpha, \beta, \gamma$  = Angles between crystallographic axes  
 $\epsilon$  = Direction of the extraordinary ray or its index of refraction  
 $2 E$  = Apparent value of axial angle in the air  
 $\gamma - \alpha$  = Maximum birefracton  
 $2 H$  = Value of axial angle when mineral is immersed in oil  
 $n$  = Index of refraction  
 $\rho$  = Axial angle of red light  
 $2 V$  = True value of angle between optic axes  
 $\nu$  = Axial angle of violet (blue) light  
 $\omega$  = Index of refraction of the ordinary ray



## INTRODUCTION

Minerals play a large part in the annual increase in wealth and in the comfort of the inhabitants of Illinois. The state is not usually thought of as a mineral-producing region, as is Colorado, Montana, or California, and the fact is not usually known that the money value of the minerals obtained in this state exceeds that of any state west of the Mississippi River.

But such is the case. During the year preceding the world-war the total value of mineral products in Illinois amounted to more than one hundred and seventeen millions of dollars, while that of California, the nearest competitor, was only one hundred and one millions, and Colorado and Montana together fell even farther behind Illinois in mineral production.

The use of minerals is an index of civilization. Man is the only member of the animal kingdom to utilize minerals; and the more primitive his place in human society, the less does he do so.

The whole fabric of civilization depends upon iron, copper, gold, and other metals, and upon coal, building stone, and clays. The people of our state produce some of these substances in great quantities and use all kinds of minerals from all corners of the globe.

Some minerals occur in extensive deposits in the state, others are scattered here and there. The majority of those described in the following pages have been found within the region and many of the others are very useful for our people.

While more than a thousand different minerals are known, only about one hundred are common enough to claim our special attention. These one hundred are well illustrated in the museum collections.

Many visitors, while having a general idea of the subject, are unable to say just what a mineral is. Upon investigation they learn that a mineral is a natural, inorganic, homogeneous, solid, liquid, or gas. When solid, it is usually crystallized.

Artificial substances such as are produced in laboratories, chemical works, iron foundries, etc., are excluded from the definition, although they often show perfection of form and purity of constitution. Mineralogy is concerned with *natural* products.

The term inorganic excludes all forms of living substance—everything that grows by internal activity, that has the power of assimilation and reproduction, that has sensibility and usually slight chemical stability. A mineral may have had an organic origin. For example, the carbon of a piece of graphite may have been at one time in a tree. The tree died and with the loss of oxygen and hydrogen was converted into peat. The loss of oxygen and hydrogen continuing, the peat or lignite was changed into bituminous coal, then into anthracite, and finally into graphite. It is not the origin but its present condition which places a substance in the mineral kingdom.

The term homogeneous indicates that the substance throughout is the same at one point as another, has the same arrangement, and shows the same properties. This separates a mineral from other inorganic substances such as rocks. A rock is made up of a mass of minerals.

Usually a mineral is a solid. Some minerals—for example, water and mercury—are ordinarily liquid but may be changed into solids by freezing: water at  $32^{\circ}$  and mercury at  $-40^{\circ}$  F. All minerals are solid under certain conditions.

Minerals are usually crystallized; that is, they have a definite internal structure which is often shown by their external form. There are a few exceptions, such as turquoise and opal, and other substances which are solidified from gases or liquids so rapidly or under such other unfavorable conditions that the molecules are unable to properly arrange themselves. These minerals are said to be amorphous. They may be regarded as minerals that are unsuccessful or are of weak molecular attractions. Ordinarily a mineral has just as definite a shape as has a bird or a flower. It has less opportunity than a flower to develop a perfect external form, since it is usually crowded by its neighbors. The growing crystal soon reaches a place where its planes touch those of its neighbors and its perfection is impaired. But though the bounding planes are distorted and irregular, the internal arrangement is so orderly and definite that the smallest fragment has the same structure as a perfect crystal.

This regularity of architecture in the mineral world is a fact of far-reaching importance. It discloses one of the great laws of the universe, a law as beautiful and universal as the law of gravitation, the conservation of energy, or the development of species.

The law of crystallization affects every particle of mineral matter in the world, and more than that, in the universe as well. The results are seen alike in the minutest forms and on a gigantic scale. The most beautiful colors in the world—the pure colors of the spectrum—are exhibited by minerals in accordance with this law.

Minerals are the most abundant and most valuable substances in the world.

If all the vegetation in the world—the great masses of weeds in Sargasso Seas, the myriads of land weeds, the flowers and grains, all the trees of the mighty forests—if all of these were placed in an immense pile and to this pile were added all the lower animals, all mankind, and all the buildings in the world, the mass would be gigantic. But, if in another pile were heaped the minerals of which the world is composed, the first pile would be as a grain of sand to a mountain, so small as to be well-nigh invisible. In quantity minerals are of the greatest importance.

In quality the same is true. They are unsurpassed in enduring beauty and value. Some diamonds like the Kohinoor, Regent, or the Cullinan are valued more highly than any other objects of the same size in the world. A ruby worth half a million dollars is so light in weight that it could be sent by mail across the continent for two cents. Mineral ornaments such as vases, tables, and columns in the palaces of the wealthy and in the great museums will remain unchanged in beauty and pleasure-giving power for many long years. Minerals are as beautiful as flowers and infinitely more permanent.

Though the same sun with all diffusive rays  
Blush in the rose and in the diamond blaze,  
We prize the higher effort of his power  
And justly place the gem above the flower.<sup>1</sup>

An acquaintance with minerals is useful in many trades and professions. The doctor of medicine and the pharmacist may be interested in minerals as the source of drugs. The lawyer may be helped by some knowledge of mineralogy, especially in mining cases. The minister furnished by this science with an insight into the structure of the universe is better able to find "sermons in stones." From

<sup>1</sup> Alexander Pope.

the study of the mineral composition of his soil the farmer is aided in soil improvement and in making "bread from stones." The physicist repeatedly uses minerals in his study of the laws of heat, light, and electricity. Even more dependent upon minerals as a source of materials for study and experiment is the chemist. For the geologist, the prospector, the miner, the assayer, and the metallurgist, mineralogy is a fundamental science, one without which they cannot well work.

Thus the mineral collections in the museum have a twofold claim upon the interest of the visitor: first, because they well illustrate the mineral resources of this state, and second, because they show the composition of the world and the uses which our people make of minerals to increase the comfort of living and their happiness.

The visitor will naturally begin his inspection of this collection with the minerals that are the most simple in their composition—those that are composed of but one chemical substance, the so-called elements. He will find that while there are about two dozen of them which occur in some abundance as minerals, not more than twelve are common enough to claim his attention. These are: diamond, graphite, sulphur, arsenic, antimony, bismuth, gold, silver, copper, mercury, platinum, and iron. Each of these is noteworthy because of its beauty or utility or because it shows some peculiar property. All of them except antimony, bismuth, mercury, and platinum have been found in the state. Diamond, graphite, and sulphur are non-metals; antimony and bismuth, brittle metals; gold, silver, copper, platinum, and iron, malleable metals; and mercury, a liquid metal under ordinary conditions. Together, these minerals constitute the most prominent representatives of Class I.





PLATE I



*a*, Prevailing forms of the diamond. G. F. Williams collection



*b*, Glass model of Cullinan diamond, the largest diamond ever found

## CLASS I. ELEMENTS

### Diamond

There are several reasons for studying the diamond first, though Illinois is not a diamond-producing state. Not more than a dozen diamonds have been found here and they are immigrants brought in by glaciers which formerly slid down from the north, carrying all kinds of minerals collected from a wide area and scattering them here and there over two-thirds of the state. The chief source of the diamond is the Kimberley region in South Africa, but no people are more partial to the diamond as a gem than are the citizens of this state. Every woman has or expects to have one, and every man should at some time buy one!

The diamond is easily premier among gems. It is a fine example of a successful mineral. Its character is positive. It deserves the most extended study. While studying it we gain an insight into the whole mineral world. The Illinois State Museum contains a few examples of the diamond, and glass models of the most famous diamonds of history. If one examines a handful of diamonds as they are taken from the mines at Kimberley (Plate I) or as they come uncut to Amsterdam, London, or New York City, he will observe that the greater number of them are shaped like two pyramids placed base to base forming an eight-sided figure called an octahedron (Fig. 1). Some of them are flat, triangular flakes, others globules, and nearly all are somewhat distorted and pitted, with some planes well developed and others small. The larger faces were formed on that side of the crystal which had




FIG. 1.—Model of octahedron; prevailing outline of diamond.

the most abundant material to draw from, while the small faces, like the smaller birds in a nest, receiving the least food, have not had equal opportunity for growth. But while the different faces vary in size, the angles between them are always the same. They are said to be constant, and illustrate the "law of constancy of angle"—a law of far-reaching importance, since because of it minerals can readily be identified and classified.

The natural shape of a mineral is one of the first characters to notice. What anatomy is to the student of the human body, cell structure to the morphologist, and architecture to the builder, crystal form is to the mineralogist. It is one of the fundamentals. The purpose of study of the form of minerals is not only to recognize and

picture the external form but to understand the internal structure as well, since they are dependent upon each other.

The architecture of the diamond may be better understood if the planes which occur on natural crystals can be represented by a drawing. Fortunately it is only necessary to measure off certain points and connect them by straight lines—a much simpler task than it would be to draw the structures seen in the plant and animal




FIG. 2.—Axes

worlds. Anyone can draw the shapes which diamonds exhibit.

First draw three lines or *axes* which intersect each other at right angles (Fig. 2). In drawing these figures we use the method most generally employed, which is called "clinographic projection." The eye is supposed to be elevated a trifle above the crystal and removed an infinite distance so that the lines in the drawing do not converge as in ordinary perspective. Those parallel in the crystal are parallel in the drawing.

To erect the axes, a templet cut out of cardboard may be constructed in the following manner (Fig. 3):

Draw a vertical line  $MM'$  and  $NN'$  at right angles to it. Divide  $NN'$  into six equal divisions. At the second and fourth divisions

draw lines parallel to  $MM'$ . From  $N'$  mark  $N'O$  equal to one division. From  $O$  draw a straight line through  $P$  to  $O'$ .  $aa'$  is the front to back axis of our crystal. From  $a$  draw  $aR$  parallel to  $N'N$ . From  $R$  draw  $RP$ . From  $S$  draw  $Sb$  parallel to  $NN'$ . From  $b$  draw  $bP$  and extend to  $\bar{b}$ .  $bb'$  is the horizontal axis extending from right to left. Twice  $OP$  gives the length of the  $c$  axis. These axes form the foundation for the construction of the axes used all through the work. An excellent discussion of the subject may be found in Tutton's *Crystallography*.<sup>1</sup> We always call the vertical line (Fig. 2)  $c$ ; the horizontal




FIG. 3.—Method of constructing crystallographic axes.




FIG. 4.—Construction of an octahedral plane.

line, extending from right to left,  $b$ ; and the one extending from the front backward,  $a$ . The upper half of the  $c$  axis, the right half of  $b$ , and the front of  $a$  are said to be positive; the others, negative. To draw any given plane mark the points at which it would intersect the axes. In the octahedron (Fig. 1) each plane intersects the three axes at points equally distant from the center. Then to draw an octahedral plane measure off equal distances on each axis and connect these points with straight lines (Fig. 4). To complete the octahedron, which has eight such planes, draw similar planes in each of the other octants (Fig. 5).

<sup>1</sup> See pages 382-439.

The relation of the axes to each other was expressed by an English mineralogist, W. H. Miller, of Cambridge (d. 1880), as a ratio,  $a:b:c$ . The portion measured off on each axis is written as a denominator. Then the ratio which represents the octahedral plane is  $\frac{a}{1}:\frac{b}{1}:\frac{c}{1}$ , and its symbol is (111) which is read as one, one, one. When simply (111) is used, the right-hand upper octant is meant. The




FIG. 5.—Completed octahedron

left-hand upper octant would have the symbol ( $\bar{1}\bar{1}1$ ) [read one, minus one, one]; the right-hand lower ( $11\bar{1}$ ), the left-hand lower ( $\bar{1}1\bar{1}$ ), the right-hand upper back ( $\bar{1}\bar{1}\bar{1}$ ), the left-hand upper back ( $111$ ), etc.


The diamond, like some other minerals, is symmetrically built. It has the same molecular structure in all directions. Light, heat, and electricity travel through it with the same ease and rapidity in the direction of all three of the axes, and the corrosive action of sol-

vents is the same in all parts. Its axes are of the same value and interchangeable, and hence a numeral like 1 may be substituted for the letters  $a$ ,  $b$ , and  $c$  and the ratio  $\frac{a}{1}:\frac{b}{1}:\frac{c}{1}$  then becomes  $\frac{1}{1}:\frac{1}{1}:\frac{1}{1}$ . This cleared of fractions yields 1:1:1. The numbers in this ratio, 1:1:1, constitute the "parameters" of the octahedral plane, since they define the position of the plane. The parameter of another plane might be 1:2:2 or 2:1:2, etc.

The plane which has the parameter 1:2:2 represents a ratio  $\frac{1}{2}:\frac{1}{2}:\frac{1}{1}$  and its symbol is (211) (Fig. 6). Since each of the three axes are equal, we can apply the 2 to each axis in turn and hence obtain three planes in every octant.



PLATE II



Construction of right-hand upper octant of a trisectahedron above, and of a trapezohedron below.

Parameters	Ratios	Symbols
$1:2:2$	$\frac{1}{2}:\frac{1}{1}:\frac{1}{1}$	$(211)$
$2:1:2$	$\frac{1}{1}:\frac{1}{2}:\frac{1}{1}$	$(121)$
$2:2:1$	$\frac{1}{1}:\frac{1}{1}:\frac{1}{2}$	$(112)$

To construct these three planes in the right-hand upper octant (Plate II, lower diagram) measure off unit's distance on  $a$  and draw a line (red) cutting  $b$  at twice unit's distance, one cutting  $c$  at twice unit's distance, and one connecting the ends of  $b$  and  $c$ . Then from unit's



FIG. 6.—Model of trapezohedron




FIG. 7.—Completed trapezohedron

distance on  $b$  draw lines (blue) cutting  $a$  and  $c$  at twice unit's distance, and connect the ends. Finally, beginning at unit's distance on  $c$  draw lines (green) cutting  $a$  and  $b$  at twice unit's distance and connect the ends. These lines will determine the position of planes which will intersect each other so as to form three trapezoids in the octant. When the same plan is followed for the other octants there results a trapezohedron (Figs. 6 and 7).

Having constructed figures with the symbol  $(111)$  and  $(211)$ , the next in order will be one with the symbol  $(221)$  (Fig. 8). Its ratio will be  $\frac{a}{2}:\frac{b}{2}:\frac{c}{1}$  and its parameter  $1:1:2$ . Just as in the case of the



trapezohedron, the numerals may be applied to each of the axes in turn. Thus writing the parameters, ratios, and symbols for the right-hand upper octant we obtain the following:

Parameters	Ratios	Symbols
1:1:2	$\frac{1}{2}:\frac{1}{2}:1$	(221)
2:1:1	$\frac{1}{1}:\frac{1}{2}:\frac{1}{2}$	(122)
1:2:1	$\frac{1}{2}:\frac{1}{1}:\frac{1}{2}$	(212)

Draw a line (red) cutting  $a$  and  $b$  at unit's distance and  $c$  at twice unit's distance (Plate II, upper diagram). From unit's distance




FIG. 8—Model of trisectahedron





FIG. 9.—Trisectahedron completed

on  $b$  draw lines (blue) cutting  $a$  at twice unit's distance and  $c$  at unit's distance. From unit's distance on  $c$  draw lines (green) cutting  $a$  at unit's and  $b$  at twice unit's distance. These lines determine the position of planes which intersect within the octant so as to produce triangles. The resulting planes are called trisectahedral planes and the figure produced by continuing the process in each octant is called the trisectahedron (Figs. 8 and 9).



# PLATE III



Construction of one octant of hexoctahedron

The next form in point of complexity is one whose planes intersect each axis at different distances (Fig. 10). For example, its parameter might be  $1:\frac{3}{2}:3$ ; its ratio then would be  $\frac{1}{3}:\frac{1}{2}:\frac{1}{1}$ , and its symbol (321). Since each number in the parameter is different, each of the three axes would be intersected at two points different from unity and there would result two planes at each corner of the octant, making six planes where the octahedron has but one.

Writing the parameters, ratios, and symbols as before, the following result:

Parameters	Ratios	Symbols
$1:\frac{3}{2}:3$	$\frac{1}{3}:\frac{1}{2}:\frac{1}{1}$	(321)
$3:1:3$	$\frac{1}{2}:\frac{1}{3}:\frac{1}{1}$	(231)
$3:1:\frac{3}{2}$	$\frac{1}{1}:\frac{1}{3}:\frac{1}{2}$	(132)
$3:\frac{3}{2}:1$	$\frac{1}{1}:\frac{1}{2}:\frac{1}{3}$	(123)
$\frac{3}{2}:3:1$	$\frac{1}{2}:\frac{1}{1}:\frac{1}{3}$	(213)
$1:3:\frac{3}{2}$	$\frac{1}{3}:\frac{1}{1}:\frac{1}{2}$	(312)

Proceed in the construction as was done with the octahedron, trapezohedron, and trisectahedron. To construct, begin at unit's distance on  $a$  (Plate III) and draw a line (red) cutting  $b$  at three halves unit's distance and one cutting  $c$  at three times unit's distance. Complete the triangle by joining  $\frac{3}{2}b$  and  $3c$ . Begin at unit's distance on  $b$  and draw a line (blue) cutting  $a$  at three halves unit's distance and one cutting  $c$  at three times unit's distance. Join  $\frac{3}{2}a$  and  $3c$ . Again from unit's distance on  $b$  draw other lines (blue dotted) cutting  $a$  at three times unit's distance and  $c$  at three halves. Join the ends. Continue the construction from  $c$  (with green) and  $a$  (with red dotted) as indicated and the six planes of the octant will be produced. They are called the hexoctahedral planes. The same operation repeated in each octant produces a hexoctahedron (Figs. 10 and 11).

The faces above described are those most characteristic of the diamond, but usually faces of any one form do not make up the whole crystal. Sometimes planes of one form predominate and small



FIG. 10.—Model of hexoctahedron




FIG. 11.—Hexoctahedron completed



FIG. 12.—Growth of unshaded planes of the wooden octahedron produce the glass tetrahedron covering it.




FIG. 13.—Right-handed tetrahedron

faces of another modify the corners. Often two crystals will interpenetrate, each crystal having only half of its faces developed. If the right-hand upper octant and every alternate octant of the octahedron should grow to the exclusion of the other faces, a tetrahedron

(Figs. 12 and 13) would result. If the left-hand upper octant and each alternate octant were developed, a left-handed or negative tetrahedron would be produced (Figs. 14 and 15).



FIG. 14.—Wooden octahedron inclosed by glass tetrahedron.




FIG. 15.—Left-handed tetrahedron

When two supplementary tetrahedrons interpenetrate, the form represented in Figure 16 results. It is called an interpenetrating tetrahedral twin. Where an octahedron would have sharp edges, these tetrahedral twins have re-entrant angles.

Now if the projecting corners of each tetrahedron were truncated by the faces of the other tetrahedron, a form resembling an octahedron would result, but the re-entrant angles, instead of the characteristic edges, would reveal its true structure. This is a very common occurrence in the diamond (Fig. 17).




FIG. 16.—Interpenetrating supplementary tetrahedrons (tetrahedral twins).

Just as with the octahedron, so also with the trapezohedron, trisoctahedron, or the hexoctahedron, a portion only of the faces might be developed. If the right-hand upper octant and every alternate octant of a hexoctahedron were produced at the expense of their neighbors, a right-handed hexatetrahedron would result (Figs. 18 and 19).




FIG. 17.—Tetrahedral twin with corners truncated

A left-handed or negative hexatetrahedron would be produced if the left-hand upper and every alternate octant were developed at the expense of their neighbors. A right-handed and a left-handed hexatetrahedron interpenetrating and having the corners truncated by tetrahedral planes give rise to one of the most characteristic forms of the diamond (Fig. 20).

Upon taking up a diamond, one first notices the prevailing octahedral form, but closer inspection reveals the re-entering angles, and in these angles the slightly inclined hexatetrahedral planes may be recognized.

Another form of twinning in the diamond is that which results when an octahedron is cut through the middle by a plane parallel to an octahedral face and one-half of the octahedron is turned  $90^\circ$  (as shown in Figs. 21 and 22). Diamonds of this sort are called "suture" stones by diamond dealers and by crystallographers "spinel twins," since they are even more commonly found among specimens of the mineral named spinel.

All of the above forms—the octahedron, the trapezohedron, trisoctahedron, hexoctahedron, tetrahedron, and hexatetrahedron—agree in this, that they are symmetrical in the same directions. If



FIG. 18.—Model of right-handed hexatetrahedron.




FIG. 19.—Construction of hexatetrahedron.

any of these forms were divided parallel to these directions, one-half would be just like the other.

These directions are: first, parallel to any one of the three crystallographic axes,  $a$ ,  $b$ , and  $c$  (Fig. 23); second, parallel to any one of the four axes perpendicular to the octahedral planes; third (Fig. 24), parallel to the six planes which would pass through the edges of a cube. Since this symmetry is best represented in a mineral called tetrahedrite, it is named the tetrahedrite class of symmetry.

By the above study we have become acquainted with facts in regard to the crystallography of the diamond and, more than that, with facts which are needed to understand the forms of a hundred



other minerals as well. All of these minerals agree in this, namely, that the molecules which compose them arrange themselves similarly along three lines of equal length at right angles to each other, the  $a$ ,  $b$ , and  $c$  axes. Hence these minerals are placed together in one of the six great groups in which all minerals are classified—the group known as the Regular System.




FIG. 20.—Interpenetrating truncated tetrahedrons beveled by hexatetrahedrons

As wood has certain directions in which it readily splits and contrary to which it breaks in an irregular manner, so the diamond has a direction in which it readily splits or “cleaves,” namely, parallel to the octahedral planes. By taking advantage of the cleavage, diamond cutters are more readily able to fashion the gem into the desired shape. Cleavage is so easily obtained that it is difficult to break or “fracture” the diamond. When it is broken and not cleaved or split, the fractured surfaces are pitted or rounded like a shell. Consequently the fracture is said to be conchoidal.

Contrary to popular report, the diamond is brittle and easily shattered. Many valuable gems have been destroyed by the finders,

who failed to recognize the difference between hardness and tenacity. For centuries it has been a tradition that if a diamond were laid upon an anvil and struck by a hammer, both anvil and hammer would fly to pieces before the diamond was broken. Pliny said that the only way to "subdue" a diamond is to "soften it in warm goat's blood"! Although the diamond is brittle, it is the hardest of minerals.




FIG. 21.—"Spinel twin" model. Twinning plane (111).




FIG. 22.—Drawing of spinel twin

#### Scale of Hardness

1. Talc
2. Gypsum
3. Calcite
4. Fluorite
5. Apatite
6. Orthoclase
7. Quartz
8. Topaz
9. Corundum
10. Diamond

To measure the hardness of minerals a scale has been arranged which consists of ten minerals so chosen that, beginning with the softest, each succeeding mineral is hard enough to scratch the one before it in the scale. Talc, which is the softest, is given as No. 1 in this list, and diamond as No. 10. The finger-nail can scratch any mineral below 3, and a knife-blade any below 6.

When the weight of the diamond is compared with that of an equal volume of water, the diamond is found to be three and one-half times as heavy as water, i.e., its specific gravity is 3.5. It is much heavier than glass (sp. gr. about 2.5), which is most commonly employed to imitate it, or quartz (sp. gr. 2.6), the most abundant mineral that resembles diamond, or phenacite, "the

#### Specific Gravity of

1. Glass 2.5
2. Quartz 2.6
3. Phenacite 2.9
4. Topaz 3.5

deceiver" (sp. gr.=3), which is sometimes worn to represent the more valuable gem.

The diamond shows color from two causes: first, because of actual coloring materials in it; and second, because it divides a ray of entering light into the colors of the spectrum. The coloring matter is usually some metallic oxide which does not change when heated. Sometimes the coloring matter is organic material which does fade when heated or held in sunlight. Many shades are seen—red, yellow, green, blue, indigo, brown. Yellow and brown are the most common among African diamonds. Brown may deepen into black, as in the opaque "carbons." Green is not so common. Blue and red are the rarest, and when these colors are pure, the diamond exhibiting them is the most valuable gem in the world. About half of all diamonds found are white or colorless.




FIG. 23.—Two kinds of axes of symmetry

The color or "fire" which they then show is due to the fact that a

ray of light which enters at an angle is refracted or turned very markedly from its course and is divided into rays of different wavelengths. The result obtained by dividing the angle which the entering light makes with a perpendicular erected to the surface of the mineral (called the angle of incidence,  $i$  in Fig. 25) by the angle which the ray makes with the perpendicular prolonged after it has entered the mineral (called the angle of refraction,  $r$  in the figure) is called the index of refraction. Thus

$$n = \frac{\sin i}{\sin r}.$$

The index of refraction  $n$  has a different value according to the kind of light used. For blue light in the diamond it is 2.465; for red, 2.402. That is, a blue ray is turned farther from a straight line than is a red ray. Now the difference between these indexes, 0.063, is called the dispersion. Both the refraction and dispersion of diamonds are high in comparison to the refraction and dispersion of other minerals. Because of its high refraction the diamond is unusually brilliant. It is said to show "life." Because of its high dispersion it has a remarkably vivid play of colors or "fire."

*Mean Refractive  
Index of*

Ice	1.31
Salt	1.54
Quartz	1.55
Topaz	1.62
Glass	1.80
Cinnabar	3.02

*Dispersion of*

Fluorite	.006
Quartz	.025
Diamond	.063




FIG. 24.—Dotted lines show direction of planes of symmetry in cube.




FIG. 25.—Angles of incidence and refraction.

Some diamonds which have a bluish tinge after being held in the sunlight emit light in the dark, i.e., become phosphorescent. Many phosphoresce after being rubbed on wood or while being subjected to an electric discharge in a vacuum or while exposed to radium emanations. Positive electricity is developed in the diamond by friction.

The diamond is worthy of its place as the leading gem, not only because of its hardness, brilliancy, and beauty of color, but also because of its permanency in the presence of corrosive gases and liquids. The air and moisture do not affect it. Ordinary acids

cannot dissolve it. A solvent composed of sulphuric acid and potassium bichromate acts upon it slowly. At a high temperature ( $900^{\circ}\text{C}.$ ) in an oxygen flame, the diamond burns to carbon dioxide just as pure charcoal does, and sometimes leaves an extremely light ash that retains the original crystal shape and consists of iron, calcium, magnesium, and silicon which were present as impurities.

Though used as a gem for many hundreds of years, it is only within comparatively recent times that cutting of the diamond has been resorted to to enhance its beauty. Cutting and polishing are accomplished by the use of diamond dust imbedded in a tin disc or an iron plate.

Besides its use as a cutting tool, in its less well-crystallized forms it is extensively used to make drills. The diamond drill is one of the most useful implements invented for piercing rock in the search for valuable ores. The drills are made from the forms of diamond called bort and carbonado. Bort has radial fibrous structure, curved drusy surface, and is dark in color. Carbonado is somewhat compact, altogether without cleavage, slightly porous, black like charcoal, and somewhat harder than ordinary diamond. It is valuable for drills.

Diamonds have been found in nearly every part of the globe, yet for some cause or other in quantities worth mentioning only in regions less than  $30^{\circ}$  from the equator. A line following their most abundant occurrence would begin in southeastern Australia and extend eastward to South America; there, dividing into two branches, would pass in one branch north of the equator to British Guiana, and in the other, south to Bahia in Brazil. The northern branch extended would finally reach India, most prolific of ancient localities, and the southern branch would reach South Africa at Kimberley, the most productive of modern regions.

In India diamonds were first found and prized as gems. Most of the famous historical gems came from there. That field is now exhausted. The New South Wales fields yield about sixty thousand dollars' worth a year, British Guiana twice that amount, and Brazil five times that amount, while South Africa so far surpasses these regions as to make them hardly worth mentioning. It easily produces twenty-five million dollars' worth annually.

In India, Australia, South America, and also in a few localities in the Urals, and in Wisconsin, Michigan, Illinois, etc., diamonds

have been found in sands and gravels in which they were imbedded when the original rock containing them was broken up and transported by glaciers or flowing streams.

In the Kimberley region, however, they occur in a greenish-blue igneous rock called *kimberlite*, a sample of which is shown in Case 1.

The kimberlite occupies crater-like basins, sometimes nearly half a mile in diameter and of unknown depth, in the Triassic rocks of that region.

In Arkansas kimberlite has been found and has yielded a few diamonds.

Meteorites often contain minute diamonds.

Glass models of diamonds of unusual size and value, several of which have been long known and owned by kings and other celebrities, are shown in Case 1. No diamond is more famous than the Kohinoor. It weighs 106 carats and is valued at more than half a million dollars. It was found in India six hundred years ago (1304 A.D.) and for centuries was fought for or purchased by various rulers. It is now exhibited among the English crown jewels in London.

The Regent, which weighs 136 carats and is valued at six hundred thousand dollars, has had a most eventful history from the time when it was found in India by a slave, stolen from him by a sailor, bought and sold at ever-increasing price, till it came to sparkle in the hilt of Napoleon's sword, and finally, as one of the gems of the French Republic, was placed where it could ever after be inspected by all who cared to see it in the priceless collections of the Louvre, Paris.

The next is the Orloff, 193 carats, valued at half a million dollars, shaped so as to fit in the eye socket of an idol in India, from which it was stolen by a French soldier, finally purchased by Prince Orloff who sold it to Catherine II of Russia; and when last heard of it was in the Winter Palace, Petrograd.

Still larger is the Jubilee, 239 carats, valued at two million dollars, found in South Africa in 1895, and now in England.

The largest diamond ever found was the Cullinan (Plate I), named after the discoverer of the Premier Diamond Mine, South Africa. It was picked up in a shallow pit in that mine in January, 1905, by a foreman, who was given ten thousand dollars for his good

fortune in recognizing it. It weighed uncut  $3,025\frac{3}{4}$  carats, or about one and one-quarter pounds, and was valued at three million dollars. The Transvaal Republic donated it to King Edward VII of England, who, it was hoped, would place it unaltered in a museum to show for all time the largest diamond ever found. But the king had it cut into eleven brilliants—four of which are yet larger than any others known. Many diamonds smaller than those mentioned above are interesting because of their history.

The South Star, the largest diamond found in Brazil (in 1853) weighs 125 carats, is valued at four hundred thousand dollars, and is now owned by a prince in Baroda, India.

The Shah of Persia, 86 carats, now in Petrograd, has the shape of a four-sided prism with inclined ends.

No diamond has had a more varied history than the Sancy. It was found in India and, after being in the possession of Charles the Bold, who lost it on a battlefield, then among the jewels of the French Count De Sancy, then of Queen Elizabeth, then of Louis XIV, then of the King of Spain, and then of a Russian Prince Demidoff, has again been taken back to India by a native prince.

All of the above-named are beautiful white stones. There are some colored ones greatly prized, as, for example, the yellow Florentine and the blue Hope diamond. The Florentine, found in India, is now among the Austrian state jewels, after having been owned by Charles the Bold, Pope Julius, and the Empress of Austria. The Hope diamond is the most famous of all colored stones. It is of a vivid blue. It has been owned by wealthy and titled people in Italy, France, and England, and now is in the possession of a family in Washington, D.C.

#### SUMMARY

*Diamond*.—C. Regular; tetrahedrite class of symmetry: (111), (321); supplementary twins of tetrahedrons and contact twins on (111). Cleavage parallel (111) perfect; brittle; fracture conchoidal.

Hardness=10; gravity=3.52.

Colorless, yellow, brown, purple, red, blue; luster, greasy; transparent; refraction very strong.  $n=2.417$ ; dispersion very strong=0.063.

Infusible; soluble in sulphuric acid and potassium bichromate.

Australia, South America, South Africa.

### Graphite

Graphite (*γράφειν*, "to write"), another form of pure carbon, though not abundant in Illinois, is scattered in flakes through gneisses and other rocks which have been strewn over the state. It is a mineral so useful in many of our activities that it could not be spared. What, for example, would the people of the state do without lead pencils, lampblack, stove polish, graphite paints, and lubricants? The specimens on exhibit came from New Jersey, Ottawa, Canada, and Ceylon, of which localities the last has yielded the greatest quantities of the finest quality.

While the diamond is very pronounced in its shape, graphite is a mineral of weak molecular attraction. Its external form is not marked and, since it is opaque and cannot be studied under the microscope, there is even some doubt as to what its system of crystallization really is, although it is classified as hexagonal.

Usually it occurs in leafy or scaly flakes disseminated in rocks that originally were like limestone but have been changed by heat into marble or into other metamorphic rocks. Often it is segregated into compact, granular, or earthy masses forming veins in gneisses and schists.

While diamond is the hardest of substances, there is no mineral softer than pure graphite; while diamond is transparent and of light color, graphite is opaque and black; while diamond is a non-conductor of electricity, graphite is a good conductor. These differences are probably due to different arrangement of the atoms in the molecule, the graphite molecule containing three atoms, while that of the diamond contains nine.

Graphite is greasy to the touch, flexible, 1 in hardness, 2 in specific gravity, infusible, and insoluble.

Its specific heat is similar to that of the diamond. By specific heat is meant the heat required to raise one gram of a substance through one degree Centigrade. Taking as the unit the amount of heat required to raise one gram of water one degree in temperature, graphite requires only .12 and diamond .18 as much.

In the electric arc, diamond can be converted into graphite or graphite to diamond by varying the conditions.

Graphite is chiefly used for pencils, stove polish, paint, crucibles, and lubricants. In early days only the purest graphite could be



employed for "lead pencils" since the "leads" were cut out of the solid material. Now material containing much foreign matter is pulverized and washed to free it from impurities, mixed with clay, and burned. The amount of clay used and the heat employed determine the degree of hardness of the pencil. Graphite is a valuable paint where heat is to be resisted. For the same reason—because of its extreme infusibility, and for its reducing action, i.e., its tendency to keep oxygen away from molten metals—it is employed for crucibles. As a lubricant it is useful for heavy machinery or wherever heat would destroy other lubricants. All of these uses are well illustrated in the graphite case.

#### SUMMARY

*Graphite*.—C. Hexagonal (?); in plates, scales, masses. Cleavage basal, perfect; flexible.

Hardness=1; gravity=2; black; streak gray; luster, metallic.

Ceylon, Siberia, Canada, Mexico.

#### Sulphur

One of the most interesting sights in one of America's most charming parks—the Yellowstone National—is a group of hills, the highest of which rises a few hundred feet above the surrounding country and is called "Sulphur Mountain." It is composed of siliceous and calcareous material mingled with vast quantities of sulphur. From the yellowish-gray mass here and there sulphurous vapors arise; and in many places sulphur springs burst forth and run in rivulets down the side of the hill, leaving behind a yellowish-white trail. In many places on the hill the sulphur is quite pure, earthy in character, and yellowish-gray in color. The cavities from which fumes are escaping are often lined with deposits of pure yellow sulphur that hang in clustered crystalline masses and have been formed by the sublimation of sulphurous vapors.

Sublimation deposits characteristic of volcanic regions do not furnish fine crystals such as may be obtained from Sicily, where in the marly limestones hot waters laden with sulphur in solution have deposited their burden under favorable conditions and produced large crystals. Samples of sulphur from the Yellowstone and from Sicily are shown. Illinois occurrences are limited to whitish masses remaining from decomposition of iron sulphide or calcium sulphate.

The structure of the sulphur crystal differs from that of a diamond, since sulphur has three well-defined directions in which light, heat, electricity, and various chemical reagents act with different ease and rapidity. Hence a sulphur crystal is represented by three axes of different lengths, which cross each other at right angles. These axes characterize the Orthorhombic System. In the regular system we found all axes equal; in the orthorhombic they are all unequal. The vertical axis  $c$  may be greater or less than the lateral axis  $b$ . Of the two lateral axes, the longer is always chosen as the  $b$  axis and the shorter as  $a$ .



FIG. 26.—Model of orthorhombic bipyramid




FIG. 27.—Orthorhombic pyramidal plane and axes.

Since these axes represent different lengths and values, they cannot be interchanged as they were in the regular system. Upon them three different kinds of planes may be constructed: first, those which intersect all the axes; second, those which intersect two axes and are parallel to the third; and third, those which intersect one axis and are parallel to two.

1. Planes which intersect three axes are called pyramid planes. They correspond to the octahedral planes of the regular system. When they intersect all three axes at unit's distance, the typical bipyramid results (Fig. 26). Figure 27 shows the construction of the pyramidal plane.

If the  $c$  axis is intercepted at one-third unit's distance, as is often the case with some planes that are found on sulphur, an obtuse bipyramid is produced (Fig. 28).




FIG. 28.—Obtuse bipyramid (113) characteristic of sulphur.

2. Besides pyramid planes occur the so-called dome planes (from *domus*, "house," since they are like a roof). They intersect two axes and are parallel to one of the lateral axes (Figs. 29–32). The plane which is parallel to the short axis is the brachydome, i.e., the short



FIG. 29.—Model of brachydomes and macropincacoids.




FIG. 30.—Upper brachydome planes (011).

dome (Figs. 29 and 30). That one parallel to the long axis is called the macrodome, i.e., the long dome (Figs. 31 and 32). The domes do not produce closed figures unless united with each other or with some other planes. In Figures 29 and 31 they are closed by planes called pinacoids.

Prism planes (Fig. 33), like domes, are parallel to one axis; but it is always the  $c$  axis to which a prism is parallel. The symbol of the prism may be (110) or (210), etc.

3. The third kind of planes consists of those parallel to two axes and intercepting one axis. They are called pinacoids (from  $\pi\nu\alpha\kappa\omicron\varsigma$ , "plane") (Figs. 34 and 35).




FIG. 31.—Model of macrodomes and brachypinacoids.




FIG. 32.—Upper macrodomes (101)

The basal pinacoid, or base, is parallel to  $a$  and  $b$  and intercepts  $c$  (001).

The brachypinacoid (short pinacoid) is parallel to the  $c$  and to the shorter of the two lateral axes, the  $a$ , but intercepts the  $b$  (010). The macropinacoid (long pinacoid) is parallel to  $c$  and to the longer of the lateral axes, but intercepts the  $a$  (100).

Figure 36 shows a combination of prism (110), brachypinacoid (010), and brachydome (011).

Pyramids, domes, prisms, and pinacoids complete the list of holohedral forms in the orthorhombic system.

If the right-hand upper octant of a pyramid and each alternate octant were developed at the expense of their neighbors, a right-handed bisphenoid would be produced (Fig. 37). A left-handed or negative bisphenoid would result if the left-hand upper octant and alternate octants grew at the expense of their neighbors.




FIG. 33.—Prism (110) and basal (001) planes.




FIG. 34.—Base, macropinacoid, and brachypinacoid.




FIG. 35.—Model showing base, macro- and brachypinacoid.



FIG. 36.—Model of prism (110), brachypinacoid (010), and brachydome (011).




FIG. 37.—Right-handed sphenoid (111).

In sulphur these bisphenoids (111) are commonly united and modified by basal (001), dome (011), and obtuse pyramid planes (113) (Figs. 38 and 39).

In Figure 39 the left-handed sphenoid predominates while the right-handed appears as a very small plane. In Figure 38 they are of nearly equal size, but an edge instead of a corner where the  $a$  and  $b$  axes are intersected shows that the prevailing form is not a bipyramid but rather two bisphenoids.




FIG. 38.—Sulphur, usual habit




FIG. 39.—Sulphur, sphenoidal habit

Study of the sulphur crystal has shown that if the  $b$  axis is taken as unity, the  $a$  axis is .8 and the  $c$  is 1.9. Therefore to construct the various planes write parameters, ratios, and symbols as before.

		Parameters	Ratios	Symbols
Ordinary bipyramid	(Fig. 28)	.8:1:1.9	$\frac{.8}{1} : \frac{1}{1} : \frac{1.9}{1}$	(111)
Obtuse bipyramid	(Fig. 30)	2.4:3:1.9	$\frac{.8}{1} : \frac{1}{1} : \frac{1.9}{3}$	(113)
Brachydome bipyramid	(Fig. 32)	$\infty$ :1:1.9	$\frac{.8}{0} : \frac{1}{1} : \frac{1.9}{1}$	(011)
Macrodome bipyramid	(Fig. 34)	.8: $\infty$ :1.9	$\frac{.8}{1} : \frac{1}{0} : \frac{1.9}{1}$	(101)
Prism bipyramid	(Fig. 35)	.8:1: $\infty$	$\frac{.8}{1} : \frac{1}{1} : \frac{1.9}{0}$	(110)

Similarly for the pinacoids.

If melted sulphur is quickly cooled, the molecules do not have opportunity to arrange themselves and the resulting mass is without

definite form. It is said to be amorphous. If it is slowly cooled, crystals are formed similar to those occurring in nature but differing in this respect, that they slant downward parallel to the  $a$  axis, so that the front angle between the  $c$  and  $a$  axis,  $\beta$ , is greater than  $90^\circ$  (Fig. 40). The basal plane, being parallel to the lateral axes, slants forward. The crystals cannot be classed in the orthorhombic system but are in the monoclinic. (Monoclinic means having one inclination.) After a time, however, these monoclinic crystals become dull and fall to pieces, since their molecules tend to arrange themselves




FIG. 40.—Axes of a monoclinic crystal.

in the more stable form of the orthorhombic crystal. Orthorhombic crystals can be obtained artificially by allowing sulphur to crystallize from solution in carbon disulphide.

Sulphur cleaves very imperfectly, parallel to the base (001) and to the prism (110). It is brittle and shows conchoidal surfaces when broken. Hardness=2; gravity=2; luster, greasy, resinous, adamantine. It allows light to pass through imperfectly, i.e., it is translucent. Its average angle of refraction is  $n=2.04$ . Since the density of its molecules varies in different directions, a ray of entering light is divided into two rays. These rays vibrate at right angles to each other and are differently refracted. The dispersion or difference between the angle of greatest and least refraction is 0.29.

The heat conductivity is so low that the warmth of the hand is enough to cause a sulphur crystal to crackle, as may be noticed when a crystal is held near the ear. Sulphur becomes electric by friction; volatilizes easily, forming sulphur dioxide; is insoluble in acids.

Hundreds of thousands of tons are mined in Sicily annually. Spain, France, and Germany produce smaller amounts. Louisiana, Texas, Nevada, and Utah are the chief source of the domestic supply. In 1916 the first two states supplied 98 per cent of the sulphur obtained in the United States.

## SUMMARY

*Sulphur*.—S. Orthorhombic; symmetry holoaxial (sulphur class): (111), (113), (011), (101), (001).

Cleavage very imperfect (001), (110); brittle; fracture conchoidal.

Hardness=2; gravity=2. Yellow, orange, white; luster resinous; translucent; refraction strong,  $n=2.04$ ; double refraction very strong, positive.

Fusible; insoluble in acid; soluble in carbon disulphide,

Sicily, Spain, France, Germany, Louisiana, Texas.

## Arsenic

A small mass of native arsenic from Austria represents the usual appearance of this mineral. It somewhat resembles slag from a metal furnace or some kinds of lava, since it shows a rounded, twisted surface, like a bunch of grapes crowded together, and is dull lead gray or blackish on the surfaces which have long been exposed to the air. The fresh surfaces are tin white and show the short radiating needles which build up individual portions of the mass. It is brittle, less than 4 in hardness, and 5.7 in specific gravity.

Native arsenic furnishes but little of the arsenic used in medicine and the manufacturing arts.

## SUMMARY

*Arsenic*.—As. Hexagonal. Cleavage (0001); botryoidal, reniform, massive; brittle, conchoidal.

Hardness=3.5; gravity=5.6. Silver white; tarnishes lead gray to black; streak white.

Volatilizes without fusing, tinges flame blue, yields dense white fumes; odor of garlic.

Colorado, Chile, Saxony, Austria.

## Antimony and Bismuth

These two brittle metals are very similar in their occurrence, properties, and uses. They do not develop well-defined crystals, but are usually found in grains, incrustations, or aggregations of scales which form masses. Bismuth is sectile and is the softer of the two, being about 2 in the scale, while antimony is 3. Bismuth is the heavier of the two, having a specific gravity of 9, while that of antimony is 6. Bismuth is somewhat reddish in hue; antimony is tin



white. Both are metallic in luster, soluble in nitric acid, easily fusible and volatile. Both are found in association with silver, iron, arsenic, sulphur, and quartz. One hunting for these minerals should examine crystalline rocks.

The localities most noteworthy on account of specimens of antimony and bismuth are Saxony, Bohemia, and Japan. Many of the ores of precious metals in our western states contain these metals.

Antimony and bismuth are used in medicine and for the manufacture of alloys for type metal, babbitt metal, and other metals of low fusing-point.

#### SUMMARY

*Antimony*.—Sb. Hexagonal; symmetry dihexagonal alternating (calcite class). Cleavage parallel (0001) perfect, parallel  $\frac{1}{2}R$  fair; brittle; fracture uneven.

Hardness=3.5; gravity=6.6. Tin white; luster metallic; opaque.

Easily fusible, volatile; oxidizes in nitric acid.

Germany, France, Japan, Australia.

*Bismuth*.—Bi. Hexagonal; symmetry dihexagonal alternating (calcite class). Cleavage parallel (0001) perfect, parallel  $\frac{1}{2}R$  fair; sectile; fracture hackly.

Hardness=2; gravity=9. White with reddish tinge; luster metallic.

Easily fusible; volatilizes; soluble in nitric acid.

Germany, Bohemia, Colorado.

#### Gold

Probably there is more general interest in this mineral than in any other that is found in the earth's crust.

It was doubtless the first metal to be used by primitive man, since it is found in the beds of streams to which men would come for water and which were their highways from earliest times. Its glitter would attract the attention. When once its acquaintance was made, it would be easily recognized again, since it does not tarnish or rust, is very heavy, being 19 times as heavy as water, and so soft and malleable that it can be given various shapes and employed in many ways.

These qualities would lead men to use it long before they would notice or use the more abundant metals such as iron. It is found in the earliest tombs, such as those at Kertsch in the Crimea, in northern

Africa, and western Asia. Cloisonne work made in Egypt three or four thousand years ago shows skill in the use of gold.

The beauty of color, ease of working, weight and permanence of gold, render it a mineral of great value. But, however great its intrinsic worth, were it as common as quartz, for example, its value would be decreased.

Thus is it over all the earth  
That which we call the fairest  
And prize for its surpassing worth  
Is always rarest.

Iron is heaped in mountain piles  
And gluts the laggard forges,  
But gold flakes gleam in dim defiles  
And narrow gorges.

The snowy marble flecks the land  
In heaped and rounded ledges,  
While diamonds hide beneath the sand  
Their starry edges.\*

Gold is found usually in quartz veins, in pyrite and other sulphides, or in sands and gravels.

In quartz it occurs as fine threads or thicker wires that run singly or are bunched into mossy or treelike masses (arborescent). Sometimes it is in scales or grains isolated at times or packed together so as to form lenses or nuggets. Wiry and granular masses alike are rounded, twisted, and so distorted as to give little suggestion of crystal faces. However, an exposed end of one of these grains or threads, one which has had opportunity to develop in a cavity uncrowded by quartz or some other mineral, may show crystal faces clearly enough developed to permit of study and to make possible the




FIG. 41.—Cube

\* J. G. Holland, "Bitter Sweet."

conclusion that the structure of gold agrees with that of the diamond, the molecules being so arranged as to place it in the regular system.

Besides the octahedron (111), two other forms appear, namely, the cube (100) (Fig. 41) and the four-sided cube (tetrahexahedron, 210) (Figs. 42 and 43). To construct the cube, write the notation as before. Since the axes are interchangeable, six planes will be produced. The parameters of the front, right side, and top are as follows:

Parameters	Ratios	Symbols
1 : ∞ : ∞	$\frac{1}{1} : \frac{1}{0} : \frac{1}{0}$	(100)
∞ : 1 : ∞	$\frac{1}{0} : \frac{1}{1} : \frac{1}{0}$	(010)
∞ : ∞ : 1	$\frac{1}{1} : \frac{1}{0} : \frac{1}{1}$	(001)

A cube with four faces in each cubic face (Figs. 42 and 43) results when one of the three axes is intersected at twice unit's distance, for example.

Parameters	Ratios	Symbols
1 : 2 : ∞	$\frac{1}{2} : \frac{1}{1} : \frac{1}{0}$	(210)
2 : 1 : ∞	$\frac{1}{1} : \frac{1}{2} : \frac{1}{0}$	(120)
∞ : 1 : 2	$\frac{1}{0} : \frac{1}{2} : \frac{1}{1}$	(021)
∞ : 2 : 1	$\frac{1}{0} : \frac{1}{1} : \frac{1}{2}$	(012)
1 : ∞ : 2	$\frac{1}{2} : \frac{1}{0} : \frac{1}{1}$	(201)
2 : ∞ : 1	$\frac{1}{1} : \frac{1}{0} : \frac{1}{2}$	(102)

Gold crystals are usually small, distorted, and so grouped that to study and decipher them is a difficult matter. The gold contained in pyrites and metallic sulphides is so finely divided as to be invisible. It is mechanically included in the sulphides and not chemically united with the sulphur. In nearly every country in which gold is mined, it

was first discovered in sands and gravels, and such deposits until within the last fifty years have been the chief source of the metal.

Like diamonds, gold has been able to withstand the friction to which it was subjected while being washed from the original ledge. Diamonds resisted the friction because of their hardness; gold because of its tenacity; both have endured because of their insolubility and slight affinity for oxygen. The condition of gold in alluvial deposits varies from dust of microscopic fineness to nuggets many pounds in size. In California a nugget weighing 161 pounds was




FIG. 42.—Tetrahexahedron model




FIG. 43.—Construction of tetrahexahedron.

found. The largest nuggets have been discovered in Australia, three weighing over 200 pounds having been found there. The largest of them, the "Welcome," weighed 248 pounds.

The origin of nuggets of such size has been a matter of much speculation, since no masses of similar size have been found in veins. It has been suggested that small particles carried downstream were welded together by the impact of water-tossed gravel until a large nugget was formed, or that the nuggets have grown by accretion of gold from some percolating solution. Polished and etched surfaces of nuggets, however, show crystalline structure. This would be wanting in welded gold, and there is an absence of the onion-like structure that would be expected if the gold were deposited by accretion from solution. Hence it may be concluded that the nuggets

were originally in quartz veins and have been rounded in the downward journey from some high ledge to the resting-place in which they were discovered.

As to the origin of gold, nothing is known. The same may be said in regard to all elements. All that is known is something of the method of their transference and deposition. Light is shed on the subject by the fact that many fresh waters and all sea waters contain gold in appreciable quantities. There is nearly one grain of gold (five cents' worth) in every ton of ocean water. Then in all the oceans there is about seventy-five billion dollars' worth.

Gold in solution, possibly as a telluride, chloride, or cyanide, was carried by waters and deposited by them upon neutralization, cooling, or evaporation. Though the surfaces of gold crystals are rounded and often look as if melted, their appearance is not due to fusion but to their manner of crystallization. The origin of the gold is the same as the origin of the vein material inclosing it—quartz, fluorite, and calcite. All of these minerals are commonly deposited from aqueous solutions.

Gold melts at  $1200^{\circ}$  C. and forms such perfectly spherical globules that by microscopical measurements it is possible to estimate the amount of gold in a globule and hence to dispense with fine balances in assaying. Gold is soluble in aqua regia only (a combination of nitric [ $\text{HNO}_3$ ] and hydrochloric [ $\text{HCl}$ ] acids).

Sulphur and oxygen do not unite with gold, and hence it remains bright in nature or when worn as an ornament.

All gold contains silver in solid solution. As the amount of silver increases, the alloy becomes paler, lighter, and more liable to dissolve in nitric acid. Most Hungarian gold contains 30 per cent of silver, California gold 10 per cent, Australian gold (Mount Morgan, Queensland), reputed to be the purest, only .3 per cent.

Platinum, copper, and iron minerals, calcite, fluorite, quartz, feldspar, amphibole, and pyroxene, mica, garnet, and zircon, are the minerals most usually found with gold.

The rocks in which gold-bearing veins are found are igneous rocks such as granites, syenites, and porphyries; or metamorphic rocks such as gneisses and schists. The richest veins are usually at places of contact of different kinds of rock.

California, Nevada, Colorado, Montana, and South Dakota have been the chief producers of gold in this country since the discovery of

the metal in California in 1848. The only gold found in Illinois is an occasional piece contained in some rock transported from northern regions by the glaciers of Pleistocene times. There are no deposits of commercial importance. In spite of this fact, the procession of people who hope to discover such deposits or think they have done so will never end. They bring to the museum iron sulphide (pyrite), decaying mica (vermiculite), and other minerals, confident that they have found valuable deposits of precious metal; and when disillusioned are dejected. At one time the United States, at another time South Africa, leads the world in gold production, while Australia ranks third.

Gold is a metal useful in all places where hardness and toughness are not desired but where insolubility, permanence in the air, beauty of color, softness, and ductility are sought. Since the earliest times it has been used for personal adornment and for ornaments for the home, the church, and the palace. It is universally favored as a medium of exchange.

#### SUMMARY

*Gold.*—Au. Regular; holosymmetric; distorted (111), (100), (210); fibers, plates, grains. Malleable; ductile; fracture hackly.

Hardness = 2.5; gravity = 19.3. Gold yellow, metallic, opaque.

Fusible at 1200° C.; soluble in aqua regia.

Western North and South America, South Africa, Australia.

#### Silver

Silver resembles gold in its mode of occurrence, crystal habit, and physical properties. Chemically it is not so stable as gold, being readily affected by acid fumes and liquids. It is rarely found in placer deposits, but occurs most commonly in wiry, mossy, flaky, or granular forms in veins.

Sometimes large pure masses are discovered. One of the most famous was an eight-hundred-pound mass found in Peru. Another from Kongsberg, Norway, weighing five hundred pounds, is preserved in Copenhagen.

Crystals of silver are usually so distorted that their form is difficult to decipher, but under favorable circumstances octahedrons (111), cubes (100), and tetrahedrons (210) can be distinguished.

Like gold, silver has no direction of easy separation (cleavage). When broken, the fractured surfaces are splintery. It is inferior to

gold in its malleability and ductility, as it is possible only to beat leaves of it so thin that it requires one hundred thousand leaves to form a pile one inch in height, and one grain can be drawn out into four hundred feet of wire. Gold, however, can be beaten into leaves thin enough to require two hundred and eighty-two thousand leaves to form an inch-high pile, and one grain can be drawn into five hundred feet of wire.

Silver is unsurpassed as a conductor of electricity. Its conductivity is placed at 100 per cent, that of copper at 93 per cent, and platinum at 16 per cent.

One thousand degrees Centigrade of heat are required to melt it. When fused it can absorb twenty times its bulk of oxygen, which it gives off upon cooling, causing it to blossom into arborescent forms.

It is readily soluble in nitric acid. It unites with sulphur so easily that to keep silver bright is a very difficult task.

The chief source of the metal is not native silver but sulphides such as argentite, proustite, tetrahedrite, etc., minerals which will be described later.

The association, occurrence, and localities of silver are nearly identical with those of gold.

The United States has for many years been one of the principal producers, as well as the chief consumer, of silver.

It is estimated that the ocean contains over two million tons of silver worth more than \$38,000,000,000.

Silver is used extensively for coinage, for making household articles, for photographic purposes, and in various other ways.

#### SUMMARY

*Silver*.—Ag. Regular, holosymmetric; (100), (210), (111); twinned on (111). Threads, wires, plates, grains, masses. Malleable; ductile; fracture hackly.

Hardness = 2.5; gravity = 10.5. White, metallic, opaque.

Fusible at 1050° C.; soluble in nitric acid.

Cordilleran states in both North and South America, Australia, Germany.

#### Copper

Copper is similar in its physical characteristics, association, and occurrence to the two minerals just described, but is more abundant.





PLATE IV



Dendritic copper from Calumet and Hecla mining region, Michigan.

In the rare crystal faces discernible, possibly the tetrahexahedron (210) and dodecahedron (110) (Fig. 44) are more common than they are in gold and silver. The dodecahedron (110) can be constructed from the following parameter (Fig. 45):

Parameter	Ratios	Symbols
$1:1:\infty$	$\frac{1}{1}:\frac{1}{1}:\frac{1}{0}$	(110)

Wiry and arborescent forms are common. Masses of remarkable size have been found. One of the largest was 45 feet long and weighed 420 tons. It was found in the "Minnesota Mine" in Michigan. That region has produced more pure copper than any other in the world. The copper is disseminated in breccias, conglomerates, and basalts, or is collected in veins of calcite, fluorite, analcite, and quartz which penetrate the basalt. Often a cavity is filled partly with copper and partly with silver. If these metals had been deposited from a fused mass, they would have been united in an alloy rather than standing side by side. Evidently they were formed from a solution, and the more difficultly soluble silver was first deposited and later the copper.




FIG. 44.—Model of a dodecahedron

Copper is redder than gold and much more soluble. It is one of the most useful of metals, being used for electrical purposes and for many domestic and commercial articles.

The United States has produced about three times as much copper as the rest of the world together. Arizona, Montana, and Michigan are the leading states in production. In the two former the ores are chiefly sulphides and carbonates; in Michigan, native copper.

Glacial drift from the north has brought nuggets of copper, some of them weighing more than 50 pounds, and scattered them widely over Illinois. One (No. 695) found in the drift in Peoria County weighs  $18\frac{5}{8}$  pounds. A hole was cut through it by the finder so that




FIG. 45.—Construction of a dodecahedron

it could be used on a rope to close a gate. No. 693 from Macon County weighs  $17\frac{3}{8}$  pounds and is covered with the fine green deposit with which time paints old copper domes of churches and palaces. This deposit is formed when carbon, oxygen, and water unite with copper to produce the copper carbonate called malachite. No. 3383 is an irregular nugget ( $11\frac{1}{4}$  pounds) which shows the scratches made by the rocks over which the nugget was pushed while frozen in a glacier. No. 259, a small nugget from Jersey County, was found farthest south of any specimen in the collection.

#### SUMMARY

**Copper.**—Cu. Regular, holosymmetric; elongated (210); twinned on (111); threads, wires, masses. Malleable; ductile; fracture hackly.

Hardness=2.5; gravity=8.9. Copper red, metallic, opaque.

Fusible at  $1100^{\circ}$  C., soluble in nitric acid.

Michigan, Arizona, New Mexico.

#### Mercury

Mercury is the only element which is liquid at ordinary temperature. It solidifies at  $-40^{\circ}$  C. and in so doing crystallizes in the regular system. It unites so readily with sulphur that it is rarely found uncombined with that element. Cinnabar, HgS, is the chief source of the metal.

Mercury is used in making medicine, in "silvering" mirrors, and in the manufacture of toys, but chiefly as a means of collecting finely divided gold in placer mining and in the free milling process.

About the same time that gold was discovered in California, fortunately quicksilver was found at New Almaden, some fifty miles south of San Francisco.

The United States is at present the leading country in the production of cinnabar, from which mercury is obtained. The famous old Spanish localities now take second rank. Nature has not provided any deposits of mercury in Illinois. Nor do we need it as much as do some other states.

#### SUMMARY

*Mercury*.—Hg. Liquid, amorphous; at  $-40^{\circ}$  C. Regular.  
Gravity = 15. White, metallic, opaque.  
Volatilizes, sublimes.  
California, Spain.

#### Platinum

Platinum is a steel-gray, metallic, moderately hard, exceedingly heavy mineral occurring in small flat grains in alluvial deposits. The world's supply has been obtained practically from the Ural Mountains alone. If its appearance and characteristics were more widely known among prospectors, other localities might be added to the list of producers.

Because of its peculiar utility and rarity, platinum is at present unsurpassed in commercial value by any metal. Its especial usefulness depends upon its resistance to heat. Over  $2000^{\circ}$  C. are required to melt it. This, in addition to its insolubility, makes it serviceable for dental purposes, for crucibles, wire, and foil to be used in chemical laboratories and manufacturing plants and for electrical purposes. The attempt to find some metal which will take the place of platinum has been unsuccessful.

The United States uses about half of all the platinum produced in the whole world.

The crystal form of platinum, being similar to that of gold, silver, and copper, presents nothing new for consideration.

Platinum is very finely disseminated in gravels derived from serpentine and syenite, and large placers may be expected only in very

old land areas which have been subjected to protracted degradation. The Ural Mountains furnish such conditions.

Small percentages of platinum are often obtained from sulphides of antimony, arsenic and copper, and in chromite. The placers of the Ural Mountains, Columbia, and California contain, associated with platinum, other minerals of high specific gravity such as gold, cassiterite, magnetite, hematite, chromite, and rutile.

#### SUMMARY

*Platinum*.—Pt. Regular; holosymmetric; grains and nuggets. Malleable; ductile; fracture hackly.

Hardness = 4.5; gravity = 19; chemically pure, 21. Steel gray.

Infusible; soluble in nitro-hydrochloric acid (aqua regia).

Nijni Taguilsk (Urals); Columbia, South Africa, Canada, Wyoming, California.

#### Iron

Iron so readily unites with oxygen, sulphur, and other elements that it rarely occurs native. Consequently, while minerals containing iron are numerous and abundant, pure iron is rare. Yet it is one of the most interesting of minerals because of its origin. Some of it is terrestrial and some meteoric in origin.

Terrestrial iron is found as small imbedded particles in basalt, peridotite, and serpentine—three kinds of dark rocks abundant in many mountain regions—and in deposits derived from the disintegration of these rocks. Gold and platinum are usually associated with terrestrial iron.

At several places on the west coast of Greenland, especially at Disco Island, large masses of iron occur which are regarded as originating from deep-seated portions of the earth, since the basalts of the region contain scattered grains and globules of iron.

Meteoritic iron illustrates the fact that the science of mineralogy is concerned not only with this earth but with the universe as well.

Until within the last one hundred years the idea prevailed that meteorites were portions of this earth which had been thrown out of volcanoes with such velocity as to reach great heights and then to fall back with enormous speed. But as the composition of meteorites became known and the circumstances connected with their fall were investigated, students of the subject were convinced that they are fragments of other heavenly bodies—the dust of the universe.



PLATE V



**Mukerop meteorite, one-sixth natural size. Fell in Amalia-Goamus, West Africa. Section mentioned on page 44 was cut from the center of this mass.**

Myriads of them enter the earth's atmosphere. At night they are seen to flash in the heavens when they are ignited by the friction generated in their fall through the earth's atmosphere. Many enter the atmosphere at such an angle that they leave it without touching the earth; many are totally consumed as they fall; some reach the earth's surface as cosmic dust, as grains, or even as masses many tons in weight. In 1894 at Cape York, in the northern part of Greenland, a mass weighing 36 tons was found and three years later brought by R. E. Peary to New York City. It was called by the Eskimos "Ahnighito" or "The Tent."

Occasionally a meteorite has been seen as it fell and has been picked up while still warm. Those which have been observed in the air and then found are called "falls." Their number is less than the so-called "finds," which are not seen to fall but are simply picked up. Several hundred "falls" and "finds" have been collected and described.

A meteorite entering the atmosphere may have an astonishingly high velocity—something like 45 miles a second—but because of the resistance of the air be reduced in velocity and strike the earth's surface with small force. Meteoric stones fell upon the ice at Hesse, Sweden, and rebounded without either breaking the ice or being themselves shattered. The heat generated by the friction with the air fuses the surface of the meteorite, especially on the front side, and causes the melted material to flow back in waves, making a kind of "varnish." Meteorites are usually pitted with thumblike impressions. Since the heating is sudden, the surface may be fused while the interior is still cold. The unequal expansion causes them to explode with loud report and to scatter over wide territory.

According to constitution there are three kinds of meteorites: first, those consisting almost wholly of iron (siderites); second, those having a cellular matrix of iron in which stony matter is imbedded (siderolites); and third, those composed almost entirely or wholly of stony matter (aerolites).

Meteoric iron is massive, but its crystalline structure can be readily discerned when it is etched with diluted nitric acid, since triangular markings usually appear on the surface. They are due to the presence of nickel. The form and the widths of the bands depend upon the percentages of nickel present. The figures resulting



are called Widmanstätten figures, after the man who first studied them.

The largest meteorite ever discovered in the United States and one of the most interesting is the Willamette iron. It was found 19 miles south of Portland, Oregon, in 1902. It weighs 15 tons and is now in the American Museum of Natural History.

Meteorites have been found in our neighboring states, Michigan, Indiana, Kentucky, Missouri, Iowa, and Wisconsin, but thus far not a single example has been reported in Illinois. All accounts of the finding of meteorites in this state have upon investigation proved to be untrue. There appears to be no reason why falls may not occur here at any time. If people are more observant, we may some time discover and preserve these messengers from the great waste spaces.

The largest meteorite exhibited in the collection (No. 4064) is a  $\frac{7}{8}$ -inch-thick section from 13 to 15 inches in diameter and weighing 13 $\frac{5}{8}$  pounds avoirdupois. It was cut from the Mukerop meteorite which fell in southwestern Africa (Plate V). The following also are shown: a dozen examples of the Canon Diablo, Arizona, meteorite (No. 3385); about fifty of the Holbrook, Arizona; one from Eddy County, New Mexico (Sacramento Mountains, No. 3384); Sheridan County, Kansas (Saline, No. 4106); Lyon County, Kansas (Admire, No. 4104); Phillips County, Kansas (Long Island, No. 4109); Iowa County, Iowa (Homestead, No. 4107, and Forest, No. 4108); Emmet County, Iowa (No. 1730); Bullitt County, Kentucky (Salt River, No. 4103); Kent County, Michigan (Grand Rapids, No. 4103); and state of Mexico, Mexico (Toluca, No. 4101).

#### SUMMARY

*Iron*.—Fe. Nickel usually present. Regular; (111), (100); massive lamellar; cleavage parallel (100) perfect; malleable; fracture hackly.

Hardness=4.5; gravity=7.5. Gray to black, metallic, magnetic Infusible; soluble in acid.

Greenland, and in meteorites of wide distribution.

## LIST OF ELEMENTS AND THEIR ATOMIC WEIGHTS

Name	Combining Weight Oxygen = 16	Name	Combining Weight Oxygen = 16
Aluminium, Al.....	27	Mercury, Hg.....	200
Antimony, Sb.....	120	Molybdenum, Mo.....	96
Argon, Ar.....	39	Neodymium, Nd.....	144
Arsenic, As.....	75	Neon, Ne.....	20
Barium, Ba.....	137	Nickel, Ni.....	59
Beryllium, Be.....	9	Nitrogen, N.....	14
Bismuth, Bi.....	208	Osmium, Os.....	191
Boron, B.....	11	Oxygen, O.....	16
Bromine, Br.....	79	Palladium, Pd.....	107
Cadmium, Cd.....	112	Phosphorus, P.....	31
Caesium, Cs.....	132	Platinum, Pt.....	195
Calcium, Ca.....	40	Potassium, K.....	39
Carbon, C.....	12	Praeseodymium.....	141
Cerium, Ce.....	140	Radium, Ra.....	226
Chlorine, Cl.....	35	Rhodium, Rh.....	103
Chromium, Cr.....	52	Rubidium, Rb.....	85
Cobalt, Co.....	59	Ruthenium, Ru.....	102
Columbium, Cb.....	94	Samarium, Sm.....	150
Copper, Cu.....	63	Scandium, Sc.....	44
Dysprosium, Dy.....	162	Selenium, Se.....	79
Erbium, Er.....	167	Silicon, Si.....	28
Europium, Eu.....	152	Silver, Ag.....	108
Fluorine, F.....	19	Sodium, Na.....	23
Gadolinium, Gd.....	157	Strontium, Sr.....	88
Gallium, Ga.....	70	Sulphur, S.....	32
Germanium, Ge.....	72	Tantalum, Ta.....	181
Glucinum, Gl.....	9	Tellurium, Te.....	127
Gold, Au.....	197	Terbium, Tb.....	159
Helium, He.....	4	Thallium, Tl.....	204
Hydrogen, H.....	1	Thorium, Th.....	232
Indium, In.....	114	Thulium, Tu.....	168
Iodine, I.....	127	Tin, Sn.....	119
Iridium, Ir.....	193	Titanium, Ti.....	48
Iron, Fe.....	56	Tungsten, W.....	184
Krypton, Kr.....	83	Uranium, U.....	239
Lanthanum, La.....	139	Vanadium, V.....	51
Lead, Pb.....	207	Xenon, Xe.....	131
Lithium, Li.....	7	Ytterbium, Yt.....	172
Lutecium, Lu.....	174	Yttrium, Y.....	89
Magnesium, Mg.....	24	Zinc, Zn.....	65
Manganese, Mn.....	55	Zirconium, Zr.....	91

## CLASS II. SULPHIDES

The next group of minerals which would naturally claim the attention of the visitor is that which embraces minerals consisting of a mixture of sulphur with some metal like antimony, molybdenum, lead, silver, copper, zinc, mercury, or iron.

From the twenty-five or more minerals in the group, thirteen are common; and while but eight of them are found in Illinois, all are used here and all are of interest, since they show marked properties.

They are stibnite, molybdenite, galena, argentite, chalcocite, sphalerite, cinnabar, pyrrhotite, erubescite, chalcopyrite, pyrite, marcasite, and arsenopyrite.

### Stibnite

Stibnite, a sulphide of antimony ( $\text{Sb}_2\text{S}_3$ ) is the chief source of the metal, antimony. Its crystals are often large and beautiful. They resemble sulphur crystals since their structure is different in three directions. The planes which are usually developed are prisms and pinacoids. Several pyramid planes are of common occurrence. The crystals are holohedral. Basal planes are wanting. The long needle-like crystals often terminate in a flat pyramid (113), as shown in Figure 46. The ratio of the axes  $a:b:c$  is .99:1:1.01, differing thus but slightly from a mineral in the regular system.


Applying these values and using the parameter as usual, the following result:

Parameters	Ratios	Symbols
2.97:3:1.01	$\frac{.99}{1} : \frac{1}{1} : \frac{1.01}{3}$	(113)
.99:1:∞	$\frac{.99}{1} : \frac{1}{1} : \frac{1.01}{0}$	(110)
∞:1:∞	$\frac{.99}{0} : \frac{1}{1} : \frac{1.01}{0}$	(010)

Some crystals of stibnite are of remarkable size and beauty. One of the finest specimens in any museum may be seen in the British Museum. It is a group of crystals eighteen inches long and terminated by lustrous pyramid faces. It came from the antimony mines



PLATE VI



*a*, Stibnite, Japan



*b*, Molybdenite from Aldfield, Pontiac County,  
Quebec, Canada.

at Shikoku, Japan, a locality which has furnished a larger number of remarkable specimens than any place in the world. Our largest specimen, No. 3784 (Plate VI *a*), comes from the same place. Other samples are from Portugal, Australia, and the western United States. Stibnite crystals are often twisted, curved, and warped. The most usual occurrence is that of massive forms with bladelike or fibrous structure.

The mineral cleaves easily parallel to the brachypinacoid, and shows nicks and horizontal lines at right angles to the  $c$  axis, indicating "glide planes" parallel to the base (001). These glide planes make it possible for the crystals to bend, and explain their curved and twisted form.

That the mineral is in the orthorhombic system can easily be illustrated by the difference in rapidity with which heat is transmitted in different directions. Senarmont's method is to coat a brachypinacoid plane with wax and touch it with the point of a hot wire. The wax is melted more rapidly in the direction of the  $c$  axis than in the direction of  $a$ . Consequently the resulting figure is an ellipse. Roentgen's method, similar in principle, is to breathe upon a face, touch it with the point of a hot wire, then sprinkle lycopodium powder upon it. When shaken, the powder drops from the mineral where it was dry. The form of the clean space is an ellipse with the long axis parallel to the  $c$  axis.




FIG. 46.—Stibnite crystal.

Stibnite is found with other sulphides (argentite, galena, sphalerite, cinnabar) and with barite and quartz in veins in granite and gneiss.

The ancients used stibnite as a pigment to darken eyebrows. Its chief use at present is as a source of antimony.

#### SUMMARY

*Stibnite*.— $\text{Sb}_2\text{S}_3$ ;  $\text{Sb}=71.8$  per cent,  $\text{S}=28.2$  per cent. Orthorhombic; (110), (111), (113), (010). Massive, bladed, fibrous, granular. Cleavage (010) perfect; glide planes (001); slightly pliable; fracture conchoidal.

Hardness=2; gravity=4.6. Steel gray; metallic; opaque.

Easily fusible (1 in the scale); volatilizes; soluble in hydrochloric acid. Japan, Hungary, Australia, California.

### Molybdenite

Molybdenite, the sulphide of molybdenum ( $\text{MoS}_2$ ), is a soft metallic mineral, bluish lead gray in color. It occurs in six-sided (hexagonal) tabular crystals in quartz veins (Plate VI *b*). In softness, color, and form it closely resembles graphite but can be distinguished by the fact that the color is bluish and the mark left on paper (the "streak") is bluish, while the color and streak of graphite are lead gray. Molybdenite (gravity=4.7) is also more than twice as heavy as graphite.

Its crystals are often striated horizontally, taper toward the top because of the decrease in the diameter of its constituent lamellae, and show glide planes. Foliated, scaly, and granular particles sometimes are scattered through the containing quartz and at other times concentrated in the masses. With it are often found other sulphides such as pyrite and chalcopyrite. It has been deposited from solution in crystalline rocks, such as pegmatite granite, gneiss, and granular limestone.

The chief sources of supply in the United States recently have been California, Colorado, Montana, Maine, and Washington. None is found in Illinois.

Molybdenum compounds are used in coloring silk, leather, and porcelain blue. They have a limited use in chemical laboratories for the determination of phosphorus; in the manufacture of steel a fraction of a per cent of molybdenum hardens the steel and changes its coefficient of expansion.

### SUMMARY

*Molybdenite*.— $\text{MoS}_2$ ; Mo=60 per cent, S=40 per cent. Hexagonal; plates, scales; cleavage parallel (0001). Flexible; sectile.

Hardness=1; gravity=4.7. Bluish gray; metallic; opaque; greasy. Infusible; soluble in nitric acid.

California, Colorado, Montana, Washington, Maine, Canada, and many European localities.

### Galena

Because of its physical properties and its importance commercially, galena, the sulphide of lead ( $\text{PbS}$ ), is an interesting mineral. It is found in great masses or disseminated in limestone, as in the Mississippi Valley region, and in veins in crystalline rock, as in the

Cordilleran region. In the Cordilleras the galena is usually argentiferous and consequently one of the chief sources of silver in this country. In the Mississippi Valley region it contains practically no silver but is associated with the zinc sulphide, sphalerite.

Galena is mined in many places in both hemispheres, but probably in no place more extensively than in Missouri and Idaho. In early days in the Mississippi Valley region the avocation of the farmers was often the quarrying of galena for lead from which to cast bullets in time of war and for making pewter ware in time of peace. Some galena is found in Pope and Hardin counties in Illinois in connection




FIG. 47.—Cube truncated by octahedron.




FIG. 48.—Model of a cube truncated by an octahedron.

with the fluorite mined there, and some is produced in the north-western portion of the state. Scattered crystals may be detected in the "Niagara" limestone at different places. The finest samples shown in the museum collection are from Jo Daviess County. No. 421 is a cube whose corners are truncated with octahedron planes. It measures over three inches each way and weighs  $7\frac{1}{8}$  pounds (Fig. 49). No. 3396 is another smaller cube, and No. 267 is a large mass which has been incrustated with iron sulphide (marcasite). Where the incrustation has broken off, the underlying galena may be seen. When one sees this pronounced crystallization he is impressed with the fact that when minerals have the opportunity they have a form as well defined as that of a flower.



Galena is soft, heavy, lead gray, metallic, and opaque. It crystallizes readily, so that even massive forms when cleaved show the structure, and well-formed crystals are very common. The usual habit is fine cubes with the corners truncated by octahedrons (Figs. 47 and 48). The octahedrons may be enlarged so as to almost displace the cube, or they may become so small as to disappear.

The cube faces are often formed by very flat four-faced cube planes ( $hko$ ),  $h$  and  $k$  representing any two different numbers. If  $h=4$  and  $k=1$ , the symbol is ( $410$ ), a tetrahedron often met with.



FIG. 49.—Galena, Jo Daviess County, Illinois.




FIG. 50.—Model of planes appearing on galena.

Crystals often exhibit the dodecahedron ( $110$ ) in combination with the cube and octahedron (Fig. 50).

Cleavage is so perfect that a single blow of a hammer will shatter a crystal into multitudes of little cubes whose faces may show striations parallel to the lower right-hand trisoctahedron ( $441$ ) due to twinning lamellae parallel to that plane (Fig. 51). Since glide planes can be produced in this direction by pressure, the striae may be due to that cause.

Galena which contains from 1 to 2 per cent of bismuth has octahedral cleavage. When heated enough to drive off the bismuth, the cleavage becomes cubic. Singularly, galena containing bismuth does not decrepitate when heated, as does ordinary galena, nor is

there a change in its specific gravity. Further, with the change in crystalline structure, there is no decrepitation such as occurs in ordinary galena.

Galena usually contains small amounts of silver sulphide, and as the amount present increases, the galena loses its coarse cubic structure and becomes finely granular.

When covered with a layer of wax and touched with the point of a hot wire, the wax melts in a circle, showing that galena is in the regular system.

Argentite, sphalerite, chalcopyrite, pyrite, fluorite, quartz, calcite, and rhodochrosite accompany galena in limestones or in crystalline rocks.

Since galena furnishes the lead of the world, it is one of the most useful of minerals.

Lead is used in plumbing, in the manufacturing of paint, medicine, alloys, shot, etc.




FIG. 51.—Twin lamellae in galena parallel to  $(44\bar{1})$ .

#### SUMMARY

*Galena*.— $\text{PbS}$ ;  $\text{Pb}$  = 86.6 per cent,  $\text{S}$  = 13.4 per cent. Regular; holosymmetric; (100), (111), (110), (221). Cleavage (100) perfect; fracture even; nearly sectile.

Hardness = 2.5; gravity = 7.5. Lead gray; metallic; opaque.

Easily fusible, decrepitates; soluble in nitric acid.

Mississippi Valley, Cordilleran region.

#### Argentite

Argentite, the sulphide of silver ( $\text{Ag}_2\text{S}$ ), is one of the chief sources of the metal. It closely resembles galena but does not occur in such distinct crystals. Usually it is in the form of dendritic, scaly, earthy, or granular masses. It does not cleave as readily as galena but is sectile. It is associated with the same minerals and is found in crystalline rocks such as are wont to contain gold, silver, and other precious minerals. The best crystal shown, No. 3822, was obtained at Guanajuato, Mexico.

## SUMMARY

*Argentite*.— $\text{Ag}_2\text{S}$ ;  $\text{Ag}=87.1$  per cent,  $\text{S}=12.9$  per cent. Regular, holosymmetric; (100), (110). Cleaves imperfectly parallel (100), (110); sectile; fracture sub-conchoidal.

Hardness=2.5; gravity=7.3. Color and streak lead gray; metallic; opaque.

Melts readily; soluble in nitric acid.

In the mountain ranges in western North and South America, and in many European and Australian localities.

## Chalcocite

In Arizona during 1918 nearly as much copper was produced as was obtained from Michigan, Montana, and Utah combined. The ore consists chiefly of chalcocite ( $\text{Cu}_2\text{S}$ ), a mineral which is dark lead gray in color, metallic, and opaque, and occurs in granular or compact masses. It resembles argentite in general appearance, but is more brittle and is often tarnished blue or green when the addition of sulphur changes the chalcocite ( $\text{Cu}_2\text{S}$ ) to covellite ( $\text{CuS}$ ), or the addition of iron changes it into erubescite ( $\text{Cu}_3\text{FeS}_3$ ). While chalcocite crystallizes in the orthorhombic system, well-formed crystals are rare. Since the angle between the prism planes (110) is  $60^\circ$ , chalcocite often looks as if it were a hexagonal mineral. When several crystals are twinned about the prism planes, the form is even more deceptive.

Chalcocite is found in connection with other sulphides at many localities in the Cordilleran range. No example has been reported in Illinois.

## SUMMARY

*Chalcocite*.— $\text{Cu}_2\text{S}$ ;  $\text{Cu}=79.8$  per cent,  $\text{S}=20.2$  per cent. Orthorhombic;  $a:b:c=0.58:1:0.97$ . Common planes (110), (001), (023), (113); twinned on (110), (032); cleavage imperfect (110); sectile; fracture conchoidal.

Hardness=2.5; gravity=5.7. Lead gray; streak black; metallic; opaque.

Easily fusible; soluble in nitric acid.

Cordilleran region, England, Germany.

## Sphalerite

Many localities in which lead is abundant are also famous because of their great deposits of sphalerite ( $\text{ZnS}$ ). The early German miners who were seeking lead were disappointed when they found sphalerite

instead, and therefore called it *Blende* from *blenden*, "to deceive." "Sphalerite," from the Greek, has the same meaning.

Pure sphalerite has the color of resin. See the specimen from Spain (No. 3765). Usually it is dark because of impurities like iron, cadmium, manganese, tin, thallium, indium, and gallium that are often present in varying quantities. Some sphalerite contains as much as 20 per cent of iron. Miners call the dark varieties "Black Jack." See specimens from Colorado, Kansas, and Missouri. Gallium and indium were first discovered in sphalerite. Sphalerite occurs, as do most of the other sulphides, when igneous rocks such as granites, diabases, and porphyries are in contact with metamorphic rocks such as gneisses, schists, and granular limestones, especially where these rocks have been fissured and subsequently cemented by vein-forming materials. In the Mississippi Valley region, however, sphalerite usually is found in beds or is scattered through the limestone.

Well-crystallized specimens are seen to follow the laws of the regular system and to illustrate the same class of symmetry as that which is shown by the diamond, the "tetrahedrite class."

Common forms such as that in Figure 52 are combinations of tetrahedrons (111), dodecahedrons (110), and trapezohedrons (311). The tetrahedrons are positive and negative, and upon the alternate octants only occur the planes which together would produce the hemihedral form called the three-faced tetrahedron (311) (Figs. 53 and 54). Supplementary tetrahedrons combined with cubes are characteristic (Fig. 55). The positive and negative tetrahedrons may be distinguished by the difference in their size, by their differing smoothness, by the different markings which their faces show when they are etched with dilute hydrochloric acid, and by a pyro-electric test. To make this test cut a plate parallel to a face of each of the two tetrahedrons in turn. Insulate, connect with an electroscope, and touch with the point of a heated wire. One tetrahedron will become positively electrified, and the other negatively.




FIG. 52.—Sphalerite

Twin lamellae parallel to tetrahedral faces are common in the sphalerite of many localities.

Stibnite, galena, argentite, pyrite, marcasite, chalcopyrite, fluorite, quartz, calcite, and barite are the associates of sphalerite.




FIG. 53.—Model of a three-faced tetrahedron, a tristetrahedron.




FIG. 54.—Construction of trigonal tristetrahedron.



FIG. 55.—Sphalerite

The region around Joplin, Missouri, has produced probably more sphalerite than has any other locality in the world, and Illinois is the leading state in zinc smelting from these ores. The museum collection contains also specimens from Alston Moor, England, and Kapnik, Hungary, two places famous for their many fine crystals.

Sphalerite is the most important source of zinc, and the metal obtained from it is used in galvanizing iron, in zinc plating, in paint manufacture, and in medicine.

#### SUMMARY

*Sphalerite*.— $\text{ZnS}$ ;  $\text{Zn}=67$  per cent,  $\text{S}=35$  per cent. Regular, tetrahedrite class;  $(111)$ ,  $(110)$ ,  $(311)$ ,  $(100)$ . Cleavage perfect parallel  $(110)$ ; brittle; fracture conchoidal.

Hardness = 3.5; gravity = 4. Yellow, adamantine, translucent; refraction  $n=2.37$ .

Fusible with difficulty; soluble in hydrochloric acid.

Kansas, Missouri, Illinois, Wisconsin, Colorado, Utah, Montana, Europe.

**Cinnabar**

Though mercury occurs sometimes uncombined in nature, the chief source of the metal is cinnabar ( $\text{HgS}$ ). Cinnabar, a word used in India two thousand years ago, means "red resin" and is well applied, since the color of the mineral is bright red and the streak vermilion. See specimens No. 3401 and No. 3893. Impurities make it brown or slaty (No. 593). Crystals of cinnabar are rare. The mineral is notable for its refractive power, the ordinary ray ( $\omega$ ) being more strongly refracted than it is in diamond.  $\omega = 2.85$ .

Further, a ray of light entering the crystal in almost any direction is divided into two rays which vibrate at right angles to each other. That is, it is "doubly refracted." One ray is called the ordinary ( $\omega$ ) and the other the extraordinary ( $\epsilon$ ). When the difference between them is great, the double refraction or "birefringence" is said to be strong. In cinnabar  $\epsilon - \omega = 0.35$ .

Of late years more cinnabar has been produced in the United States than in any other country, and of this production the greater part is furnished by California. None is found in Illinois.

**SUMMARY**

*Cinnabar*.— $\text{HgS}$ ;  $\text{Hg} = 86.2$  per cent,  $\text{S} = 13.8$  per cent. Hexagonal; "quartz class": (1010), (0001), rhombohedrons (1011);  $c = 1.145$ . Cleavage good, parallel (1010); fracture uneven.

Hardness = 2.5; gravity = 8.2. Cochineal red; streak vermilion; luster, metallic, adamantine; translucent. Refraction very strong,  $\omega = 2.85$ ; birefringence, positive, very strong ( $\epsilon - \omega = 0.35$ ). Circular polarization very strong.

Volatile; soluble in nitric acid.

New Almaden, California; Spain; and south Russia.

**Pyrrhotite**

Pyrrhotite (*πυρρός*, "reddish") is a bronze-colored, magnetic iron sulphide, which occurs in massive forms and is often lamellar in structure.

Its crystallization is so imperfect as to leave doubt concerning its true nature, and, being opaque, its optical properties can shed no light on the question. However, its structure is probably such as characterizes the hexagonal system.

There is also doubt as to the chemical composition of pyrrhotite. Different formulae have been given to it, ranging from  $\text{Fe}_8\text{S}_7$  to  $\text{Fe}_{11}\text{S}_{12}$ . The formulae all agree closely with the monosulphide  $\text{FeS}$ , troilite, which is a mineral not known on the earth but common in some meteorites.

Pyrrhotite is not so abundant as other iron sulphides. The iron which it contains cannot be separated from the sulphur without great difficulty. However, in some localities, as at Ducktown, Tennessee, immense quantities of sulphuric acid are made from it. Nickel and cobalt are often present in paying quantities, and the nickeliferous pyrrhotite of Pennsylvania, Canada, and Norway is an important source of those metals.

#### SUMMARY

*Pyrrhotite*.— $\text{Fe}_{11}\text{S}_{12}$ ; Fe=56 to 61 per cent; S=44 to 39 per cent. Hexagonal plates, masses. Brittle; fracture uneven.

Hardness=4; gravity=4.6. Bronze yellow; streak grayish black; metallic; opaque; magnetic.

Fusible; soluble in nitric acid.

Appalachian and Cordilleran systems; Europe.

#### Erubescite

Erubescite, the "blushing ore" ( $\text{Cu}_3\text{FeS}_3$ ), owes its beauty to the ease with which it tarnishes. It is called also bornite, variegated copper, horseflesh ore, peacock ore. When freshly broken it has a coppery or bronzy color, but soon tarnishes to a vivid blue or purple. Its color is its most interesting characteristic.

Granular or compact masses are the most usual, but sometimes crystals in cubes can be distinguished. As is always the case, the crystals represent the purest condition.

Cornwall, England, South Africa, and some of the Cordilleran states furnish the best crystals and the most abundant supply of erubescite. Our best samples were obtained in Colorado (No. 3753) and New Mexico (No. 2195).

#### SUMMARY

*Erubescite*.— $\text{Cu}_3\text{FeS}_3$ ; Cu=55.5 per cent, Fe=16.4 per cent, S=28.1 per cent. Regular; (100); twinned on (111); cleavage imperfect, parallel (111); slightly sectile; fracture sub-conchoidal.

Hardness=3; gravity=5. Pinchbeck brown, bronze, tarnished blue; streak grayish black; metallic; opaque.

Fusible; soluble in nitric acid.

With other copper ores in Colorado, Montana, South Africa.

### Chalcopyrite

Very closely related to erubescite in chemical composition but much more pronounced in physical characteristics and commercial importance is chalcopyrite ( $\text{CuFeS}_2$ ), i.e., copper pyrite, a name given by Henckel in 1725 when a difference between this and pyrite was




FIG. 56.—Acute primary bipyramid




FIG. 57.—Obtuse primary bipyramid

for the first time noticed. Chalcopyrite and chalcocite are the chief sources of copper today.

Lustrous, clean-cut crystals of chalcopyrite are common, and were early studied by crystallographers who thought they were in the regular system until, in 1822, accurate measurements showed that the  $c$  axis is 0.985 when  $a$  and  $b$  are unity. Hence the crystals are in the Tetragonal System, that system in which the  $c$  axis is longer or shorter than the  $a$  and  $b$ , the  $a$  and  $b$  axes are equal, and all three axes are at right angles. The symbol (111) indicates a bipyramid which is acute when the  $c$  axis is longer than the lateral axes (Fig. 56), or obtuse when the  $c$  axis is shorter than the others (Fig. 57).

Since the lateral axes are equal and interchangeable, a form whose symbol is (101) will be a secondary bipyramid, instead of one consisting of dome planes, as in the orthorhombic system (Fig. 58).



Symbols such as  $(211)$  or  $(331)$ , etc., indicate the ditetragonal bipyramid (Figs. 59 and 60), since the two or three can be applied to each lateral axis in turn, thus indicating two planes in each octant.

Similarly there are three prisms: a primary,  $(110)$  (Fig. 61); a secondary turned  $45^\circ$  to it,  $(100)$  (Fig. 62); and a ditetragonal prism,  $(210)$  (Fig. 63). These with the basal plane represent the simple holohedral forms of the system. Figure 64 shows a combination of several of these planes.

Hemihedral forms are constructed on the same plan as were those in the systems heretofore described. For example, when the alternate pyramid faces only are developed, a bisphenoid results (Fig. 65). It may




FIG. 58.—Model of a secondary bipyramid.




FIG. 59.—Ditetragonal bipyramid




FIG. 60.—Model of a ditetragonal bipyramid.

be either positive or negative. If the planes in alternate octants of a ditetragonal pyramid are developed, the tetragonal scalenohedron is produced. Tetragonal scalenohedrons possess two planes

of symmetry intersecting at right angles in the  $c$  axis, which is an axis of alternating symmetry. They are so well represented




FIG. 61.—Primary prism




FIG. 62.—Secondary prism





FIG. 63.—Model of a ditetragonal prism.

FIG. 64.—Combination of primary prism ( $110$ ), secondary prism ( $100$ ), secondary bipyramid ( $101$ ), and ditetragonal bipyramid ( $211$ ).

in chalcopyrite that the class has been called the “chalcopyrite class of symmetry.”

Figure 66 represents the most usual chalcopyrite crystal. It is composed of the positive scalenohedron. Pyramid planes with the

symbol ( $201$ ) often appear on the edges. The scalenohedrons may be either acute or obtuse.

Figure 67 represents a form composed of the prism ( $110$ ), scalenohedrons ( $111$ ) and ( $101$ ), and the basal plane ( $001$ ).

Two kinds of twins are common. In one the twinning plane is ( $111$ ) and produces a form so similar to the twin characteristic of the mineral spinel as to be called the "spinel twin." (The "spinel twin" proper is a form in the regular system.)

The faces of one of the scalenohedrons are bright, while those of the other are dull.

In the second form of twinning, a central crystal ( $201$ ) is surrounded by four other crystals which are joined on the primary pyramid plane ( $111$ ),



FIG. 65.—Model of a bisphenoid




FIG. 66.—Chalcopyrite




FIG. 67.—Chalcopyrite, French Creek, Pennsylvania.

forming a composite twin (Fig. 68). As is generally the case with all minerals, massive forms are the rule and evident crystals the exception.

The color of chalcopyrite is bronze yellow and its streak is greenish black. Because of surface alterations it readily tarnishes, and takes



PLATE VII



*a*, Group of pyrite cubes, showing striations, Central City, Colorado.



*b*, Pyrite. A pyritohedron and cubes, Colorado

on beautiful iridescent colors. The vivid blue is due to the formation of covellite ( $\text{CuS}$ ).

The Cordilleran region from Arizona to Montana furnishes large quantities of chalcopyrite. Many fine crystals have been found at French Creek, Pennsylvania, the Hartz Mountains, and Cornwall.

#### SUMMARY

*Chalcopyrite*.— $\text{CuFeS}_2$ ;  $\text{Cu}$  = 34.5 per cent,  $\text{Fe}$  = 30.5 per cent,  $\text{S}$  = 35 per cent. Tetragonal; symmetry ditetragonal alternating (chalcopyrite class);  $a:c = 1:0.985$ . Common forms (111), (101), (211), (001), (201), (114), (441); twinned about normal of (111). Brittle; fracture conchoidal.

Hardness = 4; gravity = 4.2. Brass yellow, tarnishes blue; streak greenish black; metallic; opaque.

Fusible; soluble in nitric acid.

Western United States, Pennsylvania, Harz, Cornwall.




FIG. 68.—Chalcopyrite, Neudorf. Twinned parallel to (111).

#### Pyrite

This iron sulphide is more abundant than any mineral thus far considered. It is found in all kinds of rocks, with all kinds of mineral associates, and in all parts of the world. In Illinois it occurs in the underlying rocks—the shale, limestone, and sandstone, and in the sand and gravel carried in by Pleistocene glaciers.

The name pyrite ( $\pi\rho\rho$ , Greek “fire”) was used by Dioscorides and Pliny in the first century after Christ for minerals which gave sparks when struck by the hammer, and was applied not only to minerals in which the sparks are due to the combustion of the mineral itself but to hard minerals like flint in which the sparks are due to glowing particles intensely heated by the friction.

Pyrite differs from the iron sulphide already considered, pyrrhotite, in being neither magnetic nor bronze colored, and from the copper iron sulphide, chalcopyrite, in being brass yellow and not deep yellow as in chalcopyrite.

It occurs as masses, large and small crystals, and minute yellow specks in sedimentary, igneous, and metamorphic rocks.

Two forms of crystals are common and several others abundant. One of the most typical is that which has the outline of a cube (100) (Plate VII *a*), but whose true symmetry is indicated by the striations on each face. These striations show that the cube is built up by repetition of many planes of a form which is so characteristic of pyrite as to have been named the "pyritohedron" (pentagonal dodecahedron) (Plate VII *b*). The pyritohedron is formed when the alternate quarters of the four-faced cube (the inner part of




FIG. 69.—Pyritohedron derived by disappearance of tetrahedral planes darkened, and growth of the other planes.




FIG. 70.—Model of a diploid

Fig. 69) are developed, beginning with the plane (210). The black strips with which the glass faces of the outer figure are bound mark the pyritohedron. Clear-cut pyritohedrons are so common that every collector can obtain them. Occasionally a form shown in Figure 70, called a diploid, is found. It results when the plane (321) and each alternate plane in the right-hand octant of the hexoctahedron and the planes of the like symbol in the other octants are developed to the exclusion of their neighbors.

Both the diploid and the pyritohedron agree in this, that if revolved around any one of the four octahedral axes, i.e., the lines extending through the center of the crystal and perpendicular to an octahedral plane (Fig. 23), each of its faces would be in the position previously occupied by the adjoining face three times during a complete revolution. The faces are said therefore to have four trigonal

axes. If revolved around the crystallographic axes ( $a$ ,  $b$ ,  $c$ ) the planes are in similar positions twice during a complete revolution and hence these axes are called digonal axes. Since planes through any two of these digonal axes ( $a$  and  $c$ , or  $b$  and  $c$ , or  $a$  and  $b$ ) are planes of symmetry, the digonal axes are called didigonal axes. Pyrite crystals have three didigonal axes. Their faces are in pairs about a center. Hence the pyrite class of the regular system has a center, three planes, three didigonal and four trigonal axes of symmetry. This symmetry is called tesseral central symmetry.

Cube, octahedron, pyritohedron, and diploid appear in various combinations. The edges of the pyritohedron (210) are truncated by the cube (100) (Fig. 71).

Etching with aqua regia produces figures symmetrically arranged in respect to the cube planes.

Sulphides of nickel and cobalt are often found in pyrite as isomorphous intermixtures, i.e., mixtures of substances having the same crystalline form. Chalcopyrite, marcasite, and silver sulphide are often associated with it.

The most important impurity, however, is gold, which is often present as a metal scattered through the pyrite in invisible particles. Much of the gold of the world is now obtained by crushing, roasting, smelting, and cyaniding pyrites, and much of the placer gold may have originally come from the same source.

The chief use of pyrite is in the manufacture of sulphuric acid, sulphur, and iron oxide to be employed as polishing powder and paint. Iron for steel manufacture cannot be obtained from it, since thorough separation of the sulphur is almost impossible and a fraction of 1 per cent remaining in the iron renders it brittle while hot.



FIG. 71.—Model of a combination of pyritohedron (210) and cube (100).



## SUMMARY

*Pyrite*.— $\text{FeS}_2$ ; Fe=46.6 per cent, S=53.4 per cent. Regular; pyrite class: (100), (111), (210), (321), (421). Supplementary twins; granular; massive; brittle; fracture conchoidal.

Hardness=6; gravity=5.1. Pale brass yellow; streak greenish black; metallic; opaque.

Burns on charcoal and gives off  $\text{SO}_2$ ; fuses to magnetic globules; soluble in nitric acid.

Ubiquitous.

## Marcasite

The same chemical composition is ascribed to marcasite as to pyrite, namely,  $\text{FeS}_2$ . But there are pronounced physical differences. Marcasite crystallizes in the orthorhombic system and in the holosymmetric class. The crystals have a center of symmetry; three planes of symmetry intersecting at right angles in the crystallographic axes; and the  $c$  axis is a didigonal axis of symmetry, that is, if revolved around the  $c$  axis the planes assume similar positions twice in one complete revolution, making the  $c$  axis a digonal axis. Since planes of symmetry intersect in this axis, it is called a didigonal axis. A form




FIG. 72.—Marcasite




FIG. 73.—Marcasite

common to marcasite is composed of the prism (110), base (001), and brachydomes (011) and (013) (Fig. 72). Prisms of marcasite are usually terminated by various brachydome planes (Fig. 73). Isolated crystals are rare. Because of multiple twinning they generally show jagged outlines and re-entrant angles. Before marcasite was distinguished from pyrite, these forms were called "spearhead pyrites," "cockscorn pyrites," "radiated pyrites," "hepatic pyrites,"




PLATE VIII



Marcasite, Jo Daviess County, Illinois

PLATE IX



Marcasite disks, Gulf Mine, Sparta, Randolph County, Illinois



etc. Four or five individuals consisting of various dome and basal planes twinned parallel to the prism produce the "spear-head pyrites" (Fig. 74). "Cockscomb pyrites" result from repeated twinning parallel (110) so as to produce individuals parallel to each other (Fig. 75). The prisms are short and the striated basal plane long. Radiated, nodular, and stalactitic forms are abundant (Plate VIII). More marcasite than pyrite is found in Illinois. No. 3287 from Sparta shows disks which are not surpassed in abundance and perfection by any locality (Plate IX).

Plate X shows a portion of a large radiated mass and Plate XI a coating of marcasite on cubes of galena which are resting upon a




FIG. 74.—"Spearhead pyrites"




FIG. 75.—"Cockscomb pyrites"

botryoidal mass of sphalerite, well illustrating the association of the sulphides.

Haidinger (1845), recognizing the orthorhombic form of marcasite, reserved Pliny's term pyrite for the regular form and applied the old Moorish word "marcasite" to the orthorhombic. It is a common error to use the name pyrite when marcasite is meant. Marcasite is orthorhombic, white in color and streak, not as heavy and more easily decomposed than pyrite. Even in museum cases it disintegrates and becomes covered with white efflorescent iron sulphate (melanterite) and forms sulphuric acid which attacks the material upon which it rests. Its instability may be due to minute spicules of troilite ( $\text{FeS}$ ), a mineral heretofore identified in meteorites only.

When marcasite is heated to  $200^{\circ}\text{C}$ . in a sealed tube with a copper sulphate solution, it yields a solution entirely ferrous. Pyrite

treated in the same way yields a solution 19.9 per cent ferrous and

80.1 per cent ferric. Hence the formula of marcasite is  $\text{FeS}_2$ , while that of pyrite is  $\text{FeS}_2$  and  $\text{FeS}_2$ .

Since marcasite is most common in limestones and shales, and pyrite in crystalline rocks, their physical differences are doubtless due to their origin—marcasite having been hastily deposited from cold solutions, and pyrites slowly deposited from hot solutions, pyrite representing the more successful molecular grouping and showing that metamorphism produces in the lower zones of the earth's crust minerals of more complete symmetry, higher specific gravity, and greater hardness than those found in the upper zones.

Marcasite has been made in the laboratory from an acid solution and with temperatures not above  $300^\circ\text{C}$ . When the solution was neutral, pyrite crystals were formed. The most favorable conditions were found to be an acidity amounting to about 1.2 per cent free sulphuric acid and a temperature of  $100^\circ$ . At  $450^\circ$  marcasite changes to pyrite.

Both marcasite and pyrite are common fossilizing material because of the reducing action which decaying organisms exert upon iron sulphate solutions. Marcasite decomposing in moist air forms sulphuric acid, which can change the limestones surrounding it to gypsum.

#### SUMMARY

*Marcasite*.— $\text{FeS}_2$ ; Fe=46.6 per cent, S=53.4 per cent. Orthorhombic; holosymmetric;  $a:b:c=0.766:1:1.234$ ; forms (110), (001), (011), (018); twinned on (110). Crystals grouped, nodular, stalactitic, radiated, massive. Cleavage imperfect (111); brittle; fracture uneven.

Hardness=6; gravity=4.8. Pale brass yellow; streak greenish gray; metallic; opaque.

Soluble in nitric acid. Fuses readily.

Ubiquitous.

#### Arsenopyrite

Arsenopyrite resembles marcasite in its crystallography (Figs. 76 and 77) but is whiter, has a black streak, and is softer and heavier. It often contains as high as 9 per cent of cobalt in the form of an isomorphous intermixture of the cobalt sulphide, glaucodot. Arseno-






PLATE X



Marcasite, showing radiated internal structure

PLATE XI



Marcasite coating galena, Marsden Mine, Jo Daviess County, Illinois



pyrite is the chief source of arsenic, a metal used principally in the manufacture of Paris green, in various medicinal compounds and embalming fluids, and in glass and enamel manufacture.




FIG. 76.—Arsenopyrite




FIG. 77.—Arsenopyrite

#### SUMMARY

*Arsenopyrite*.— $\text{FeAsS}$ ; Fe=34.3 per cent, As=46.0 per cent, S=19.7 per cent. Orthorhombic; holosymmetric;  $a:b:c=0.677:1:1.08$ ; (110), (011), (014); twinned on (101); massive; cleavage fair (110); brittle; fracture uneven.

Hardness=5.5; gravity=6. Silver white; streak grayish black.

Soluble in nitric acid.

Freiberg, Cornwall, Ontario, Washington.

### CLASS III. SULPHANTIMONITES, SULPHARSENITES

#### PYRARGYRITE GROUP

Pyrargyrite, a silver sulphantimonite, and proustite, a silver sulpharsenite, called the "ruby silver ores" because of their wine-red color when fresh, are excellent examples of isomorphism, since they crystallize in forms very similar and with angles nearly identical, though one contains antimony and the other arsenic. Their structure places them in the hexagonal system, and their symmetry is said to be "ditrigoal polar" (tourmaline class). It is polar, inasmuch as the crystals are different at different ends. If the difference is not shown by developed planes, it may nevertheless be disclosed by the




FIG. 78.—Symmetry planes of a ditrigoal polar crystal.




FIG. 79.—Axes of hexagonal system

striations on the prism planes, since the striations are not symmetrical toward both ends. The symmetry is ditrigoal, since three planes of symmetry intersect in the  $c$  axis, and if the forms are revolved around this axis the planes are in a similar position three times in one complete revolution (Fig. 78).

In the hexagonal system are grouped those crystals which have three lateral axes of equal length intersecting each other at  $60^\circ$ , and perpendicular to them a vertical axis longer or shorter than they are. The method of naming the axes can be understood from Figure 79. The holosymmetric (holohedral) forms are three pyramids and three

prisms, and the ratios are always given thus:  $a:b:\bar{a}_1:c$ . The sum of the intercepts on the first three is always 'zero'. If  $(hkl)$  represents any



FIG. 80.—Model of a primary pyramid

symbol, then  $h+k+i=0$ . The primary pyramid (Fig. 80) is a form whose parameter is  $1:\infty:1:1$  and whose symbol is  $(10\bar{1}1)$ .

Parameter	Ratio	Symbol
$1:\infty:1:1$	$a : b : \bar{a}_1 : c$ $1 : \infty : 1 : 1$	$(10\bar{1}1)$

The diagonal (or "secondary") pyramid, whose relation to the primary pyramid is shown in Figure 81, is one whose parameter is  $2:2:\bar{1}:2$ .

Parameter	Ratio	Symbol
$2:2:\bar{1}:2$	$\frac{1}{2} : \frac{1}{2} : \frac{1}{2} : 1$ $1 : 1 : \bar{2} : 2$	$(11\bar{2}1)$




FIG. 81.—Basal section showing relation of primary, secondary, and dihexagonal pyramids and prisms. I =  $(10\bar{1}1)$ , II =  $(11\bar{2}1)$ , III =  $(21\bar{3}1)$ .




FIG. 82.—Model of a dihexagonal bipyramid.

The dihexagonal pyramid (Fig. 82) is a form whose parameter may be  $\frac{3}{2}:3:\bar{1}:3$ .

Parameter	Ratio	Symbol
$\frac{3}{2}:3:\bar{1}:3$	$\frac{a}{2}:\frac{b}{1}:\frac{a}{3}:\frac{c}{1}$	$(21\bar{3}1)$




FIG. 83.—Model of a primary hexagonal prism.




FIG. 84.—Secondary hexagonal prism



FIG. 85.—Model of a dihexagonal prism.

The relation of all three of these pyramids to each other is shown by a ground plan of the lateral axes (Fig. 81). The  $c$  axis is simply a point.

The three prisms are similar and their symbols are identical with those of the three pyramids, save that the number applied to the  $c$  axis is always zero (see Figs. 83, 84, 85).

Two hemihedral forms are common; first, that which results when the unit pyramid planes in alternate sextants only are developed. If the start is made with the front sextant  $(1101)$ , a positive rhombohedron ( $R$ ) results (Fig. 86). If the start is made with  $(01\bar{1}1)$ , a negative rhombo-

hedron ( $-R$ ) is produced. If the two planes in each alternate sextant of a dihexagonal pyramid are developed, a scalenohedron is formed



FIG. 86.—Rhombohedron ( $R$ ) resulting from disappearance of darkened planes of the interior figure—a hexagonal bipyramid.



FIG. 87.—Model of a positive scalenohedron.



FIG. 88.—Model of a scalenohedron truncated by a rhombohedron ( $R$ ).



FIG. 89.—Model of a prism truncated by negative  $\frac{1}{2}R$ .

( $21\bar{3}1$ ) (Fig. 87). A positive rhombohedron and scalenohedron are united in Figure 88, while in Figure 89 a prism is truncated by the negative  $\frac{1}{2}$  rhombohedron,  $-\frac{1}{2}R$ .



Crystals of pyrrargyrite and proustite occur in forms which are combinations of the secondary prism ( $11\bar{2}0$ ) terminated above with the plus rhombohedron ( $10\bar{1}1$ ), the flat minus rhombohedron ( $01\bar{1}2$ ), and the flat scalenohedron ( $21\bar{3}4$ ), while below appear the scalenohedrons ( $21\bar{3}\bar{1}$ ) and the rhombohedrons ( $01\bar{1}2$ ) and ( $01\bar{1}4$ ) (Fig. 90).




FIG. 90.—Pyrrargyrite, crystal form.

As usual with most minerals, well-developed crystals are rare. Ordinarily pyrrargyrite and proustite occur in masses.

Pyrrargyrite varies in color from the darkest varieties, which are of a deep ruby shade in thin splinters, to the lightest varieties, which are clear wine color. Upon exposure to light, pyrrargyrite becomes dead black. It should therefore be sheltered from light if the reddish color is to be preserved. The streak of pyrrargyrite is purplish red, while that of proustite is scarlet.

Like minerals in all systems other than the regular, these minerals divide entering light into two rays vibrating at right angles to each other and hence differently refracted. One of the rays is called the ordinary ( $\omega$ ) and the other the extraordinary ( $\epsilon$ ).

#### SUMMARY

*Pyrrargyrite*.— $\text{Ag}_3\text{SbS}_3$ ; Ag = 59.8 per cent, Sb = 22.5 per cent, S = 17.7 per cent. Hexagonal; ditrigonal polar (tourmaline class);  $a:c = 1:0.789$ ; ( $11\bar{2}0$ ), ( $11\bar{0}1$ ), ( $01\bar{1}2$ ), ( $21\bar{3}1$ ), ( $21\bar{3}4$ ), ( $01\bar{1}4$ ); massive. Cleavage imperfect ( $11\bar{0}1$ ) and ( $1010$ ); brittle; fracture conchoidal.

Hardness = 2.5; gravity = 5.8. Black to ruby, streak purplish red; metallic; adamantine; translucent. Refraction strong; mean refractive index in sodium light is 2.98 (stronger than that of diamond, and surpassed by cinnabar alone, 3.02).

Easily fusible; soluble in nitric acid.

Guanajuato, Mexico; Chili; Cordilleran states.

*Proustite*.— $\text{Ag}_3\text{AsS}_3$ ; Ag = 69.4 per cent, As = 15.2 per cent, S = 19.4 per cent. Crystallography similar to pyrrargyrite;  $a:c = 1:0.304$ .

Physical properties similar to pyrrargyrite, but color lighter and streak scarlet.  $\omega = 2.94$ .

Localities similar to pyrrargyrite.

## TETRAHEDRITE GROUP

In this group are two well-crystallized copper minerals, tetrahedrite ( $\text{Cu}_3\text{SbS}_3$ ) and tennantite ( $\text{Cu}_3\text{AsS}_3$ ), which are related to each other as were the two silver minerals, pyrrargyrite and proustite. So definite are they in crystal habit that they have been chosen to furnish the name of a crystal class, the tetrahedrite class (already illustrated by the diamond).

The tetrahedron ( $111$ ) is always developed, sometimes alone (Fig. 91), but usually combined with other tetrahedrons, trapezohedrons, and dodecahedrons. A common combination is a positive




FIG. 91.—Prevailing form of tetrahedrite.




FIG. 92.—Characteristic form of tetrahedrite.

tetrahedron ( $111$ ) with edges beveled by positive and negative three-faced tetrahedrons ( $211$ ), and with corners truncated by the minus tetrahedron ( $\bar{1}\bar{1}\bar{1}$ ) and beveled by the dodecahedron ( $110$ ) (Fig. 92). The negative tetrahedron appears as a small triangle on each corner and is usually dull or pitted with triangular markings, while the positive faces are bright.

When complementary forms of a three-faced tetrahedron ( $211$ ) are present, the positive form is often striated perpendicularly, while the negative is striated parallel to the dodecahedron edge which it truncates.

The edges and corners of one tetrahedrite crystal often project from the faces of another—a kind of twinning derived by a half-turn of the projecting crystal about a line normal to ( $111$ ).

In tennantite, dodecahedral or cubic faces usually predominate (Fig. 93).

The old German miners called both of these minerals (tetrahedrite and tennantite) *Fahlerz* ("pale ore"), and that term included several varieties which were later divided and named according to the locality in which they are found or according to some peculiarity due to varying composition. They contain not only copper, antimony, arsenic, and sulphur, but often bismuth, lead, silver, mercury, zinc, and iron in varying amounts, vicariously replacing each other.

Some of the varieties are the following: (1) freibergite (Freiberg, Saxony) often contains as much as 30 per cent of silver and is lighter than ordinary tetrahedrite in color; (2) schwartzite (Schwatz,




FIG. 93.—Tennantite (schwartzite)




FIG. 94.—Model of sandbergerite

Tyrol) contains 16 per cent mercury and occurs in black, drusy, dodecahedrons; (3) binnite (Binnenthal, Switzerland) appears in brilliant cubic crystals; (4) sandbergerite contains zinc and shows a tendency to develop large faces of the three-faced tetrahedron (211) (Fig. 94).

Tetrahedrite crystals are often coated with brassy, drusy chalcopyrite (Cornwall). They can all be recognized by polar symmetry, metallic luster, absence of cleavage, and reactions for copper, together with either arsenic or antimony.

Massive tetrahedrite is worked in Germany, Cornwall, and in many places in the Cordilleran states as a source of copper and silver.

## SUMMARY

*Tetrahedrite*.— $\text{Cu}_3\text{SbS}_3$ ; Cu=46.8 per cent, Sb=29.6 per cent, S=23.6 per cent. Regular; symmetry ditrigonal polar; (111), (11 $\bar{1}$ ), (110), (211); twinning axis the normal to (111); brittle; fracture sub-conchoidal.

Hardness=3.5; gravity=4.7. Lead gray; streak dark brown; metallic; opaque.

Fusible; soluble in nitric acid.

Germany, Bohemia, Cornwall, western United States.

## CLASS IV. HALOIDS

### THE SALT GROUP

The two most important minerals of this group are halite, sodium chloride (common salt), and sylvite, potassium chloride. Being soluble in water, they may be distinguished by their taste. While both are saline, sylvite is bitter.

Sylvite—the *sal digestivus sylvii* of the old pharmacists—has long been used for medicinal and chemical purposes. It was first discovered in the volcanic sublimations of Vesuvius, but larger quantities and finer specimens are now obtained at Stassfurt, Germany.

Halite has the distinction of being the mineral most largely used as a food. Other minerals furnish food for plants and thus indirectly sustain the life of man, but halite is the only mineral which is eaten in its natural state. It is also one of the most useful of minerals in chemical and manufacturing industries—glass manufacture, chlorine and soda works, etc.

Halite is the most abundant salt in ocean water, and in many seas in arid regions in various parts of the world. Salt Lake, Utah, contains 20.19 per cent sodium chloride; the Dead Sea, Palestine, only 7.8 per cent. The Dead Sea contains more magnesium chloride, about 11 per cent, and about 2 per cent of potassium chloride. Its total content of salts exceeds that of Salt Lake, being about 25 per cent.

By the evaporation of such seas in preceding geological periods, great beds of salt have been laid down. Such are those of New York, Michigan, Louisiana, Kansas, Nevada, and other states. The New York bed most utilized is seventy-five feet thick and lies at a depth of from one thousand to two thousand feet below the surface. The salt is obtained there, as it is in Michigan, Louisiana, Kansas, and other places, by driving pipes down to the bed, forcing hot water down to dissolve the salt, and carrying the brine thus produced up to evaporating basins, where it is collected, purified, and made ready for market. In some places salt is mined just as is coal. The great chambers remaining after the removal of mountainous masses of salt

in some of the mines of Germany, Austria, and Russia are among the most interesting and beautiful underground caverns that are to be found.

Salt obtained by evaporation of brines often exhibits skeletal cubes with cavernous faces (Fig. 95). Natural crystals show quite perfect cubes. Ordinarily both halite and sylvite occur in granular and massive condition and contain magnesium chloride, magnesium sulphate, and calcium sulphate as impurities.

When pure, salt is transparent and colorless. But varying tints of yellow, red, and blue are common. The coloring material is usually some iron oxide. It has been suggested that the deep blue color in salt (see No. 3288 from Stassfurt) may be due to metallic potassium.

Both halite and sylvite are highly diathermanous, allowing free passage of heat as other transparent bodies allow passage of light, and cleavage blocks are used to inclose gases in a tube with transparent ends which will readily transmit heat rays.




FIG. 95.—Halite cube from salt brine.

#### SUMMARY

*Halite*.— $\text{NaCl}$ ;  $\text{Na}=39.4$  per cent,  $\text{Cl}=60.6$  per cent. Regular; (100); massive. Cleavage perfect (100); brittle; conchoidal.

Hardness=2.5; gravity=2.2. Colorless; streak white; vitreous; transparent; refraction weak,  $n=1.54$ .

Soluble in three volumes of water; taste saline; fusible.

Lakes in arid regions, New York, Michigan, Louisiana, Kansas, Germany, Poland, Russia.

*Sylvite*.— $\text{KCl}$ ;  $\text{K}=52.4$  per cent,  $\text{Cl}=47.6$  per cent. Regular; symmetry holooxial; (100); massive. Cleavage perfect (100); brittle; fracture uneven.

Hardness=2; gravity=1.9. Colorless; streak white; vitreous; transparent; refraction,  $n=1.49$ .

Soluble in three volumes of water; taste saline, bitter; fusible.

Volcanic regions; Stassfurt.

### Fluorite

Fluorite ( $\text{CaF}_2$ ) occurs commonly in beautiful, clear-cut cubic crystals. Few minerals show their crystal form so plainly. The edges of the cubes are often beveled by the four-sided cube ( $310$ ) (Fig. 96) or by a hexoctahedron ( $421$ ) (Fig. 97). The flat four-faced cube is so characteristic that it has been called the "fluoroid."

The most typical twinning of fluorite is that where two cubes interpenetrate about a line normal to  $(111)$  with the result that the corners of one cube project from the faces of the other (Fig. 98). At a point where these corners emerge, the cube face is raised into low "vicinal" faces which form a flat four-faced cube with very high parameters, for example,  $32:1:0$ . "Vicinal" faces often replace




FIG. 96.—Fluorite




FIG. 97.—Fluorite

simple faces with low parameters. Multitudes of minute cubic crystals often cover the faces of the large cubes without detracting from their luster, since the minute faces are parallel to the large ones. Fluorite also occurs in granular and compact masses.

Its cleavage is remarkably perfect, yielding octahedrons (Plate XIII). This trait, together with its vitreous luster, aids in the ready identification of the mineral. The cubic faces have a higher luster than have the cleavage faces. Natural octahedrons are usually dull. Fluorite displays many beautiful colors. Large yellow cubes with corners beveled by  $(421)$  are found at Mehenoit, Cornwall; and transparent yellow cubes (see No. 503), at Durham. Beautiful purple crystals, No. 502 from Alston and No. 3852 from Cumberland, are shown. Pink and rose-red octahedrons are found near Chamonix, and at the island of Siglio (near Elba). Beautiful green, plum-





PLATE XII




Fluorite group from Rosiclare, Hardin County, Illinois

PLATE XIII



*a*, Fluorite cubes, Rosiclare, Illinois



*b*, Octahedrons cleaved out by ten-year-old boy, showing ease and regularity of cleavage.



colored, and amethystine crystals from the north of England adorn many museums. Since the colors are in layers parallel to cubic faces, some lilac cubes have a yellow center (Derbyshire) and some a deep green center. Amethystine (Nos. 3296 and 3297), green (Nos. 2278 and 2567), and colorless (No. 1788) specimens, all from Rosiclare, Hardin County, give an idea of the color of Illinois occurrences. Large crystal groups are shown in Nos. 708 and 958. No. 908 is a typical massive specimen. The most complete exhibit of Illinois fluorite is shown in the cases devoted to the economic exhibits.

When specimens of fluorite are heated, they lose in weight and color, and hence are thought to owe their color to hydrocarbon compounds. No relation has been traced between composition and color. Green and red crystals are strongly phosphorescent, that is, if heated above  $212^{\circ}$  F. or held in sunlight, when taken into a dark room they are luminous. The phenomenon called "fluorescence" is named from this mineral, since it is best illustrated when some richly colored specimens are held in the sunlight. They become hazy at a slight depth below the surface and diffuse from this superficial layer a plum-blue color. When viewed by transmitted light, they are light green. This fluorescence is due to transformation of light rays within the mineral, so that those emitted are of greater wave-length than those which entered.

Fluorite is low in refraction ( $\omega=1.43$ ). Were colorless isotropic crystals more abundant, the mineral would find extensive use in the manufacture of lenses and microscopic objectives where achromatic light is desired. It is used in the production of ultra-violet rays, for making vases and ornaments (Derbyshire, England), and for enamel and glass manufacture; but the largest quantities are employed as a



FIG. 98.—Model of fluorite twin

flux in iron smelting and in similar operations where a fluid slag is sought.

Being the only common mineral which contains fluorine in any large proportion, fluorite is the chief source of hydrofluoric acid. Moissan used it for vessels and stoppers in experiments on the isolation of fluorine, since it is one of the few substances which is not attacked by the gas.

Fluorite occurs chiefly associated with metallic ores, calcite, barite, and in tin-bearing veins which are marked by the presence of other minerals containing fluorine, such as topaz, lepidolite, tourmaline, and apatite. Illinois leads all other states in the production of fluorite, some years more than a million dollars' worth having been sold, greatly to the advantage of the steel industry, the manufacture of enameled bath tubs, the production of hydrofluoric acid, etc.

#### SUMMARY

*Fluorite*.— $\text{CaF}_2$ ; Ca=51.1 per cent, F=48.9 per cent. Regular; holosymmetric; (100), (310), (421). Massive; interpenetrant twins on axis normal to (111). Cleavage, perfect (111); brittle; fracture sub-conchoidal.

Hardness=4; gravity=3.2. Purple, blue, green, yellow, brown; streak white; vitreous; transparent. Refraction weak ( $\omega=1.44$ ); dispersion weak.

Fusible; soluble in nitric acid.

England, Germany, France, Illinois, Kentucky, Colorado.

#### Cryolite

Cryolite is a colorless or pure white translucent mineral composed of sodium and aluminium fluoride ( $\text{Na}_3\text{AlF}_6$ ). It is well named "ice-stone" ( $\chi\rho\nu\sigma\varsigma$ , "frost" and  $\lambda\iota\theta\sigma$ , "stone") because of the translucence of its white masses, because of its low melting-point (a splinter fuses in a candle flame), and because it has been obtained in the greatest amounts in the land of ice, West Greenland, where it was discovered near the town of Ivigtut in 1795. It was long the only source of aluminium and is still an important ore, though today bauxite, a brownish, earthy, hydrated aluminium oxide ( $\text{Al}_2\text{O}_3 + 2\text{H}_2\text{O}$ ) found in quantities in the southern states, furnishes most of the aluminium of commerce. Free crystals of cryolite are so rare that the author has never noticed one. Optical examination of crystalline

masses and etching with sulphuric acid, however, show the crystals to be repeatedly twinned and to belong to the "triclinic system." In masses they resemble cubes placed in parallel position. Their cleavage also appears to be parallel to cubic planes, hence it is easy to mistake their crystallography.

The cleavage, oblique striations, and hardness (2.5) distinguish cryolite from colorless fluorite and similar minerals. Heated with sulphuric acid, cryolite gives off hydrofluoric acid (HF).

#### SUMMARY

*Cryolite*.— $\text{Na}_3\text{AlF}_6$ ; Na=32.8 per cent, Al=12.8 per cent, F=54.4 per cent. Triclinic; (110), (001), (101); massive; cleavage, perfect, parallel (001), nearly perfect parallel (110), ( $\bar{1}$ 01); brittle; fracture uneven.

Hardness=2.5; gravity=3. Colorless; vitreous; transparent; refraction weak,  $\beta=1.36$ ; birefringence weak, positive.

Easily fusible; soluble in sulphuric acid.

Greenland.

## CLASS V. OXIDES

Of the minerals which consist of one or more basic elements united with oxygen, about seventeen are abundant and important. The oxides of silicon, that is, quartz, chalcedony, and opal, will be considered first; and second, the oxides of the metals, such as cuprite, zincite, corundum, hematite, spinel, magnetite, franklinite, chromite, cassiterite, rutile, pyrolusite, manganite, goethite, and limonite, will be taken up next.

### Quartz

Quartz ( $\text{SiO}_2$ ) is the most abundant mineral in the world. It is the chief constituent in the sands of the deserts and of the ocean shores, in the great layers of sandstone and quartzite which underlie the plains and outcrop in the mountains, and in most of the rocks that form the cores of mountain ranges.

No mineral has received more study than quartz. It furnished that thoughtful Danish physician, theologian, and geologist, Steno (1669), material with which to establish the "law of the constancy of a crystal angle," and it has been the subject of study for mineralogists ever since, seeming still to be able to reward the investigator with new facts.

The Greeks thought quartz to be ice so thoroughly frozen as to have lost the power of melting, and hence named it *κρύσταλλος*, that is, "ice," and today many persons say crystal when they mean quartz. The name "quartz" is an old German mining term used since the sixteenth century and now common to many languages.

Crystals of quartz are more abundant than those of any other mineral. They occur in the hexagonal system, and their symmetry is trigonal holoaxial, i.e., they have no plane nor center of symmetry but if revolved around the  $c$  axis their planes occupy similar positions three times during a complete revolution.

Prism (1010) and rhombohedron (1011) planes are nearly always present and combined, as shown in Figure 99. In Hungary and Brazil are found crystals which contain these planes only. It will be seen that when revolved around the  $c$  axis an upper rhombohedron





PLATE XIV



*a*, Smoky quartz, "cairngorm," from Montana.



*b*, Quartz, Montgomery County, Arkansas

**PLATE XV**



**Quartz group, Montgomery County, Arkansas**



would be in the front three times during a complete revolution; hence the crystals are "trigonal."

Several planes occur with such regularity that, for sake of abbreviation, to represent them a letter is used instead of the symbol, for




FIG. 99.—Quartz; prism and rhombohedron.




FIG. 100.—Quartz




FIG. 101.—A form of quartz common at Alston Moor, England.




FIG. 102.—Quartz

example,  $R$  stands for the plus rhombohedron ( $10\bar{1}1$ ),  $\bar{z}$  for the minus rhombohedron ( $01\bar{1}1$ ),  $m$  for the prism ( $10\bar{1}0$ ),  $S$  for the right trigonal pyramid ( $11\bar{2}1$ ), and  $x$  for the right plus trapezohedron ( $51\bar{6}1$ ) (Figs. 100-102). The construction of the right and left plus trapezohedrons

is explained below (see Figs. 103 and 104). Rarely is a crystal terminated by a single rhombohedron, as in Figure 102. The usual termination is a combination of plus and minus rhombohedrons,  $R$  ( $10\bar{1}1$ ) and  $z$  ( $0111$ ), of different sizes. Sometimes  $R$  and  $z$  are so nearly of the same size as to resemble a unit pyramid (Fig. 101).

In some crystals, like those from Alston Moor, England, the prism planes are very small or disappear, and the result resembles a bipyramid (Fig. 101). But their trigonal character can be discovered by heating them and plunging them into water, when they cleave




FIG. 103.—Quartz; positive left trigonal trapezohedron.




FIG. 104.—Quartz; positive right trigonal trapezohedron

into imperfect rhombohedra. In some crystals the right edge between  $R$  and  $m$  is truncated by the plane  $s$  whose symbol is ( $11\bar{2}1$ ) (Fig. 102). It has the direction of the diagonal pyramid, yet occurs but three times instead of six, so is recognized as a hemihedral form of the diagonal pyramid called the "right trigonal pyramid." The left trigonal pyramid appears on the left-hand side. The trigonal pyramid is often accompanied by a trapezohedral face  $x$  ( $51\bar{6}1$ ). Figures 103 and 104 show right-handed and left-handed trapezohedrons. They result when the alternate upper sextants of a scalenohedron and the corresponding sextant below are developed. The trapezohedral planes are marked by the letter  $x$ . The right- and left-handed crystals may be distinguished in three ways: first, in a



PLATE XVI



Quartz, Bourg de Oisans twin, Hot Springs, Arkansas

right-handed crystal the  $x$  plane is below the right-hand corner of the rhombohedron; second, the direction of the zone  $z s x m$  is that of the thread of a right-handed screw; and third, the striae on  $s$  are parallel to the edge  $sR$ .

Twinning in quartz crystals is common according to three laws: first, two crystals of the same sort, both right-handed or left-handed, may be united parallel to the  $c$  axis in interpenetrant twins (Fig. 105). Thus  $x$  may appear at each corner and  $R$  and  $z$  in the same plane. However, since  $R$  is usually smooth and bright, while  $z$  is pitted or




FIG. 105.—Two right-handed quartz crystals twinned parallel to  $c$  axis.




FIG. 106.—Brazil twin. Right- and left-handed quartz crystals interpenetrating.

coated, they may be distinguished from each other. The boundaries of the two interlacing crystals show a zigzag pattern. Second, right-handed and left-handed crystals interpenetrate parallel to the  $c$  axis and at the same time parallel to the diagonal prism ( $11\bar{2}0$ ). In this case twinning is betrayed by the  $x$  and  $z$  faces (Fig. 106). This is called the Brazil twin. Figure 107 shows two right juxtaposed Brazil twins. Third, Bourg de Oisans in Dauphiny (France) has long been noted for fine quartz crystals twinned in the manner shown in Figure 108 (Nos. 3307, 3308, and 3315). Recently Japan has furnished the museums of the world with a large number of these twins. They are united parallel to the diagonal pyramid ( $11\bar{1}22$ ) so that the  $c$  axis and the line of union form a zigzag.



Quartz illustrates not only the geometrical but the optical, electrical, thermal, and chemical features of crystals as well. Optical properties throw much light upon its internal structure. The connection between geometrical and electrical properties may be illustrated as follows:




FIG. 107.—Brazil twin. Two right-handed crystals juxtaposed.

The three horizontal axes are polar (i.e., not symmetrical around the center), for one end of each axis emerges through a prism edge that is truncated by the planes  $s$  and  $\alpha$ , and the other through a prism lacking these planes. Since the horizontal axes are polar, they exhibit pyro-electric polarity. Finely powdered red lead and sulphur are sifted through a piece of cloth, thus becoming electrified by friction. The red lead is positive, the sulphur negative. When they touch the heated quartz crystal, the  $s$  and  $\alpha$  faces become negatively electrified on cooling, i.e., attract the red lead.




FIG. 108.—Quartz twinned on (1122). Bourg de Oisans twin.



FIG. 109.—Quartz etched with hydrofluoric acid; A, left-handed; B, right-handed.

The action of caustic alkalis or hydrofluoric acid in nature or in the laboratory produces different effects on different planes, and thus the right-handed and left-handed nature of the crystals can be made evident (Fig. 109). The etching shows that the forms are related to each other as are a right and left glove, and hence they are said to be entantiomorphous.

The manner of crystal growth is shown by some specimens of quartz which within the transparent outer crystal have different layers of cloudy material forming outlines of the planes at different stages of the crystal's development. This is called "ghost quartz." "Capped quartz" has an inner kernel separated from the outer by a layer of clay or other substance which permits them to be taken apart. "Twisted quartz," while seeming to consist of a single warped crystal, in reality is composed of many individuals, each turned through a small angle so as to produce a spiral effect. Quartz is practically lacking in cleavage. Only by heating and plunging in cold water can rough rhombohedrons be obtained.

Intergrowths of right- and left-handed quartz break with a wavy surface, producing "ripple fracture," while the ordinary fracture is conchoidal.

As might be expected in a mineral so abundant and so varied in its surroundings and mode of formation, quartz exhibits great variety in form and appearance.

Rock crystal, or mountain crystal, as it was called by indefatigable collectors who sought fine specimens in the mountain fastnesses of Switzerland and the Tyrol, is a clear, transparent variety well marked in crystallization. It was formed in non-resistant rocks like clay and sandstone, or in cavities in igneous or metamorphic rocks, which afforded it opportunity to develop its own planes (see Nos. 1772 to 1779, 3896, etc.). Crystals of remarkable size have been found in the Alps, Brazil, Japan, and Madagascar. A crystal twenty-five feet in circumference was found in Madagascar. A famous cave in the Berner Oberland in Switzerland yielded five hundred tons of quartz crystals. Herkimer County, New York, is noted for its beautiful, transparent crystals. Little Rock, Arkansas, has annually furnished countless souvenirs of this kind to the tourists in that region (Nos. 473, 1775, etc.). Multitudes of geodes found at Warsaw in Hancock County, Illinois, varying in size from a hazelnut to nodules a foot or more in diameter are lined with clear quartz crystals. There is hardly a state in the United States in which fine quartz crystals have not been found.

Clear crystals have long been used for ornaments. Beautiful transparent globes as much as six inches in diameter have been cut from quartz found in Japan.

Inclosures of foreign substances may add to the beauty of the mineral. Crystals from Herkimer County, New York, often inclose anthracite flakes. Spangles of mica and of hematite produce the shimmer seen in aventurine quartz. Fibrous actinolite, asbestos, or rutile needles produce beautiful effects. Silky fibers of asbestos or of quartz replacing them give a peculiar band of color to the opalescent quartz called "cat's eye" (a name more properly applied to a variety of chrysoberyl). The golden yellow crocidolite from South Africa owes its beauty to this cause. Cavities shaped like a quartz crystal and containing water or liquid carbon dioxide are often seen.



FIG. 110.—Airy's spiral in right-handed crystal.

There are several varieties based on color: milk quartz (No. 1773) is white and opaque, morion (No. 4645) is black, smoky quartz (No. 3312) brown. Morion and smoky quartz, abundant in the Alps, are colored by a hydrocarbon which disappears upon heating. When cut into gems smoky quartz is called "cairn-gorm." A clear yellow quartz also colored by hydrocarbon is named citrine. Prase (No. 3225)

is green from needles of actinolite. Rose quartz abundant in the Black Hills is pale red from solution of salts of titanium or manganese (Nos. 1681, 3323, 4513). Amethyst, a violet quartz, has long been one of the most popular of semiprecious stones (Nos. 4458, 588, 589, 3319, etc.). Its color is thought to be due to manganese, though upon heating to a temperature of 250° it changes to yellow. The color is often irregularly distributed, white, opaque layers alternating with transparent violet and brown layers (Plate XVII). Microscopic examination shows alternating layers of right- and left-handed lamellae. Where the right and left layers are mingled, converging light produces lines known as "Airy's spirals" (Fig. 110). The interpenetration of right- and left-handed layers produces roughly striated surfaces. The more strongly colored parts are "biaxial," that is, they have two directions in which light is not doubly refracted.



**PLATE XVII**



**Amethyst, Thunder Bay, Lake Superior**

Upon heating, the biaxial character disappears, showing that it, as well as the color, is due to easily destructible material. Beautiful violet specimens are found in Brazil, Ceylon, the Urals, Colorado, and north of Lake Superior. By some authors all quartz showing Airy's spirals is called amethyst, whatever the color.

Quartz quite commonly replaces organic substances, producing such objects of permanent beauty as silicified wood; or fills cavities formerly occupied by fluorite, calcite, barite, etc., assuming the shape of these minerals, thus producing pseudomorphs. A cube of quartz may be a pseudomorph after fluorite; fibrous quartz, a pseudomorph of fibrous gypsum; cellular quartz, the capping of calcite crystals later dissolved by water. Quartz is the great repairing agent of nature, since it so commonly cements crevices in rock layers and microscopic fissures in minerals.

#### SUMMARY

*Quartz*.— $\text{SiO}_2$ ; Si=46.7 per cent, O=52.3 per cent. Hexagonal; symmetry, trigonal holoaxial (quartz class).  $a:c=1:1.1$ .  $m=(10\bar{1}0)$ ,  $R=(10\bar{1}1)$ ,  $z=(01\bar{1}1)$ ,  $s=(11\bar{1}2)$ ,  $x=(51\bar{6}1)$ . Twinned on  $m$  ( $10\bar{1}0$ ) or ( $11\bar{1}2$ ). Massive, cleavage parallel  $R$  very imperfect; brittle; fracture conchoidal.

Hardness=7; gravity=2.65. Colorless; vitreous; transparent,  $\omega=1.544$ . Uniaxial; double refraction, positive weak; rotary polarization, strong.

Infusible before blowpipe; insoluble in acid.

Ubiquitous.

#### Chalcedony

Chalcedony is identical with quartz in chemical composition and in many physical properties, but differs in several respects. First, it never shows crystal planes but occurs in translucent or opaque botryoidal, reniform, or stalactitic masses composed of microscopic fibers. Second, the fibers composing it are optically biaxial. Quartz is uniaxial. The refractive index and fusing-point of chalcedony differ from those of quartz.

Chalcedony is waxy or greasy in luster, and somewhat splintery in fracture. It is much more soluble in potassium hydrate than is quartz. It is deposited from aqueous solution, and is found in veins or other cavities in various kinds of rocks. Usually it has banded structure

and shows a great variety of colors. The banding is due to alternation of different-colored layers of chalcedony, quartz, and opal.

There are many varieties. The translucent, waxy, cream-colored, slightly banded variety is chalcedony proper. The red variety is called carnelian; the brownish-red, sard; the leek-green, plasma; the apple-green, chrysoprase (No. 3324); chalcedony with blood-red spots of jasper, heliotrope (No. 3329). Agate (Nos. 496-99, etc.) is composed of successive bands of chalcedony, carnelian, jasper (No. 568), smoky quartz, amethyst, etc., that have been laid down in the lining of a cavity, the outer band being formed first and the others successively until the cavity has been more or less filled. Usually the last stages permit of the formation of good quartz crystals. The fineness of some of the layers is a cause of wonder, and as the cavities often have no visible outlet, the manner in which the silicon reached its resting-place is enigmatical unless it be explained as being due to the solidification of colloidal silica. Sometimes the solution has deposited curvilinear layers first, and later parallel bands perfectly horizontal (No. 496). Such deposits are prized for cameo cutting. The figure is cut in one layer and the background in a layer of different color. When the layers are black and white, the material is called onyx. When red or brown, sardonyx. In moss agate (Nos. 4398 and 1851), banding is inconspicuous, but dendritic inclusions of chlorite, manganese oxide, and other substances occur. Flint (No. 3330) is a translucent to gray, brown, or nearly black chalcedony consisting largely of the remains of diatoms, sponges, and other marine organisms. The best variety is found in the chalk cliffs of England. Hornstone, as its name implies, resembles horn in color and in streaked appearance. It is more brittle, splintery, and soluble than flint, and less pure. Chert is still farther removed from flint in these respects. Its color is white or gray, and the impurities are calcareous and arenaceous substances. Jasper is a creamy, brown, or red chalcedony containing ferruginous, calcareous, or arenaceous substances as impurities.

These last four-named varieties border very closely upon rock species because of their impurities. A final step is represented by granular to massive silica, which occurs in large bodies and forms the rock called quartzite.

PLATE XVIII



Moss agate, India





## SUMMARY

*Chalcedony*.— $\text{SiO}_2$ ; Si=46.7 per cent, O=53.3 per cent. Cryptocrystalline, apparently amorphous, concretionary, botryoidal, stalactitic, arborescent. Brittle; fracture conchoidal.

Hardness = 7; gravity = 2.6. Waxy, yellow, red, brown, translucent to opaque; optically biaxial, thus differing from quartz, which is uniaxial.

Infusible; insoluble.

Ubiquitous.

## Opal

Opal differs from quartz and chalcedony in constitution, since water forms a part of its molecule ( $\text{SiO}_2$  and  $\text{H}_2\text{O}$ ). When water occurs in quartz it is mechanically, not chemically, included and is given off with a cracking noise (decrepitation) when the quartz is heated above the boiling-point ( $100^\circ \text{C}$ ). In opal it is chemically combined in definite proportion, constituting most commonly about 10 per cent of the mass. At the moment of solidification a molecule of silica attracts one or more molecules of water, called the water of constitution, in distinction to mechanically included water. It is held scarcely more firmly than are the different molecules which constitute water itself, and is given off upon heating to  $100^\circ \text{C}$ .

Opal differs from quartz not only in composition but in physical characteristics also. It lacks crystal form, lines cavities with layers which on their free sides are botryoidal, reniform, or stalactitic, is softer and lighter than quartz, has a greasy luster, and is completely soluble in hot potassium hydrate.

Of the several varieties, the most important is precious opal (Nos. 3741 and 4725), which is highly prized for gems in spite of its opacity, low refraction, and moderate hardness. Its beauty depends upon the wonderful play of colors which the thin films composing it cause by their difference in refractive index.

For centuries near Czerwenitz, Hungary, beautiful yellow, red, green, and blue opals have been found disseminated through an altered trachyte. Similar rock near Queretaro, Mexico (No. 3334), furnishes a fiery red opal without much play of color, the so-called fire opal. Blue and green opals of great beauty are obtained from nodules of brown jaspery limonite in Queensland (No. 3741). At all these localities opals of various other shades also are found.

A perfectly transparent colorless and glassy opal found in botryoidal masses is called hyalite. Because of strain due to the solidification of the original jelly mass, it often shows double refraction. Siliceous sinter is fibrous, stalactitic, porous, or powdery opal deposited by hot waters in Yellowstone Park, Iceland, and New Zealand. Its structure is influenced by the algae which live in the water, just as blades of grass determine the form of ice deposited on them by winter's rain. The shapes of the sinter or geyserite are often fantastic and beautiful.

#### SUMMARY

*Opal*.— $\text{SiO}_2 \cdot \text{H}_2\text{O}$ . Amorphous; brittle; fracture conchoidal.

Hardness=6; gravity=2. Colorless, white, red, brown, yellow, green, blue. Vitreous.

Infusible; yields water; soluble in potassium hydrate.

Hungary, Bohemia, Australia, Mexico, Idaho, Montana.

#### A. THE MONOXIDES

Minerals composed of an equal number of atoms of a bivalent metal and of oxygen are called monoxides. The two leading examples are cuprite and zincite.

##### Cuprite

Cuprite ( $\text{Cu}_2\text{O}$ ), or "red copper ore," contains the largest per cent of copper (88.8 per cent) of any mineral except native copper.




FIG. III.—Cuprite

It occurs in well-formed crystals which exhibit the cube (100), octahedron (111), dodecahedron (110), trapezohedron (211), and trisectahedron (221). A typical crystal is shown in Figure III. Granular aggregates and dense masses are common. Fresh surfaces of cuprite have a shining red appearance like proustite, but the streak is brownish red, while that of proustite is scarlet. Cuprite is the harder of the two

minerals. It is found in mineral veins where chalcocite and chalcopyrite have been subject to oxidizing agencies. Hence many veins

which above the water line contain cuprite are composed of sulphides, chalcocite, chalcopyrite, etc., below that line. Cuprite is often coated with malachite, a green copper carbonate, and at times the change extends throughout the crystal without destroying the external form. There are three varieties of cuprite: first, the ordinary crystallized form described above; second, "chalcotrichite"—"plush copper ore," consisting of slender fibers as fine as a hair, elongated in the direction of a cube edge or a cube diagonal; third, "tile ore," a brick-red, earthy mixture of cuprite and limonite.

## SUMMARY

*Cuprite.*— $\text{Cu}_2\text{O}$ ; Cu=88.8 per cent, O=11.2 per cent. Regular (111), (100), (110); brittle; fracture uneven.

Hardness=3.5; gravity=6. Cochineal red; streak brownish-red. Metallic, adamantine; translucent; refraction very strong,  $\omega=2.85$ .

Fusible; soluble in strong hydrochloric acid.

In Cordilleran states hydrochloric acid.

## Zincite

In Sussex County, New Jersey, zincite ( $\text{ZnO}$ ) occurs in quantities sufficient to make it a profitable source of zinc (Nos. 314, 315, 524, 747). It is of brownish or deep red color, and has a characteristic orange-yellow streak. It is usually massive, but when crystals are found they are in the hexagonal system and "hemimorphic" or "polar," since their planes are not symmetrically arranged around the center (Fig. 112). Some specimens of zincite contain as much as 7 per cent of manganese oxide ( $\text{MnO}$ ), which is present in solid solution as an isomorphous mixture. That it is a solid solution is suggested by the fact that with increase of the amount of manganese the mineral becomes more yellow.




FIG. 112.—Zincite

## SUMMARY

*Zincite.*— $\text{ZnO}$ ; Zn=80.3 per cent, O=19.7 per cent. Hexagonal. Symmetry, dihexagonal polar. Cleavage, perfect (0001), (10\bar{1}0). Brittle; fracture subconchoidal.

Hardness=4; gravity=5.6. Blood red, streak orange; luster, sub-adamantine; translucent; double refraction positive.

Infusible; soluble in hydrochloric acid.

New Jersey.

#### B. THE SESQUIOXIDES

Minerals having three atoms of oxygen to two of another element, usually a metal, are called sesquioxides. As examples we select corundum and hematite.

##### Corundum

Since very early in human history corundum ( $\text{Al}_2\text{O}_3$ ) has been recognized and used by man. The words sapphire and ruby occur in literature two thousand years old. Sapphire is blue corundum, and ruby the red variety. Ordinary corundum, being non-transparent and unattractive in color, did not early arrest attention. When cleaved, its fragments resemble feldspar, one of the most abundant of minerals, and hence would not be easily recognized. Though the name corundum is an old Hindu word, the mineral was first brought from China to Europe, and was first analyzed by a German, Klaproth, in 1787.




FIG. 113.—Corundum

A third variety of corundum, emery, is granular and dark in color because mixed with iron oxides such as magnetite. The name "emery" is of Greek origin, and some of the isles of Greece, especially Naxos, were long the chief source. The ancient lapidaries used it for grinding and

polishing, and today its use is very extensive, although pure corundum is preferable, being superior to emery in hardness.

Sapphire, ruby, ordinary corundum, and emery differ only in purity of chemical composition, color, and in structure. Emery is the least pure, is opaque, and occurs only in grains. Corundum and the transparent varieties crystallize in good hexagonal crystals of rhombohedral symmetry (calcite class). While the sapphire and ruby are rare, corundum is abundant and may best be taken as the representative of the species. Figure 113 shows a common and typical

form. It is composed of a rhombohedron truncated with a basal plane. In such a crystal it is evident that there are three planes of symmetry intersecting in the  $c$  axis, and that if the mineral were revolved around this axis the planes would repeat themselves three times during a complete revolution. Thus the  $c$  axis is an axis of trigonal symmetry. The three horizontal axes are axes of binary symmetry.

More common than the simple crystals are those composed of rhombohedrons  $R$  ( $10\bar{1}1$ ), dihexagonal pyramids ( $24\bar{2}3$ ), diagonal prisms ( $11\bar{2}0$ ), and base ( $00\bar{0}1$ ) (Fig. 114).

Repetition of dihexagonal pyramid planes of different inclination to the  $c$  axis produces undulating prisms and rounded crystals. Often




FIG. 114.—Corundum



FIG. 115.—Corundum with twinning lamellae parallel to  $R$ .

twinning lamellae parallel to the rhombohedron ( $10\bar{1}1$ ) cause striations on the basal plane and divide the crystal into sextants (Fig. 115). When the basal plane is polished, these striations reflect the light in such a manner as to produce a beautiful six-rayed star ("asterism").

The largest amount of corundum has been found as water-worn pebbles, which, though rounded, usually retain their crystal shape because of their great hardness. Coarse pieces and granular masses intergrown with mica are also characteristic. In the Boston Society Natural History Museum is a group of large corundum crystals weighing several hundred pounds.

True cleavage is wanting, but there is a lamellar separation parallel to  $R$  and the base, due to layers of twin crystals.

Vitreous luster is characteristic. Few minerals have as great variety of color. Colorless examples are rare. When the transparent or translucent varieties are red, they are called ruby. If of the shade known as "pigeon blood," they are more highly prized than any other gem. When deep blue to lilac in color, they are called sapphire; yellow to brown, oriental topaz; green, oriental emerald; purple, oriental amethyst—a good example of the common tendency to call one object by the name of another in order to enhance its value—a tendency which should be resisted. The color is due to oxides of iron or chromium. Sapphire exposed to action of radium bromide assumes successively green, light-yellow, and dark-yellow tints, while ruby develops in succession shades of violet, blue, green, and yellow. They regain their original color upon heating.




FIG. 116.—Cross-section of dichroscope.




FIG. 117.—Direction of vibration of two rays of light passing through calcite prism.

In corundum (as in other transparent minerals not belonging to the regular system) the color depends upon the direction in which light passes through the crystal. Hence false gems can be readily separated from genuine ones. The mineral needs only to be examined through a dichroscope ( $\delta\iota\varsigma$ , "two,"  $\chi\rho\omega\mu\alpha$ , "color") (Fig. 116), an instrument consisting of a metal tube three inches long fitted with a weak lens at the end *a* to be held to the eye and having a small square opening at the end *b* against which the mineral to be examined is held. In the tube is a cleavage rhomb *c* of calcite ( $\text{CaCO}_3$ ), a transparent mineral which has the power of dividing a ray of entering light into two rays that vibrate in planes at right angles to each other and consequently make a single point appear double (Plate XXI). Such a thickness of calcite is chosen as to make the square opening appear like two openings side by side. The transmitted light vibrates in the direction indicated by the arrows in Figure 117. One ray of light is called the ordinary ( $\omega$ ) and the other the extraordinary ( $\epsilon$ ).

When a piece of colored glass is placed on the square opening, both images have the same color. If a ruby is placed on the opening, the ordinary ray ( $\omega$ ) looks deep red, while the extraordinary ( $\epsilon$ ) is violet red. A sapphire appears deep blue for the ordinary ray ( $\omega$ ) and greenish blue for the extraordinary ( $\epsilon$ ) (Figs. 118 and 119).

Rubies and sapphires are found in granite, basalt, gneiss, mica and chlorite schist, granular limestone and dolomite, and in gravels derived from them. The finest sapphires have been obtained in Ceylon, the most valuable rubies in Burma. Montana and North Carolina furnish valuable sapphires, rubies, corundum, and emery.




FIG. 118.—Ruby



FIG. 119.—Sapphire

The examples shown come from widely distributed localities: rubies from Burma (No. 3339), sapphires from Kashmir (No. 3341) and from Queensland (No. 3340), corundum from North Carolina (No. 3342) and New Jersey (Nos. 506 and 519), emery from Massachusetts (No. 1254).

#### SUMMARY

*Corundum*.— $\text{Al}_2\text{O}_3$ ; Al = 52.9 per cent, O = 47.1 per cent. Hexagonal, symmetry rhombohedral (calcite class).  $a:c = 1:1.363$ .  $R$  (1011),  $(24\bar{2}3)$ ,  $(11\bar{2}0)$ . Parting or pseudo-cleavage; brittle; fracture conchoidal.

Hardness = 9; gravity = 4. Many colors, vitreous, transparent, dichroic, refraction strong,  $\omega = 1.768$ . Double refraction, negative weak.  $\omega - \epsilon = 0.008$ .

Infusible; insoluble.

North Carolina, Montana, Idaho, India.

#### Hematite

Of all minerals in the mineral kingdom, none is more important from a human standpoint than hematite ( $\text{Fe}_2\text{O}_3$ ), inasmuch as it is the mineral which furnishes the greatest quantity of iron—a metal upon which modern civilization is founded and which may be said to furnish a standard of development of a people.



In appearance hematite varies greatly with its physical condition. When well crystallized it is metallic, black, 6.5 in hardness and 5 in specific gravity. When earthy and friable it is submetallic, red, soft, and of low specific gravity.

Hematite is isomorphous with corundum, that is, it has the identical shape, though it is a different chemical substance. The best crystals are found in igneous rocks. The island of Elba has for many years furnished beautiful crystals which show the same rhombohedral symmetry already studied in corundum.

The rhombohedron  $R$  ( $10\bar{1}1$ ) occurs alone (Fig. 120) or combined with a negative flat rhombohedron  $-\frac{1}{2}R$  ( $01\bar{1}2$ ). Rounded crystals (Fig. 121) composed of a flat rhombohedron  $\frac{1}{2}R$  ( $10\bar{1}4$ ), the ordinary rhombohedron  $R$  ( $10\bar{1}1$ ), and the bipyramid ( $22\bar{4}3$ ) are characteristic. Tabular crystals composed of a broad basal plane ( $0001$ ), truncating short rhombohedrons  $R$  ( $10\bar{1}1$ ), and secondary prism planes ( $11\bar{2}0$ ) often twinned parallel to a prism plane are common (Fig. 122). They are often so grouped as to form rosettes, "iron roses."




FIG. 120.—Model of a rhombohedron ( $10\bar{1}1$ ) =  $R$ .

Individual crystals of minute size occurring in myriads sometimes constitute great masses of ore and form the variety called micaceous hematite. The fine scales of this variety of hematite are usually imperfectly cemented so that they easily rub off and give a false impression of the hardness of the mineral.

When heated in a reducing flame, the mineral loses its red color and becomes magnetic but does not melt. The electrical conductivity of hematite has been accurately measured and is found to be two times as great in the vertical as in the horizontal direction of the crystal. Hematite is one of nature's most universal paints. It colors the rocks red and brown and yellow as it varies in amount and in its degree of oxidation. About 6 per cent of the



PLATE XIX



*a*, Botryoidal hematite, Cumberland, England



*b*, Limonite, Hardin County, Illinois

earth's crust is iron and a large part of iron is in the form of hematite. Illinois has no workable deposits, but the mineral occurs in flakes, incrustations, or red ochreous balls, or as coloring matter in all parts of the state and in massive micaceous or compact pieces in the drift.

In many parts of the world granular hematite (No. 1581) forms such extensive deposits as to furnish the largest source of iron. Well-crystallized lustrous hematite is capable of receiving a high polish and reflects the light as does a looking-glass, and hence has been called "specularite" (Nos. 3894 and 3895). The fibrous and columnar varieties are composed of individual threads or pencils more or less parallel and ending in rounded grapelike (botryoidal) (No. 3343) or




FIG. 121.—Hematite




FIG. 122.—Tabular hematite crystal twinned parallel to the prism.

kidney-shaped (reniform) surfaces and exhibiting curvilamellar markings in various places (No. 3345). Botryoidal masses break up into conical forms known as "pencil ore." Masses that are earthy and soft enough to adhere to the fingers like graphite, and usually bright red in color, suggested a name for the mineral (hematite, "blood-stone") to Theophrastus three hundred years before Christ (Nos. 343, 608, and 4496). Thin flakes such as may be seen in microscopic slides appear blood red in transmitted light, just as the powder does in ordinary light.

The great deposits of hematite in Michigan, Minnesota, Wisconsin, and Alabama make it possible for the United States to lead the world in the production of iron.

## SUMMARY

*Hematite*.— $\text{Fe}_2\text{O}_3$ ; Fe = 70 per cent, O = 30 per cent. Hexagonal; symmetry dihexagonal, alternating (calcite class).  $a:c=1.3656$ .  $R$  (1011),  $-\frac{1}{2}R$  (0112),  $\frac{1}{2}R$  (1014), (1120), (2243), (0001). Cleavage (parting) parallel  $R$  and (0001) imperfect; brittle; fracture uneven.

Hardness = 6; gravity = 5.2. Iron black to blood red; streak brownish red or purple; metallic; in thinnest pieces translucent and red.

Infusible alone before blowpipe; powder difficultly soluble in concentrated hydrochloric acid.

Elba, Switzerland, New York to Alabama, Minnesota, Wisconsin, and Missouri.

## C. HYDRATED SESQUIOXIDES

Three minerals may be chosen to represent this group. They are:

Manganite

$\text{Mn}_2\text{O}_3 \cdot \text{H}_2\text{O}$

Goethite

$\text{Fe}_2\text{O}_3 \cdot \text{H}_2\text{O}$

Limonite

$2\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$

**Manganite**

From a mineralogical point of view manganite ( $\text{Mn}_2\text{O}_3 \cdot \text{H}_2\text{O}$ ) is of more importance than pyrolusite, though not so commercially, since




FIG. 123.—Manganite




FIG. 124.—Manganite twinned parallel to (011).

pyrolusite and another manganese oxide, psilomelane, are mined in great quantities, while manganite is comparatively rare. Manganite is well defined in its physical and chemical characteristics. It occurs in steel gray to black, moderately hard (hardness, 4), orthorhombic prisms, which are usually grouped in bundles and striated vertically. The prisms end in basal planes and are striated hori-

zonally (Fig. 123). The customary planes are (210), (110), (120), (010), (111), (101), (011), and (021). Twins parallel to (011) are common (Fig. 124). Fibrous, radiated, and granular forms are representative (No. 3356).

Manganite is formed by deposition of manganese oxide in many springs and often replaces other minerals, assuming their forms, i.e., becoming a pseudomorph. For example, at Ilfeld, Germany, manganite (No. 3357) replaces calcite, and since the calcite has been deposited from an aqueous solution it is natural to conclude that manganite has a like origin. On the other hand, manganite changes into pyrolusite by loss of water. The process which occurs in nature is imitated by slow heating of manganite with free access of air. Pyrolusite is often found with a manganite core, showing that the process is but partially completed.

#### SUMMARY

*Manganite*.— $\text{Mn}_2\text{O}_3 \cdot \text{H}_2\text{O}$ ;  $\text{Mn}_2\text{O}_3 = 89.7$  per cent,  $\text{H}_2\text{O} = 10.3$  per cent. Orthorhombic.  $a:b:c = 0.844:1:0.545$ ; (001), (111), (011), (101), (110), (010), (210), (120), (021); twinning parallel (011); cleavage parallel (010) perfect; brittle; fracture uneven.

Hardness=4; gravity=4.4. Steel gray; streak reddish black; sub-metallic; opaque.

Infusible; soluble in hydrochloric acid with evolution of chlorine.

Hartz Mountains, Michigan, Colorado.

#### Goethite

Goethite ( $\text{Fe}_2\text{O}_3 \cdot \text{H}_2\text{O}$ ), named in honor of the poet Goethe, who was interested in mineralogy as well as in other natural sciences, is an iron hydrate occurring in lustrous brown or black orthorhombic prisms terminated with pyramids (No. 547).

The usual planes (Fig. 125) are (110), (210), (010), (111), (011), and rarely (001). Prism planes are often striated. Columnar forms and capillary crystals radially grouped are common. The last are called "needle iron stone." Columnar and capillary crystals bunched together in radiated and concentric masses which end in rounded surfaces are said to be "reniform." Goethite also occurs in thin red scales composed of (100), (010), (401), (011) (Fig. 116). Multitudes of these fine scales attached on one side produce a mass with velvety luster. Because of their color they are called "ruby mica," and in

the finest scales are transparent, reddish yellow, and under the microscope dichroic, i.e., exhibit two different colors when the light traversing them is allowed to vibrate first in one and then in another direction through a dichroscope (see p. 96). Thus are they easily distinguished from scales of hematite, which are monochroic. Goethite crystals are common alteration products in secondary cavities, and give rise to a bronze sheen and opalescent tint. When goethite is heated it gives off water, becomes red, and changes to hematite.




FIG. 125.—Goethite




FIG. 126.—Goethite

Further heating with some reducing agent makes it black and magnetic, and heating continued until all the oxygen is removed produces pure iron. Goethite is much less abundant than either hematite or magnetite, but is a common associate of these and other iron ores in veins. Bohemia, Cornwall, Connecticut, the Lake Superior region, and Colorado yield the best crystals.

#### SUMMARY

*Goethite*.— $\text{Fe}_2\text{O}_3 \cdot \text{H}_2\text{O}$ ;  $\text{Fe}_2\text{O}_3 = 89.9$  per cent,  $\text{H}_2\text{O} = 10.1$  per cent. Orthorhombic.  $a:b:c = 0.913:1:0.607$ ; (110), (210), (011), (111), (001), (401), (010), (100). Cleavage parallel (010) perfect; brittle; fracture uneven.

Hardness=5; gravity=4. Brown to black; translucent; double refraction positive; dichroic.

Fusible with difficulty to magnetic bead; soluble in hydrochloric acid.

Bohemia, Cornwall, Connecticut, Michigan, Colorado.

**Limonite**

Much more common than goethite is the fibrous, dense, or earthy iron hydrate, limonite ( $2\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$ ) named from *λειμών*, the Greek for a moist grassy place, since it is found as a brownish-yellow deposit in bogs. It causes the iridescent slime seen in sluggish streams and pools, replacing the decaying vegetable matter. The rusting of iron is simply a change to limonite. There are several varieties founded upon form, origin, and condition of the mineral in deposits. First, there is the fibrous, radial, curvilaminar limonite, often with black glazed (No. 1881) or opalescent lustrous surface (No. 3359), lining cavities and geodes, and hanging in stalactites in caves (Nos. 2571 and 244); second, dense compact limonite (No. 298), in veins where it has been deposited by circulating waters which have gathered it from the surrounding decomposing rocks; third, extensive beds formed by waters circulating above ground and emptying into ponds. These beds are often oölitic (No. 1747), i.e., composed of myriads of small grains, among which are found fragments of algae, foraminifera, bryozoans, etc. Such deposits are analogous to the fourth variety—bog iron ore, forming today in swamps and making granular, nodular, concretionary, earthy, or sandy masses. On the bottom of many lakes is a black mud from which small grains of limonite are separating. Measurements made in Sweden show that deposits six inches thick have been formed in twenty years. The fifth variety consists of grains as large as a pea (pisolitic). The grains often show concentric structure and fill clefts in limestone and are cemented together in clumps. Sixth, constantly associated with the denser limonite and other iron ores is a yellowish-brown, soft, porous mass called yellow ochre. It is porous because it is a remnant left after the dissolution of other materials.

That limonite is a secondary mineral derived from such minerals as siderite and also from pyrite, hematite, and magnetite, is evident because the form of the original crystal is often retained—the rhombohedrons of siderite, cubes of pyrite, octahedrons of magnetite, and hexagonal plates of hematite. Different stages of the process show different proportions of the original crystals still unaffected. All the steps of the transition from original to derived material can be traced.

The brown streak of limonite, its inferior hardness and weight, and the presence of water distinguish it from hematite. It is



harder and lighter than the crystallized goethite and contains more water.

Because of the ease with which limonite fuses, it was probably the first mineral to be used by man as a source of iron.

#### SUMMARY

*Limonite*.— $2\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$ ;  $\text{Fe}_2\text{O}_3 = 88.5$  per cent,  $\text{H}_2\text{O} = 14.5$  per cent. Amorphous, fibrous, concentric, dense, earthy.

Hardness = 5.5; gravity = 3.8. Dark brown; streak yellow brown; submetallic; opaque.

Fusible to magnetic bead; soluble in hydrochloric acid.

Scotland, Sweden, Connecticut, New York, Pennsylvania, Alabama, Ohio, Illinois.

#### D. ALUMINITES, FERRITES, MANGANITES, CHROMITES

The chief minerals of this group, all of which crystallize in the regular system with the octahedron as a common form, are

Spinel	$\text{MgO} \cdot \text{Al}_2\text{O}_3$
Manganite	$\text{FeO} \cdot \text{Fe}_2\text{O}_3$
Franklinite	$(\text{Fe}, \text{Zn}, \text{Mn})\text{O} \cdot (\text{Fe}, \text{Mn})_2\text{O}_3$
Chromite	$\text{FeO} \cdot \text{Cr}_2\text{O}_3$

#### Spinel

Spinel is a mineral useful as a gem because of its beauty and hardness. The minute quantities of various elements which replace a part of the Mg or Al in the typical formula  $\text{MgO} \cdot \text{Al}_2\text{O}_3$  produce different colors. For example, the dark green to black opaque spinel (ceylonite) contains Fe. The yellowish- or greenish-brown variety (picotite) contains Fe and Cr, and the grass-green variety (chlor-spinel) contains Fe and Cu. But the typical spinel approaching most nearly the formula  $\text{MgO} \cdot \text{Al}_2\text{O}_3$  is of a beautiful clear red color, generally transparent, and called precious spinel. The purest red is called ruby spinel; the orange red, rubicelle; and the violet, the almandine spinel.

From earliest times precious spinel was prized as a gem, but was not distinguished from ruby until Romé de l'Isle studied it (1783), though these minerals differ in crystal form, cleavage, optical properties, hardness, and density. The specific gravity of spinel is 3.5, while that of ruby is 4; spinel is only 8 in hardness, while ruby is 9.

Spinel shows no pleochroism and is isotropic, i.e., it allows the light to pass through it similarly in all directions, as would be expected of a mineral crystallizing in the regular system.

In gem-bearing sands of Ceylon, Burma, and Siam, which have long been the source of precious spinel, are found small, sharp-edged octahedrons and typical spinel twins, where the octahedral face is the twinning plane (Fig. 127). The corners of the octahedrons are




FIG. 127.—Spinel twin




FIG. 128.—Spinel

often beveled by trapezohedrons and the edges by dodecahedrons, giving the crystal a rounded appearance. Dark-colored varieties occur in abundance at Vesuvius, in New York, New Jersey, and North Carolina.

#### SUMMARY

*Spinel*.— $\text{MgO} \cdot \text{Al}_2\text{O}_3$ ;  $\text{MgO}$ =28.2 per cent,  $\text{Al}_2\text{O}_3$ =71.9 per cent. Regular; holosymmetric; (111). Cleavage imperfect parallel (111); brittle; fracture conchoidal.

Hardness=6; gravity=3.5. Red, yellow, green, black; streak white; luster vitreous; transparent; refraction high,  $n=1.715$ .

Infusible; soluble with difficulty in sulphuric acid.

Burma, Ceylon, Appalachian region.

#### Magnetite

The third mineral in importance as a source of iron is magnetite, which derived its name, according to Pliny, from the shepherd Magnes, who found his iron-pointed staff attracted by the mineral while he was wandering over Mount Ida. It is the most magnetic of all

minerals, sometimes possessing polarity and attracting particles of iron to itself (loadstone) (No. 334). Usually it is simply itself attracted by a magnet.




FIG. 129.—Magnetite

Because of its magnetism it is easily separated from the sands of the ocean or lake or the streams in which it is found in abundance. In various sedimentary deposits in igneous and in metamorphic rocks it occurs as grains, granules, and masses.

Crystals of magnetite show most commonly octahedral (No. 3346) and dodecahedral forms in which the dodecahedron is striated parallel to the octahedral edges (No. 548), because of oscillatory combination (Fig. 129). The magnetism and the black streak of magnetite distinguish it from hematite.

#### SUMMARY

*Magnetite*.— $\text{FeO} \cdot \text{Fe}_2\text{O}_3$ ;  $\text{FeO}$ =31 per cent;  $\text{Fe}_2\text{O}_3$ =69 per cent. Regular; holosymmetric; (111), parting, parallel (111); brittle; fracture uneven.

Hardness=6; gravity=5.8. Black; streak black; metallic; opaque. Magnetic, sometimes polar.

Fusible with difficulty; powder easily soluble in hydrochloric acid.

Scandinavia, Urals, Altai Mountains, New York, Pennsylvania, New Mexico, North Carolina.

#### Franklinite

Franklinite closely resembles magnetite in form, color, hardness, and weight, but has a browner streak, is more commonly rounded on its octahedral edges, and is but slightly magnetic. Its usual association with the red zinc oxide (No. 3348), zincite, renders its determination by physical means less difficult, but chemical test (search for a zinc incrustation on charcoal or the amethystine color of manganese in the borax bead) is necessary for its accurate determination.

The mineral receives its name from Franklin Furnace, New Jersey, where it has been found in great quantities.

#### SUMMARY

*Franklinite*.— $(\text{Fe}, \text{Zn}, \text{Mn})\text{O} \cdot (\text{Fe}, \text{Mn})_2\text{O}_3$ . Regular; holosymmetric; (111); rounded grains. Resembles magnetite in physical properties, but is slightly magnetic and browner in streak.

Hardness = 6; gravity = 5.

Infusible, soluble in hydrochloric acid.

Franklin Furnace, New Jersey.

### Chromite

Chromite resembles magnetite and franklinite in form and color (No. 543), but is slightly softer (hardness 5.3) and lighter (gravity, 4.5). The best means of identifying it is to test for the green color which it gives to a cold borax bead.

Chromite owes its importance to the fact that it furnishes practically all the chromium used in the arts and manufactures. Chromium compounds are used to color porcelains and enamels green, and to dye calicoes, etc. Their most important use of late years, however, has been to harden steel. Before the world-war nearly a million dollars' worth of chromite was imported annually, a few thousand dollars' worth only being produced in this country. As a result of government investigation and encouragement production of domestic chromite was greatly increased. It is found in rocks consisting chiefly of olivine and serpentine.

### SUMMARY

*Chromite*.— $\text{FeO} \cdot \text{Cr}_2\text{O}_3$ ;  $\text{FeO}$  = 32 per cent,  $\text{Cr}_2\text{O}_3$  = 68 per cent. Regular; holosymmetric (111); granular, massive; uneven; fracture brittle.

Hardness = 5.5; gravity = 4.5. Black, yellowish red in very thin sections; dark brown.

Infusible; insoluble in acids, decomposing when fused with sodium sulphate.

New Caledonia, Bohemia.

### E. DIOXIDES

The minerals in this group contain two atoms of oxygen to one of the basic element. Those which most merit attention are:

Cassiterite

$\text{SnO}_2$

Rutile

$\text{TiO}_2$

Pyrolusite

$\text{MnO}_2$

### Cassiterite

This mineral, the only important source of tin, has been known since earliest times and was used by the ancients to make bronze. In color it is usually dark brown or black. It is hard (hardness, 6.5),

heavy (gravity, 7), insoluble, and infusible, and is usually in the form of rounded grains and pebbles or short stout crystals.

The color of cassiterite depends upon impurities such as iron oxide ( $\text{Fe}_2\text{O}_3$ ), tantalum oxide ( $\text{Ta}_2\text{O}_5$ ), etc. Pure varieties, which are rare, are colorless, transparent, and lustrous, and, were they a little harder, would be much prized for gems. In Mexico yellowish varieties are found, and Australia has yielded some fine red specimens; but most cassiterite is black.

Owing to its hardness, weight, and stability, it occurs in stream deposits as "stream-tin" and has been successfully mined in the Malay Peninsula, Australia, the Black Hills, and California. In primary deposits it is persistently associated with certain acidic




FIG. 130.—Cassiterite




FIG. 131.—Cassiterite twinned on (011)

igneous rocks, such as granites and pegmatites, where it has crystallized in short, stout tetragonal crystals, usually twinned. Simple crystals (Fig. 130) composed of (111), (110), (100), (101) are rarer than the twin forms which are so characteristic. The twinning plane is parallel (011), as shown in Figure 131. Prism planes are usually striated parallel to  $c$ . Basal planes are almost unknown. Slender prisms, having acute, ditetragonal pyramids such as (321) in addition to the more usual pyramids, occur rarely and are called "needle tin." Some of the massive varieties have a radiated fibrous structure, are arranged in curvilaminar layers of different shades, and in a botryoidal surface lack the luster of the ordinary crystal. Being remarkably dull and wooden, they are called "wood tin." Little brown-banded spherical nodules with the same fibrous structure are called "toad's

eye tin" in Cornwall, which has long been the most productive tin region.

The minerals associated with cassiterite suggest its origin. They are commonly apatite, fluorite, zinnwaldite, topaz, and tourmaline—all of which contain fluorine and lead to the thought that vapors containing fluorine were influential in the deposition of cassiterite. Daubree produced cassiterite artificially by the action of steam on tin fluoride. But cassiterite is produced in two other ways also. Violet-colored simple crystals have been made in tin works by the oxidation of metallic tin, and cassiterite has been found replacing organic remains and cementing nodules, as would be the case were it deposited from solution, or from vapors.

The chief uses of cassiterite are as a source of tin for plating and the manufacture of alloys.

#### SUMMARY

*Cassiterite*.— $\text{SnO}_2$ ; Sn=78.6 per cent, O=21.4 per cent. Tetragonal; holosymmetric.  $a:c=1:0.672$ . (111), (100), (110), (101), (210), (321); twinned on (101); cleavage parallel (100) imperfect; brittle; fracture sub-conchoidal.

Hardness=6.5; gravity=7. Brown or black; streak gray; adamantine; translucent;  $\omega=1.997$ ; double refraction positive;  $\epsilon-\omega=0.097$ . Insoluble; with soda on charcoal yields tin.

Cornwall, Malay Peninsula, Wyoming, and Dakota.

#### Rutile

Rutile ( $\text{TiO}_2$ ) is a source of titanium, an element used for giving a yellow color to glass, for hardening steel, and for various chemical purposes. Its hardness is the same as that of cassiterite (6.5), and its color and form are very similar, but it is redder (*rutilus*, Latin for "red") and has a yellowish-brown streak instead of a grayish streak. It is only 4.3 in specific gravity and cleaves readily to (100) and (110), hence is easily distinguished from cassiterite. It occurs in stout crystals (Nos. 3354 and 3355), in acicular and twin crystals, and in masses (No. 3197). The stout crystals (Fig. 132), nearly duplicating those of cassiterite, consist of the following planes: (100), (110), (011), all of which may be vertically striated, and (111) and (101).




FIG. 132.—Rutile

Twinning parallel to (011) is very common (Fig. 133) and the twins are often repeated six or eight times till they form a complete ring with the different individuals inclined to each other in a zigzag with angles of  $65^{\circ} 35''$  (Fig. 134).

Acicular crystals varying from the finest threads to needles and blades of some thickness often penetrate other minerals such as quartz.




FIG. 133.—Rutile triplet twinned on (011).




FIG. 134.—Rutile octet twinned on (011).

The beautiful yellowish-red or brown fibers in quartz are called *fleches d'amour*. In some groups the needles cross each other at the twinning angle and form a reticulated skeletal plate called "sagenite" (= net).

#### SUMMARY

**Rutile.**— $\text{TiO}_2$ ; Ti=60 per cent, O=40 per cent. Tetragonal; holosymmetric.  $a:c=1:0.644$ . (100), (110), (310), (111), (101); twinned on (101); cleavage, parallel (100), (110); brittle; fracture uneven.

Hardness=6.5; gravity=4.3. Reddish brown; streak yellowish brown; metallic; adamantine; translucent;  $\omega=2.616$ ; double refraction positive strong,  $\epsilon-\omega=0.287$ .

Infusible; insoluble.

Switzerland, Virginia, North Carolina, Florida, Arkansas, Alaska.

#### Pyrolusite

Pyrolusite ( $\text{MnO}_2$ ) is an amorphous, black, soft (hardness, 2) mineral used in glass manufacture to clear the glass from green and brown colors (Nos. 541 and 1838). Because of its usefulness in this

respect it has received its name ( $\pi\upsilon\rho$ , "fire";  $\lambda\upsilon\epsilon\iota\nu$ , "to wash"). Large quantities are employed also as a flux in iron manufacture.

It has no crystal form of its own, but borrows its form of manganite, from which it is derived by the loss of water. Brazil and Russia before the war supplied the United States with the greatest part of the manganese ore needed. About one million tons of ore came from Russia in 1913. In 1916 more than that amount was produced in the United States.

#### SUMMARY

*Pyrolusite*.— $\text{MnO}_2$ ; Mn=63.2 per cent, O=36.8 per cent. Pseudomorph after manganite, showing radiated fibrous structure, but usually massive, earthy, soiling the fingers.

Hardness=2; gravity=5. Gray to black; streak black.

Infusible; soluble in warm hydrochloric acid.

Minnesota, Arkansas, California, Virginia, Russia, Brazil.



## CLASS VI. CARBONATES

### CALCITE GROUP

CALCITE GROUP	HEXAGONAL
Calcite	$\text{CaCO}_3$
Dolomite	$\text{CaMg}(\text{CO}_3)_2$
Magnesite	$\text{MgCO}_3$
Siderite	$\text{FeCO}_3$
Rhodochrosite	$\text{MnCO}_3$
Smithsonite	$\text{ZnCO}_3$

#### Calcite

Calcite is one of the most important and interesting minerals in the world, both because of its beauty and abundance, and because of its usefulness from a scientific and practical standpoint. The history of calcite is the history of mineralogy. In abundance it is surpassed by quartz alone. Its crystals occur in such profusion, variety, and beauty as easily to have attracted the attention of mineralogists and to have continually furnished them with material for study. This study has led to important results. About the time that the Dane, Steno, noted the regularity of the angles on quartz and announced the law of the constancy of angle, a countryman of his, Erasmus Bartholinus (1670), was working with the splendid calcite crystals then recently discovered in Iceland (No. 3832); and in his book *Experimenta Crystalli Islandici* described the remarkable cleavage and the double refraction which calcite shows more satisfactorily than does any other mineral.

Twenty years later the Hollander Huygens, famous for his undulatory theory of light, extending Bartholinus' study of calcite, was able to formulate the laws of double refraction—the laws of a phenomenon which could not be explained by the corpuscular theory of Newton. For many years following, while discussion of the corpuscular and wave theories of light was at its height, calcite was carefully studied by the advocates of both theories. As the result of such study Malus (1808) discovered the polarization of light. Today calcite is much used in optical researches because of its effect on light, being employed for “nicol prisms” in microscopes, both for purely scientific and for commercial purposes.





Calcite, Webb City, Missouri; twinned on (100)

PLATE XXI



*a*, Calcite, "dog-tooth spar," Joplin, Missouri



*b*, Calcite, "Iceland spar," showing double refraction



No mineral shows more planes and combinations of planes than does calcite. More than two hundred forms and seven hundred combinations have been described. There are four distinct habits of crystallization—rhombohedral, scalenohedral, prismatic, and tabular. The fundamental form is the rhombohedron,  $R$  ( $10\bar{1}1$ ), (Fig. 135), in which the mineral always cleaves, and so readily that it is difficult to produce a fracture in any other direction (Nos. 3460 and 3832). As an independent form this plane is rare but is found on crystals from near Bologna, Italy, and is a predominant form on the calcite from Iceland ("Iceland spar"). The obtuse rhombohedron,  $-\frac{1}{2}R$  ( $01\bar{1}2$ ) (Fig. 139), is common. Figure 137 represents another common acute rhombohedron. The scalenohedron which furnishes the so-called "dog




FIG. 135.—Calcite. Positive rhombohedron ( $10\bar{1}1$ ) =  $R$ ; the cleavage rhombohedron.




FIG. 136.—Calcite. Negative rhombohedron ( $01\bar{1}1$ ) =  $-R$ .

tooth spar" (Fig. 138) (Nos. 3446, 3458, etc.) is a form of frequent occurrence. Prism planes also appear (No. 3450), modified usually with rhombohedron planes as in Figure 139, where the rhombohedron is negative,  $-\frac{1}{2}R$ . If the prisms are short and a basal plane is present, tabular crystals similar to Figure 140 result. Sometimes they are as thin as paper and grouped parallel to one another so as to give the effect of cleavage which is peculiar to slate, hence the variety is called "slate spar." Scalenohedrons are usually modified by rhombohedrons (Fig. 141).

All these forms agree in having three planes of symmetry, which are diagonal to the lateral crystallographic axes, and intersect in the vertical axis  $c$ , the axis of trigonal symmetry. Such symmetry is so typical as to have been named after calcite the "calcite class."

There are four types of twins: (1) A common type is that in which two crystals are united by juxtaposition on the basal plane. Figure 142 shows a rhombohedron and Figure 143 a scalenohedron twinned according to this law. If the crystals overlap, filling the




FIG. 137.—Calcite. Negative acute rhombohedron ( $02\bar{2}1$ ) =  $-2R$ .




FIG. 138.—Calcite, scalenohedron.




FIG. 139.—Calcite, prism and negative obtuse rhombohedron.




FIG. 140.—Calcite, showing prism, negative obtuse rhombohedron, and base.




FIG. 141.—Calcite; combination of scalenohedron and rhombohedron.




FIG. 142.—Calcite. Rhombohedron twinned on ( $0001$ ).

re-entering angles, cleavage lines will disclose the twinning. (2) More common than the foregoing is that type whose twinning plane is  $-\frac{1}{2}R$  ( $01\bar{1}2$ ). In this case the cleavage planes of the two individuals are parallel. Figure 144 shows a juxtaposed twin of this sort characteristic of crystals from Guanajuato, Mexico. Twinning lamellae





PLATE XXII



Calcite, Joplin, Missouri;  $(21\bar{3}1)$  and  $(31\bar{4}5)$

PLATE XXIII



Calcite scalenohedron, Rossie, St. Lawrence County, New York



parallel to  $-\frac{1}{2}R$  have been commonly produced in calcite by pressure, and in thin sections under the microscope are so characteristic as to furnish the best means of distinguishing the mineral. They well illustrate secondary twinning such as may be artificially produced in calcite and is especially pronounced in "Iceland spar." (3) The type of twinning parallel to the cleavage rhombohedron  $R$ , though rare, is shown in scalenohedrons and prisms (Figs. 145 and 146). In these twins one cleavage plane only is parallel to the two individuals.




FIG. 143.—Calcite scalenohedron twinned parallel to  $\{0001\}$ .




FIG. 144.—Calcite scalenohedron twinned parallel to  $\{0112\} = -\frac{1}{2}R$ .




FIG. 145.—Calcite prism twinned parallel to  $R$ .

(4) The type where the twinning plane is the acute negative rhombohedron  $\{02\bar{2}1\}$  produces forms which closely resemble those of the second class, but here the cleavage planes of the different individuals are not parallel (Fig. 147).

Some calcite crystals exhibit asterism, i.e., when a candle flame is viewed through them it appears as a radiating star of light. This is due to systems of hollow tubes parallel to each other in three directions, and is produced where the "gliding surfaces" of the negative rhombohedron  $-\frac{1}{2}R$  intersect each other.

Calcite is useful for nicol prisms because the ordinary ray while passing through it is so greatly refracted ( $\omega = 1.658$ ). The extraordinary ray is allowed to pass through the prism, being but slightly affected by the Canada balsam whose index is nearly that of the extraordinary ray. (For balsam  $\epsilon = 1.536$ .)

A microscopic section of calcite rotated above the polarizer when the analyzer is removed shows high relief if the ordinary ray is allowed to pass through, and relief so low as to be almost invisible when the extraordinary ray passes through.

When calcium carbonate crystallizes from aqueous solution in veins or other cavities, it forms the ordinary variety of calcite. If it is deposited from springs or streams by evaporation in a more or less granular condition it forms travertine, calc tufa, stalactites, and stalagmites. If it is composed of fragments or organic remains cemented by calcareous or other cements, it forms chalk, oölite, and




FIG. 146.—Calcite scalenohedron twinned parallel to  $R$ .




FIG. 147.—Calcite scalenohedron twinned parallel to  $(02\bar{2}1)$ .

limestone. If the limestone has been metamorphosed by heat and pressure so as to become crystallized, it forms marble. Among some of the localities the following are famous because of the abundance and beauty of their crystallized varieties. In Iceland near Eskifjörður a cavity 36 feet long, 15 feet wide, and 10 feet high in dolomite rock was found filled with clear crystallized calcite. The prevailing forms were rhombohedrons  $(10\bar{1}1)$  with edges beveled by scalenohedrons  $(21\bar{3}1)$  and  $(3145)$ , and scalenohedrons terminated by  $(10\bar{1}1)$  or  $(3145)$ . Their surfaces were often corroded or coated with other minerals such as stilbite.

In England the lead, iron, and fluorite mines of Derbyshire, Dunham, and Cumberland (Nos. 3450, 3451, and 3452) have furnished fine crystals which now ornament museums in all parts of the world. Many beautiful crystals come from the Hartz Mountains. They are



PLATE XXIV



*a*, Calcite, Joplin, Missouri;  $(21\bar{3}1)$  and  $(31\bar{4}5)$



*b*, Quartz geode with large flat rhombohedral crystals, St. Francisville, Missouri

commonly prismatic planes and tabular forms. The silver mines of Guanajuato, Mexico, have furnished twin crystals of great beauty and variety. Among many famous localities in the United States may be mentioned St. Lawrence County, New York (No. 4657); the Lake Superior copper mines with their complex crystals which often contain spangles and wires of copper; and the Wisconsin, Illinois (Nos. 698 and 699), and Missouri (No. 3459, etc.) lead and zinc mines with their rhombohedrons and scalenohedrons. The geodes of Keokuk contain numerous large flat crystals (Nos. 686, 691, 672). At Joplin, Missouri, many large, beautiful, honey-yellow scalenohedrons ( $21\bar{3}1$ ) terminated by rhombohedrons  $R$  ( $10\bar{1}1$ ) and the striated  $-\frac{1}{2}R$  have been found. The acute terminal edges of these scalenohedrons ( $21\bar{3}1$ ) are often replaced by striated and rounded faces (Nos. 3886, 3899, 3890, also Plate XXIII).

#### SUMMARY

*Calcite*.— $\text{CaCO}_3$ ;  $\text{CaO}$ =56 per cent,  $\text{CO}_2$ =44 per cent. Hexagonal; symmetry dihexagonal alternating (calcite class);  $a:c=1:0.854$ .  $R$ ,  $-\frac{1}{2}R$ ,  $4R$ ; twinned on (0001), (0112), (1011), (0221). Cleavage parallel  $R$  perfect; brittle; fracture conchoidal.

Hardness=3; gravity=2.72. Colorless; vitreous; transparent; refraction strong,  $\omega=1.658$ ; double refraction very strong, positive,  $\omega-\epsilon=0.172$ .

Infusible; soluble with effervescence in cold hydrochloric acid, diluted to one-third strength.

Ubiquitous.

#### Dolomite

Dolomite can be distinguished from calcite, which it very closely resembles, from the fact that it is harder (hardness, 3.5), heavier (gravity, 2.85), and does not effervesce in cold hydrochloric acid diluted to one-third strength, except when finely powdered. Though strongly resembling each other in crystal form, calcite and dolomite differ in this respect, that while calcite has dihexagonal alternating symmetry, dolomite has hexagonal alternating symmetry (dioptase class), i.e., it lacks all planes of symmetry, and the vertical axis is an axis of trigonal symmetry. This becomes evident when the two minerals are etched with acid and when their axes of elasticity are measured. If rhombohedrons of calcite and dolomite are placed in dilute hydrochloric



acid, upon the surface appear depressions which show the symmetry of the crystals. The depressions on calcite (Fig. 148) indicate three planes of symmetry, since each etched figure has one line of symmetry parallel to the shorter diagonal of the rhombohedron, showing that a plane of symmetry is perpendicular to that face. A dolomite rhombohedron treated in the same manner is marked with pits unsymmetrical in outline (Fig. 149), indicating that there is no plane of symmetry perpendicular to the rhombohedron. The figures on the three upper faces are related to each other as are a right-handed and a left-handed glove, the lower ones appearing as if they were reflections of the upper (enantiomorphous). This indicates that the crystal has one hexagonal axis of alternating symmetry, a




FIG. 148.—Calcite etched with dilute hydrochloric acid.




FIG. 149.—Dolomite etched with dilute hydrochloric acid.

fact which is also shown by rhombohedron planes which are sometimes developed upon the alternate edges of the usual rhombohedron.

The difference in the symmetry of calcite and dolomite is also indicated by their coefficients of elasticity. These coefficients are obtained by cutting bars of the minerals, supporting them on knife edges, applying a weight sustained by a knife edge, and measuring the amount of bending by microscopic or other means. When lines are drawn on a rhombohedral face proportional to the amount of bending and the ends connected, curves shown in Figures 150 and 151 are produced. In calcite the elasticity is symmetrically arranged parallel to the diagonal of the rhombohedron. In dolomite it is unsymmetrical. Thus etching and measuring of elasticity show that the rhombohedrons are not perpendicular to planes of symmetry in dolomite. Further, in calcite it was seen that  $\frac{1}{2}R(01\bar{1}2)$  is a glide plane, or plane of secondary twinning, as shown by the series of

hollow tubes arranged parallel to this plane and appearing as fine lamellae under the microscope. The presence of this glide plane can be discovered by pressing a knife into a cleavage rhombohedron across one of the terminal edges. By the pressure the molecules are revolved  $180^\circ$  into a new twinning plane, so that the other lamellae are parallel to  $-\frac{1}{2}R$  (Fig. 152). In dolomite  $-\frac{1}{2}R$  is not a plane of secondary twinning.

Dolomite is a double salt of calcium and magnesium—a molecule of each carbonate being united to form it. If it were an isomorphous mixture of the two carbonates in molecular proportions, its crystallization would be the same as that of calcite and magnesite. However, it is different, and it would have a specific gravity of 2.843. But it




FIG. 150.—Elasticity coefficient of calcite.




FIG. 151.—Elasticity coefficient of dolomite.

is heavier (gravity, 2.85), just as would be expected, since while the double salt is forming there is a contraction of the two carbonates which increases their specific gravity.

Dolomite occurs in well-crystallized forms deposited from solution and in masses made of fragments of organic remains which have been more or less altered and cemented by chemicals in solution. The massive variety forms extensive beds which extend for miles over the country.

Vermont, New Jersey, and New York (No. 3217, etc.) furnish many crystals. Saddle-shaped crystals are abundant in Joplin, Missouri (No. 3464). The greater part of the limestone of Illinois is dolomitic.




FIG. 152.—Glide planes in calcite.

## SUMMARY

*Dolomite*.— $\text{CaMg}(\text{CO}_3)_2$ ;  $\text{CaO}=30.4$  per cent,  $\text{MgO}=21.7$  per cent,  $\text{CO}_2=47.9$  per cent. Hexagonal; symmetry hexagonal alternating;  $a:c=1:0.832$ . ( $10\bar{1}1$ ); cleavage parallel ( $10\bar{1}1$ ) perfect; brittle; fracture conchoidal.

Hardness=3.5; gravity=2.85. Colorless, streak white; vitreous; transparent. Refraction strong,  $\omega=1.682$ ; double refraction strong, negative,  $\omega-\epsilon=0.189$ .

Infusible; soluble with effervescence in warm acids.

Tyrol, Switzerland, England, Vermont, New Jersey, New York, Missouri.

## Magnesite

The pure magnesium carbonate is a white brittle mineral, usually massive, granular, and earthy. It is harder than either of the two other members of the group thus far described (hardness, 4) and heavier (gravity, 3.1). Before the world-war this mineral, obtained mainly from Greece (No. 3466), furnished a large part of the magnesium needed in the arts and manufactures. Recently magnesite from Quebec, California, etc., has been used; also dolomite and the waters from which sodium chloride had been extracted.

## SUMMARY

*Magnesite*.— $\text{MgCO}_3$ ;  $\text{MgO}=47.6$  per cent,  $\text{CO}_2=52.4$  per cent. Hexagonal;  $a:c=1:0.8112$ . Massive, granular, earthy; brittle; fracture sub-conchoidal. White; vitreous; silky; transparent to opaque.

Infusible; effervesces in warm hydrochloric acid. Common decomposition product of ferromagnesian silicates.

Greece, Canada, California, Washington, Maryland.

## Siderite

Siderite,  $\text{FeCO}_3$ , which furnishes almost no iron in the United States, is an important iron ore in Germany (No. 4512), and the most important source of iron in England. When pure it occurs in brown, vitreous, translucent rhombohedrons, or in fibrous botryoidal or globular forms. The rhombohedrons are often curved and sometimes acute. Large basal planes give rise to a tabular variety which often has zonal structure caused by hexagonal bands. Being liable to oxidation, the mineral readily loses its gray color, becomes brown,

and changes into limonite. The dehydration of limonite produces hematite and finally magnetite.

Among European localities, Cornwall and Freiburg are the most productive of siderite, while in the United States the Appalachian regions have furnished the largest supplies. Of recent years Ohio has been the leading producer.

#### SUMMARY

*Siderite*.— $\text{FeCO}_3$ ;  $\text{FeO}=62.1$  per cent,  $\text{CO}_2=37.9$  per cent. Hexagonal; symmetry, dihexagonal alternating;  $a:c=1:0.818$ .  $(10\bar{1}1)$ ,  $(0001)$ ; faces curved; fibrous, globular. Cleavage parallel  $(10\bar{1}1)$  perfect; brittle; fracture sub-conchoidal.

Hardness=3.5; gravity=3.8. Brown; vitreous, translucent. Refraction very strong,  $\omega=1.878$ ; double refraction very strong, negative  $\omega-\epsilon=0.241$ .

Fuses at 4.5 to black magnetic globule. Effervesces in warm hydrochloric acid.

Cornwall, Freiburg, Ohio, Appalachians.

#### Rhodochrosite

Beautiful rose-pink rhombohedrons of rhodochrosite are found in, Colorado (Nos. 3471 and 3472), the Ural Mountains, and other places where solutions carrying manganese carbonates have for some cause slowly given up their burden. The reddish color of rhodochrosite easily distinguishes it from the other carbonates.

#### SUMMARY

*Rhodochrosite*.— $\text{MnCO}_3$ ;  $\text{MnO}=61.7$  per cent,  $\text{CO}_2=38.3$  per cent. Rounded  $(10\bar{1}1)$ , massive, compact. Cleavage parallel  $(10\bar{1}1)$  perfect; brittle; fracture uneven.

Hardness=4; gravity=3.5. Rose red; translucent; negative.

Infusible. Effervesces in warm hydrochloric acid.

Russia, Hungary, Saxony, Belgium, New Jersey, Colorado.

#### Smithsonite

The physical condition, that is, the form, cleavage, fracture, hardness, weight, luster, diaphaneity, and optical properties of the zinc carbonate, smithsonite, closely resemble that of the other members of the group. Smithsonite is sometimes colorless, but more

usually green (No. 3751), blue, or brown from the presence of copper, iron, or other foreign substances. It is used as a zinc ore in the Mississippi Valley region, as well as in other places. In northern parts of Illinois it formerly produced upward of a thousand dollars' worth of zinc annually.

#### SUMMARY

*Smithsonite*.— $\text{ZnCO}_3$ ;  $\text{ZnO}$ =64.8 per cent,  $\text{CO}_2$ =35.2 per cent. Usually botryoidal, reniform, granular, earthy.

Hardness=5; gravity=4.4. White, green, blue, brown, vitreous.

Infusible; gives zinc coating with soda on charcoal. Effervesces in warm hydrochloric acid.

Many European localities, Wisconsin, Illinois, Iowa, Missouri, Arkansas.

The calcite group furnishes one of the best illustrations of isomorphism which the mineral kingdom affords, since the carbonates of calcium, magnesium, iron, manganese, and zinc, all different chemical substances, assume practically the same form. All the members of the group are rhombohedral in form, practically identical in cleavage, very similar in hardness and gravity. All effervesce in warm hydrochloric acid. The following group, the aragonite group, all of whose members are orthorhombic, is another illustration of isomorphism.

#### ARAGONITE GROUP

##### ARAGONITE GROUP

Aragonite  
Witherite  
Strontianite  
Cerussite

##### ORTHORHOMBIC

$\text{CaCO}_3$   
 $\text{BaCO}_3$   
 $\text{SrCO}_3$   
 $\text{PbCO}_3$

#### Aragonite

The orthorhombic form of calcium carbonate, named from Aragon in Spain where it was first found, is much less common than calcite. Its comparative rarity may be due to two causes: (1) to the conditions necessary for its formation; and (2) to its instability. One of the conditions necessary for its formation is that the solution from which it is deposited must be hot, whereas calcite is usually deposited from cold waters. This is shown when the two minerals are made in the laboratory and by the conditions which surround aragonite



PLATE XXV



*a*, Aragonite crystals four inches in diameter, Cianciana, Sicily



*b*, Stalactites, Bisbee, Arizona

in the field and by its associations. Aragonite very often accompanies sulphur crystals, which are commonly deposited in volcanic regions from hot solutions. There are exceptions, such as the calcium carbonate deposited by living organisms in the shells of mollusks, which is in the form of aragonite. Further, if a calcium carbonate solution contains a minute quantity of a soluble sulphate or orthorhombic carbonate, aragonite crystals may be formed. Aragonite is less stable than calcite, readily changing its crystal form at ordinary temperature.

When one mineral changes into another of the same composition by simply altering its form, it is called a paramorph. When it changes into the form of a mineral of different composition it is called a pseudomorph. For example, a quartz crystal which assumes the cubic shape of fluorite is said to be a pseudomorph after fluorite. An aragonite crystal which takes the form of calcite is said to be a paramorph. A paramorph can be detected, for example, when an aragonite crystal is but partially paramorphosed, the inner portion being aragonite and the outer calcite. Paramorphism is possible only among minerals which exhibit dimorphism or polymorphism. Aragonite and calcite were the earliest recognized examples of dimorphism in the mineral kingdom.

Aragonite rarely occurs in simple orthorhombic crystals. It is nearly always twinned parallel to the prism in such a manner as to produce seemingly hexagonal forms. Nearly all calcium carbonate which is fibrous, stalactitic, botryoidal, or concretionary is classified as aragonite. For example, the stalactites of Mammoth Cave and other caves (Nos. 2118, 2119), the pisolites of Carlsbad and other hot springs, and the beautiful white coral-like groups from Wind Cave, South Dakota (No. 3176), and Bisbee, Arizona, are aragonite.

Aragonite can be distinguished from calcite when the crystal form is not evident by its superior hardness and weight.

Simple crystals (Fig. 153) are composed of prisms, brachypinacoids, and dome planes. Groups of crystals with predominance of acute pyramids (441), (991), and (081), (091), showing horizontal striations, are common. If three crystals such as shown in Figure 154




FIG. 153.—Aragonite



interpenetrate parallel to the prism plane and the re-entering angles are filled in, a form having the cross-section shown in Figure 155 and resembling a hexagonal prism is produced. Large, white, yellowish prisms of this sort are found at Girgenti and Cianciana, Sicily (No. 3902), and at the sulphur mines in Hungary. At Aragon, Spain, the crystals are corroded and are found in red ferruginous marl with gypsum and quartz. Nos. 2116 and 2120 show stalactites from Chester, Illinois; No. 2117, aragonite as fossilizing material at the same locality; Nos. 2672, 2685, examples from Rock Island; No. 3469




FIG. 154.—Basal section of aragonite triplet.




FIG. 155.—Basal section of aragonite; interpenetrant triplet.

shows botryoidal masses from Quincy; No. 3176 from the Black Hills; Nos. 3467 and 3468 are from Fort Collins. Carlsbad, Bohemia, is also represented. No. 3902 is an unusually fine group consisting of two large crystals of interpenetrating triplets and several smaller imperfect crystals. The largest crystal measures about four inches in diameter and is two inches high.

#### SUMMARY

*Aragonite*.— $\text{CaCO}_3$ ;  $\text{CaO}$ =56 per cent,  $\text{CO}_2$ =44 per cent. Orthorhombic.  $a:b:c=0.628:10.721$ . (110), (010), (011), (001), (111); twinned on (110); cleavage (010), (110) imperfect; brittle; fracture sub-conchoidal.

Hardness=3.5; gravity=2.9. Colorless; vitreous; transparent; mean angle of refraction,  $\beta=1.682$ , the least,  $\alpha=1.530$ . Double refraction very strong, negative, i.e., difference between the greatest angle of refraction,  $\gamma$ , and the least,  $\alpha$ , is 0.156.

Infusible; effervesces in hydrochloric acid.

Spain, Sicily, Cordilleran states.

**Witherite**

This carbonate of barium occurs in white, heavy, not very abundant crystals and masses. The crystals appear to be hexagonal pyramids (Fig. 156), but a thin section cut parallel to the base (Fig. 157) shows that the seemingly simple crystal is composed of three orthorhombic crystals, twinned parallel to prism planes and hence crossing each other at angles of  $62^\circ$ —the angle between the prism planes. Upon the pyramid planes are more or less prominent striations caused by the growth of a succession of different capping




FIG. 156.—Witherite



FIG. 157.—Cross-section of witherite

pyramids. The pyramid planes vary in their intercepts on the  $c$  axis and produce both acute and obtuse forms. In addition to well-formed crystals there are aggregates made up of acicular crystals, grouped into botryoidal and reniform shapes. Compact masses are the most characteristic form of the mineral.

The luster of the fresh crystals is greasy, but is changed when sulphur fumes or solutions coat the surface with barium sulphate, changing it into a dull white.

Witherite may be distinguished from minerals of similar appearance by the fact that it is heavy, effervesces in cold hydrochloric acid, and colors the blowpipe flame green. Great quantities are produced in two northern counties in England, where it was discovered in 1783 by the mineralogist after whom it was named, Withering, and where it has been mined for more than one hundred years. Witherite is used for medicinal and industrial purposes. Large quantities are employed in the manufacture of rat poison.

## SUMMARY

*Witherite*.— $\text{BaCO}_3$ ;  $\text{BaO}=77.7$  per cent,  $\text{CO}_2=22.3$  per cent. Orthorhombic; holosymmetric.  $a:b:c=0.603:1:0.730$ . (110), (010), (011). Common form pseudohexagonal bipyramid produced by interpenetration twinning of three individuals at angles of  $62^\circ$ ; twinning plane (110). Brittle; fracture uneven.

Hardness=3.5; gravity=4.3. Colorless; streak white; vitreous; translucent. Double refraction negative, weak; axial plane parallel (010); acute bisectrix perpendicular to (001).  $2E=26^\circ 30'$ .

Fusible (2); effervesces in hydrochloric acid.

Northumberland and Cumberland, England; Kentucky, Michigan.

## Strontianite

Strontianite very closely resembles aragonite in color, streak, luster, and form, but differs in being heavier (gravity, 3.7) and in yielding the intense red color characteristic of strontium when heated in the blowpipe flame.

Strontianite is sometimes colorless and transparent, but more often translucent and white, green, yellow, or brown. Its fibrous, acicular, or columnar crystals are rarely well defined or terminated. They are usually vertically striated. The same kind of twinning occurs as is so common for aragonite and witherite, viz., interpenetrant triplets forming pseudohexagonal prisms or pyramids.

At Strontian on the west coast of Scotland in 1791 a mineral was found that contained a new element. The element was named strontium and the mineral strontianite.

Strontianite finds limited use as a source of red lights for fireworks and in the refining of beet sugar.

## SUMMARY

*Strontianite*.— $\text{SrCO}_3$ ;  $\text{SrO}=70.1$  per cent,  $\text{CO}_2=29.9$  per cent. Orthorhombic; holosymmetric.  $a:b:c=0.6090:1:0.7239$ . (110), (010), (011); fibrous, acicular, columnar, granular; cleavage parallel (110) nearly perfect; brittle; fracture uneven.

Hardness=3.5; gravity=3.7. Colorless, white, green, yellow, brown. Optically negative; axial plane parallel to (100); bisectrix perpendicular to (001).  $2E=12^\circ 17'$ .

Fusible; soluble in hydrochloric acid.

Scotland, Appalachian states.

### Cerussite

Cerussite, like aragonite, witherite, and strontianite, simulates crystals of the hexagonal system, while in reality its molecules arrange themselves in accordance with the laws of the orthorhombic system. Simple crystals of cerussite are more abundant than are those of witherite and strontianite. The most common habit is the form produced by pyramids, domes, short prisms, and pinacoids (Fig. 158), and the tabular crystals like those in Figure 159, in which the domes are elongated parallel to the  $a$  axis and the pinacoid is the predominant plane.

The commonly occurring interpenetrant twinning of three crystals parallel to prism planes produces raylike pyramids (Fig. 160).




FIG. 158.—Cerussite




FIG. 159.—Cerussite




FIG. 160.—Three cerussite crystals interpenetrating parallel to prism planes.

Besides pyramids and tabular crystals, single and combined, acicular and fibrous masses are of common occurrence. All these crystals when fresh have smooth, bright, lustrous surfaces. Silky or adamantine luster is characteristic of lead minerals. The best specimens have been obtained from Bohemia, Hungary, New South Wales, and Idaho, where they are found on galena and other lead ores, from which they have resulted by the decomposition of galena.

### SUMMARY

*Cerussite*.— $\text{PbCO}_3$ ;  $\text{PbO}=83.5$  per cent,  $\text{CO}_2=16.5$  per cent. Orthorhombic; holosymmetric;  $a:b:c=0.610:1:0.723$ .  $(110)$ ,  $(010)$ ,  $(111)$ ,  $(021)$ ; twinned on  $(110)$ ; cleavage parallel  $(110)$ ,  $(021)$  imperfect; brittle; fracture conchoidal.

Hardness=3.5; gravity=6.5. Colorless; streak white; adamantine; transparent;  $\beta=2.076$ ,  $\alpha=1.804$ ; double refraction strong, negative;  $\gamma-\alpha=0.274$ ; axial plane (010); acute bisectrix perpendicular to (001).  $2E=17^{\circ}8'$

Fusible; soluble in nitric acid.  
Cordilleran states.

### Malachite

Malachite ( $\mu\alpha\lambda\acute{\alpha}\chi\eta$ , "mallow or willow tree") is a basic copper carbonate, which is readily recognized because of its vivid green color. Fine compact nodular masses composed of radial fibers have been found in such quantities in the Ural Mountains (No. 3407) and

so rarely in other European localities that they have furnished the rulers of Russia a unique and much-prized material for gifts. In palaces and museums in all the capitals of Europe the tourist sees vases, tables, and other ornaments made of this striking green mineral. They are usually recorded as "a gift from the Czar of Russia." In many European and American localities malachite (Nos. 3405, 3406) occurs in such quantities as to furnish a useful source of copper.




FIG. 161.—Axes of monoclinic crystal.

When crystallized under favorable conditions, in cavities, for example, it forms fine fibrous needles which build tufts as soft in appearance as velvet (velvet malachite). Radiating fibers fill winding cavities so as to look like roots of trees when exposed. The granular and earthy forms of malachite are the most abundant. Well-formed crystals are almost unknown, but acicular prisms disclose the fact that the molecules have so arranged themselves as to produce forms characteristic of the monoclinic system.

In this system the molecular structure is represented by three axes of unequal length (Fig. 161). The angle  $\beta$  which the  $a$  axis makes with the  $c$  axis is greater than  $90^{\circ}$ . The  $\alpha$  and  $\gamma$  angles are right angles. Planes constructed upon these axes produce figures having a plane of symmetry parallel to  $c$  and  $a$ , and an axis of symmetry

which is the  $c$  axis. The forms which constitute the system are analogous to those of the orthorhombic system. They are pyramids (111), prisms (110), orthopinacoids (straight pinacoids) (100), clinopinacoids (inclined pinacoids) (010), orthodomes (101), and clinodomes (011). As in the orthorhombic system, closed forms are obtained by combinations of two or more kinds of planes in all cases except that of the pyramid (Fig. 162). In Figure 163 orthodomes are united with clinodomes to produce a complete form. Figure 164 shows a combination of prism and base.

Malachite does not well illustrate the crystallography of the monoclinic system. To understand its crystals it is necessary to




FIG. 162.—Monoclinic bipyramid




FIG. 163.—Orthodomes and clinodomes

resort to the microscope. Indeed, with any transparent mineral employment of optical means of investigation contributes greatly to the knowledge of its crystallography. Sections of minerals cut at various angles are cemented with Canada balsam to pieces of glass and ground till thin enough (about one-hundredth of an inch in thickness) to permit light to pass through them. Examined under the microscope, first with the light vibrating in all directions, then with light made to vibrate in but one direction by means of a calcite prism ("Nicol prism"), and studied with light passing through the mineral with parallel rays and then with converging rays, the crystal structure becomes clear. Such an examination of malachite reveals the fact that there are two directions in which light is not doubly refracted, that these directions are in the plane (axial plane) parallel

to the clinopinacoid, that they form an angle of nearly  $90^\circ$  with each other ( $2 E=89^\circ 18'$ ), that the line which divides this angle (called the acute bisectrix) forms an angle of  $32^\circ 50'$  with the  $c$  axis, and



FIG. 164.—Model of prism and basal plane.




FIG. 165.—Malachite section parallel to (010).

that the angle between  $c$  and  $a$ , the  $\beta$  angle, is  $61^\circ 50'$ . A cross-section of a malachite crystal (Fig. 165) parallel to the clinopinacoid shows the relationship of these various directions.

#### SUMMARY

*Malachite*.— $\text{CuCO}_3 \cdot \text{Cu(OH)}_2$ ;  $\text{CuO}=71.9$  per cent,  $\text{CO}_2=19.9$  per cent,  $\text{H}_2\text{O}=8.2$  per cent. Monoclinic;  $a:b:c=0.881:1:0.401$ .  $\beta=61^\circ 50'$ . (110), (001); twinned parallel (100); cleavage (010), (001) perfect; brittle; fracture uneven.

Hardness=3.5; gravity=4. Green; adamantine; silky; dull; translucent.  $\beta=1.88$ ; double refraction negative;  $\gamma-a=0.2$ .

Easily fusible; soluble in hydrochloric acid.

Urals, Cordilleras.

## Azurite

Azurite is another basic carbonate of copper conspicuous because of the beauty of its color, which is a deep azure blue. In composition it differs from malachite in having less copper oxide. Azurite has 69.2 per cent, malachite 71.9 per cent. Azurite by increase in water content changes to malachite, increasing about one-third in bulk, thus affording an illustration of a chemical action which would tend to rend inclosing minerals or rock. Azurite crystals partially changed into malachite are of common occurrence. Both of these minerals result from decomposition of copper sulphides, are useful ores of copper, and are found in the same localities. Azurite occurs in good monoclinic crystals in which the prism (110) and base (011) predominate, modified by pyramids (111) and domes (013) (Fig. 166). Chessy near Lyons, France, and Bisbee, Arizona (Nos. 3409, 3410), are noted for their fine crystals.




FIG. 166.—Azurite crystal, showing also position of optic axes and axial plane.

## SUMMARY

*Azurite*.— $2\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$ ;  $\text{CuO}$ =69.2 per cent,  $\text{CO}_2$ =25.6 per cent,  $\text{H}_2\text{O}$ =5.2 per cent. Monoclinic;  $a:b:c=0.85:1:1.76$ ;  $\beta=87^\circ 36'$ . (110), (001), (111), (103); cleavage parallel (011) fair; brittle; fracture conchoidal.

Hardness=4; gravity=3.8. Deep to light blue; vitreous; translucent in thin pieces. Double refraction positive;  $\gamma-a=0.2$ . Axial plane perpendicular to (010); acute bisectrix makes an angle of  $75^\circ$  with  $a$  and lies in the supplementary angle to  $\beta$ .

Easily fusible (2); soluble in hydrochloric acid.

Urals, France, Arizona.



## CLASS VII. SILICATES

The carbonates are a very important group of oxygen salts, as we have just seen, but the silicates are even more important, especially if we consider their number of species and their quantity. Their quantity is so great that they constitute nine-tenths of the mass of the earth's crust. There are several hundred species, from which we select about forty only. These forty are most worthy of our attention because of perfection of form, beauty of color, durability, utility, or abundance. Some of them are so abundant as to be the chief minerals in great mountain chains. Some are highly prized as gems, others as minerals useful for chemical and various commercial purposes. But it is as rock-forming minerals *par excellence* that they most strongly bespeak the attention. The most abundant representatives of the section are the minerals first to be considered, namely, the feldspars.

### FELDSPAR GROUP

Though the word feldspar was used by the Germans as early as 1750, the minerals to which we now give that name have been distinguished for not much more than one hundred years and have been thoroughly studied only within the last seventy-five years. Today no minerals are better described and understood. Their investigation has contributed largely to the science of mineralogy. They deserve the attention which they have received, both because of their scientific and because of their commercial importance.

They afford excellent illustration of the relation of crystal structure to physical properties, such as the transmission of light and heat, and form an important feature in the classification of igneous rocks. Commercially they are of importance because of their use in manufacture and agriculture. They are the source of clay—a mineral substance valuable in soils; useful for paving and building brick, porcelain and china; and essential to the artist and artificer for modeling, to the manufacturer of woollens as fuller's earth, and to the chemist and smelter as fire clay.

They are found in nearly all parts of the country, but are most characteristic of mountain regions and of areas covered by glacial drift. For this reason the southern Mississippi Valley is about the only portion of the country in which feldspar may rarely be found. All of the feldspars in Illinois are found in the drift which covers the larger part of the state. For the best specimens one turns to igneous rocks in mountain regions.

All of the feldspars are aluminium silicates of potassium, sodium, or calcium, and rarely barium. Their prevailing color is white or light shades of red; they are about 2.5 in specific gravity, 6 in hardness, and split readily in two directions. They are divided into two sections because some of them crystallize apparently in the monoclinic system and others in the triclinic system.

The most important feldspar with the monoclinic habit is orthoclase.

## THE ORTHOCLASES

### Orthoclase

This is a potassium feldspar ( $\text{KAlSi}_3\text{O}_8$ ) in which a little of the potassium may be replaced by sodium. The color of orthoclase varies from colorless, glassy adularia and sanidine (No. 3477) to white, gray, yellowish, or reddish individuals (Nos. 3413 to 3415), or masses more or less opaque owing to partial conversion into kaolin.

Orthoclase is insoluble in acids without previous fusion. Pure orthoclase fuses at 5 in the scale of fusibility (about  $1150^\circ \text{C.}$ ), but with the increase of sodium, which is probably due to an intermixture of a sodium feldspar like albite, the melting-point is lowered.

Orthoclase is the chief constituent of the granitic rocks which occur in such great masses in the Rocky Mountains, the Alps, and other mountain regions. When mingled with the quartz and mica which constitute a large part of granite, the crystal outlines of orthoclase may be undeveloped, but may still be readily recognized by the cleavage planes which are parallel to the base (001) and clinopinacoid (010) and are at right angles to each other. The quartz in granite is without cleavage and the mica cleaves in one direction only and is elastic.

When the orthoclase is in druses (cavities lined with crystals), well-developed forms like those in Figure 167 show its pronounced




FIG. 167.—Model of an orthoclase crystal.




FIG. 168.—Orthoclase




FIG. 169.—Adularia orthoclase




FIG. 170.—Model of Carlsbad twin, interpenetrating parallel (010); twinning axis  $c$ .




FIG. 171.—Baveno twin, composition face (021).




FIG. 172.—Manebach twin, composition face (001).

tendency to assume clearly marked outlines. Such crystals are found abundantly in Colorado and are composed of prisms (110), brachypinacoids (010), domes (201), and basal planes (001). The dome (201) is nearly at right angles to the base. Quite similar to them in shape but usually thinner, in boardlike forms parallel to (010), is sandidine (from *savils*, "a table"), which is found as limpid glassy crystals in the feldspar blocks thrown out of Vesuvius and as large, dull-gray crystals in trachyte at the Drachenfels, Germany.

Forms less columnar but elongated in the direction of the edge between the base and clinopinacoid and having in addition pyramid planes (111) and clinodomes (021) (Fig. 168) are customary in the pink feldspar of the Baveno granite quarries.

Glassy crystals, called adularia from Adula, the old name of St. Gothard, Switzerland, where they are found in abundance, have the form shown in Figure 169. In all these crystals the cleavage parallel to (001) is most perfect, that parallel to (010) is but little inferior, while that parallel to (110), though barely evident, is important in the orientation of the crystal. When prismatic cleavage lines




FIG. 173.—Orthoclase section parallel to (010); axial plane, angle of extinction.

are placed vertically and the basal plane turned until it slants downward toward the observer, the crystal is in conventional position. The clinopinacoid is almost always vertically striated. Cleavage cracks on (001) often produce a pearly luster. The pale-blue opalescence of the Ceylon "moonstone" is due to cleavage cracks, to inclusion of feldspathic material, or to decomposition.

Twinned crystals are as abundant as are simple forms, and the twinning follows three laws which have been named after three localities where multitudes of good specimens are found and were

first studied. The localities are Carlsbad, Bohemia; Baveno, Piedmont, northwestern Italy; and Manebach, Saxe-Gotha, Germany.

Carlsbad law: Two individuals interpenetrate parallel to  $(010)$ , one of them being turned around on the  $c$  axis until the back side is toward the front (Fig. 170). The twinning axis is  $c$  and the composition face is  $(010)$ . If the crystal is terminated by  $(\bar{1}01)$  and  $(001)$  instead of  $(\bar{1}01)$  only, the two planes will be nearly in the same plane but may be distinguished, since the base is a plane of cleavage while the dome is not. Also while the base is smooth and bright, the dome may be dull.




FIG. 174.—Orthoclase (adularia);




FIG. 175.—Orthoclase; positive and negative direction of extinction on  $(001)$  and  $(010)$ .

Baveno law: Two individuals are united parallel to the clinodome  $(021)$  and one is revolved  $180^\circ$  so that the original back planes are turned toward the front. Twinning and composition plane is  $(021)$ . Since the angle formed by  $(021)$  and  $(001)$  is of nearly  $45^\circ$  and that between  $(021)$  and  $(010)$  is the same, Baveno twins may appear to be rectangular prisms (Fig. 171). But cleavage planes will show that two basal planes  $(001)$  come together on one edge and two clinopinacoids  $(010)$  on the other. At the ends will be observed a diagonal line of union. At one end are small salient angles formed between  $(\bar{2}01)$  and  $(201)$  ( $13^\circ 42'$ ) and between  $(110)$  and  $(\bar{1}\bar{1}0)$  ( $10^\circ 34'$ ). At



PLATE XXVI



*a*, Microcline, "Amazon Stone," Pike's Peak, Colorado



*b* Microcline, Pike's Peak, Colorado

the other end are re-entering angles. Baveno twins are often repeated to produce quartets.

Manebach law: Two individuals are twinned and composed on (001). The two clinopinacoids then fall in the same plane, but the end discloses the twinning (Fig. 172).

#### SUMMARY

*Orthoclase*.— $\text{KAlSi}_3\text{O}_8$ ;  $\text{K}_2\text{O}=16.9$  per cent,  $\text{Al}_2\text{O}_3=18.5$  per cent,  $\text{SiO}_2=64.6$  per cent. Monoclinic;  $a:b:c=0.658:1:0.555$ .  $\beta=116^\circ 3'$ . (110), (001), (010), (101), (201), (021), (111). Carlsbad law, twinning axis  $c$ : composition plane (010). Baveno law, twinning plane and composition plane (021). Manebach law, twinning plane and composition plane (001). Cleavage parallel (001), (010) perfect, (110) imperfect. Brittle; fracture conchoidal.

Hardness=6; gravity=2.5. Colorless to red; vitreous; transparent;  $\beta=1.524$ ; double refraction negative, weak;  $\gamma-a=0.006$ . Axial plane perpendicular to (010). Acute bisectrix  $5^\circ$ , above  $a$  axis on (100): extinction on (010)  $-5^\circ$ , on (001)  $0^\circ$ .  $2E$  varies but usually is in the neighborhood of  $120^\circ$ .

Insoluble; fusible (5).

In granite, gneiss, trachyte. More common in metamorphic rocks than are plagioclases. Mountain regions, Colorado, Switzerland, Italy; confined to the drift in Illinois.

#### Microcline

Closely related to orthoclase and connecting it crystallographically with plagioclase is microcline. In orthoclase ( $\delta\rho\theta\sigma\varsigma$ , "straight";  $\kappa\lambda\acute{\alpha}\omega$ , "to cleave") the basal plane and cleavage parallel to it form a right angle to the clinopinacoid plane and the cleavage parallel to it. In microcline ( $\mu\iota\kappa\rho\acute{o}\varsigma$ , "small";  $\kappa\lambda\iota\nu\epsilon\iota\nu$ , "to incline") these planes deviate but about  $15'$  to  $35'$  from a right angle; that is, the angle which (001) makes with (010) is generally about  $89^\circ 30'$ . In plagioclase ( $\pi\lambda\acute{\alpha}\gamma\iota\omega\varsigma$ , "oblique";  $\kappa\lambda\acute{\alpha}\omega$ , "to cleave") the angle between the basal plane (001) and brachypinacoid (010) is about  $86^\circ$ .

The chemical composition, hardness, specific gravity, and general character of microcline (No. 3418) are the same as those of orthoclase, and were it not for the inclination of the  $b$  axis to the  $c$  the two species would be classed as one. But because of their inclination microcline is a triclinic feldspar. Its crystals are generally composed of two sets of twins, one parallel to the brachypinacoid (010)



and the other at right angles to it, namely, parallel to the edge formed by (001), the base, and (101), the macrodome. This produces a curious cross-hatching or grating structure best seen under the microscope when the nicols are crossed as two series of lamellae in the basal plane. It is called the "microcline structure." It would be possible for the twins which cause the cross-hatching to be so fine as to be invisible. Then the inclination of (001) to (010) might disappear, the cross-hatching become indistinguishable, and microcline then resemble orthoclase completely. For this reason some authors think that orthoclase is simply an extreme form of microcline and that all of the feldspars are triclinic.

Green varieties of microcline are called Amazon stone. Pike's Peak, Colorado (Nos. 3416, 3245, 3420, and 1251), and the Ilmen Mountains in Russia have long been reputed for fine specimens of Amazon stone which they have furnished. Museums are quite generally supplied with specimens from these localities.

#### SUMMARY

*Microcline*.—Like orthoclase, except that the *b* axis is inclined about half a degree, the basal cleavage often shows fine striations and between crossed nicols cross-hatching, and the angle of extinction on base is  $+15^{\circ} 30'$ , while in orthoclase it is  $0^{\circ}$ .

*Anorthoclase*.— $(\text{Na}, \text{K})\text{AlSi}_3\text{O}_8$ . Triclinic; resembles microcline. Cross-hatching often so fine as to be scarcely visible; (201) and (110) often elongated.

In alkaline lavas (Pantellaria) and porphyries (Christiania).

*Hyalophane*.— $\text{K}_2\text{BaAl}_4\text{Si}_8\text{O}_{24}$ . A double salt composed of two molecules of orthoclase ( $\text{KAlSi}_3\text{O}_8$ ) and one of barium alumino-silicate ( $\text{BaAl}_2\text{Si}_4\text{O}_{16}$ ). It occurs in monoclinic glassy crystals which closely resemble adularia (Fig. 174).

In dolomite in Tyrol, etc.

---

Of the more than threescore minerals thus far considered, but one before microcline has been found to crystallize in the triclinic system, and that one, cryolite, is so poor a representative of the system that no description of triclinic symmetry was given in that connection. But microcline, and more especially the plagioclases now about to be described, are the most important minerals constructed with triclinic symmetry.

In the triclinic system are grouped such forms as have three axes of unequal lengths crossing each other at angles not right angles.

The angle between  $c$  and  $b$  is called  $\alpha$ , that between  $c$  and  $a$ ,  $\beta$ , and that between  $a$  and  $b$ ,  $\gamma$ , as indicated in Figure 176. Upon these axes planes are constructed which cut all three axes, pyramid planes (111); or which cut two axes and are parallel to the third, prisms (110) and domes (011), (100); or which cut one axis and are parallel to two, pinacoids (010), (100), (001). The longer of the two lateral axes is chosen as  $b$  and the dome and pinacoid parallel to it are the macrodome and macro-pinacoid. The brachydome and pinacoid are parallel to  $a$ . The figures produced by any one kind of plane are not closed forms. They have neither planes nor axes of symmetry but a center of symmetry only. The figures may readily be drawn by following the method employed for preceding systems. This system is the sixth and completes the crystallographic groups which embrace all crystallized minerals.




FIG. 176.—Triclinic axes of copper sulphate; no axes at right angles.

### THE PLAGIOCLASES

The feldspars which exhibit between the two chief cleavage planes angles markedly different from a right angle are grouped together as plagioclases. They compose two species widely separated chemically, one being the sodium feldspar, albite ( $\text{NaAlSi}_3\text{O}_8$ ), and the other the lime feldspar, anorthite ( $\text{CaAl}_2\text{Si}_2\text{O}_8$ ). Between these two is a series formed by insomorphous mixtures of albite and anorthite molecules in varying proportion. If Ab represents the albite molecule and An the anorthite molecule, the typical composition of the chief varieties may be expressed as follows:

Albite	Ab
Oligoclase	Ab <sub>3</sub> An
Andesine	Ab An
Labradorite	Ab An <sub>3</sub>
Bytownite	Ab An <sub>4</sub>
Anorthite	An

In physical as well as chemical properties these six minerals form a continuous series and afford a fine illustration of the variations of optical characters with molecular constitution. They may readily be distinguished from each other by microscopic or by chemical tests.

Plagioclase strongly resembles orthoclase in crystallographic and physical properties, but may usually be recognized by the fine striations on the basal plane. These striations are due to repeated twinning of thin leaves parallel to the brachypinacoid (010).

Some plagioclases are more common than orthoclase in igneous rocks, both as constituent of the ground mass and as inclosed crystals (phenocrysts). The six different kinds are shown in their order, beginning with albite.

### Albite

Albite (*albus*, "white") occurs in white masses or as crystals in cavities in igneous rocks and crystalline schists in the Appalachian (No. 3124) and Cordilleran regions and in all the mountain ranges of the world.




FIG. 177.—Albite




FIG. 178.—Albite twinned on (010); albite law.

The simple crystal (Fig. 177) resembles orthoclase in outline, as do also some of the twins which follow the Carlsbad, Baveno, and Manebach laws. But the most common and typical crystals are twinned according to the albite and the pericline laws.

According to the albite law the twinning plane and the composition face are (010) and the basal planes form re-entrant angles with each other (Fig. 178). The different crystals are usually thin as

paper and repeated so as to produce polysynthetic twins which give rise to fine striations and pearly luster best seen on the basal plane.

According to the pericline law (*Περικλινής*, "sloping," so called because of the oblique appearance of the crystals which are often elongated parallel to the  $b$  axis) (Fig. 179), the  $b$  axis is the twinning axis and the twin crystals are united along a face called the rhombic section, which is parallel to  $b$  but slopes backward forming an angle of  $27^\circ$  with




FIG. 179.—Albite, pericline twin




FIG. 180.—Albite, (001), (010), (110), (101); axial plane.




FIG. 181.—Albite section parallel (100)




FIG. 182.—Albite section parallel (010)

the edge (001) and (010). The portions of (010) above and below the line of union form an angle with each other.

Albite resembles orthoclase in hardness and fusibility but is heavier and may readily be distinguished by use of a heavy liquid like methylene iodide (sp. gr. = 3.3).

It often incloses minerals prized as gems, such as the phenacite, topaz, tourmaline, and beryl of the Urals, Ilmen, and Rocky Mountains.

#### SUMMARY

*Albite.*— $\text{NaAlSi}_3\text{O}_8$ ;  $\text{Na}_2\text{O}$  = 11.8 per cent,  $\text{Al}_2\text{O}_3$  = 19.5 per cent,  $\text{SiO}_2$  = 68.7 per cent. Triclinic; symmetry holosymmetric;  $a:b:c = 0.6335:1:0.5577$ .  $\alpha = 94^\circ 3'$ ,  $\beta = 116^\circ 29'$ ,  $\gamma = 88^\circ 9'$ .  $(110)$ ,  $(010)$ ,  $(001)$ ,  $(20\bar{1})$ ,  $(11\bar{1})$ . Common twins; twinning plane and composition face  $(010)$  for albite law; axis  $b$  is the twinning axis and the rhombic section the composition plane for pericline law. Cleavage  $(001)$ , perfect;  $(010)$ ,  $(110)$ , imperfect; brittle; fracture uneven.

Hardness = 6; gravity = 2.63. Colorless; vitreous, transparent;  $\beta = 1.533$ . Double refraction positive, weak;  $\gamma - \alpha = 0.011$ . Acute bisectrix  $c$  in zone  $(010)$ :  $(001)$  inclined at  $16^\circ$  to the perpendicular of  $(001)$ . Obtuse bisectrix  $a$  inclined at  $70^\circ$  to the perpendicular of  $(001)$ . Red light less dispersed than violet ( $\rho < \nu$ ).

Insoluble; fusible at 4.

Rocky Mountains, Ural Mountains, etc.

#### Anorthite

The plagioclase farthest removed from albite in composition and physical character is the comparatively rare mineral anorthite ( $\text{CaAl}_2\text{Si}_2\text{O}_8$ ) found in basalt in Japan, in limestone and augitic blocks ejected from Vesuvius, etc. The crystals are limpid glassy forms exhibiting a larger number




FIG. 183.—Rhombic section of anorthite.




FIG. 184.—Anorthite, showing position of axial plane and bisectrix.

of faces than albite and having less marked striations on the brachypinacoid. The optical properties are very different from those of albite, as may be seen in the table (p. 144). Carlsbad, Manebach, pericline, and albite laws of twinning are exhibited. When twinned according to the pericline law the rhombic section slopes down forward from the basal plane (Fig. 183). Such a direction of slope is called negative; the backward slope such as that shown by albite is called positive.

Anorthite is heavier than albite and more difficultly fusible (5), but more easily soluble.

#### SUMMARY

*Anorthite*.— $\text{CaAl}_2\text{Si}_2\text{O}_8$ ;  $\text{CaO}$ =20.12 per cent,  $\text{Al}_2\text{O}_3$ =36.72 per cent,  $\text{SiO}_2$ =43.16 per cent. Triclinic; symmetry holosymmetric;  $a:b:c=0.6347:1:0.5501$ .  $\alpha=93^\circ 13'$ ,  $\beta=115^\circ 55'$ ,  $\gamma=91^\circ 12'$ . (110), (010), (001), (100), (10 $\bar{1}$ ), (20 $\bar{1}$ ), (021), (023), (111), (11 $\bar{1}$ ), (1 $\bar{1}$ 1), (207), (130). Twinned according to Carlsbad, Baveno, Manebach, albite, and pericline laws. Cleavage (001), perfect; (010), fair. Brittle; fracture conchoidal.

Hardness=6.5; gravity=2.75. Colorless; vitreous; transparent;  $\beta=1.58$ . Double refraction weak, negative;  $\gamma-\alpha=0.013$ . Obtuse bisectrix  $\gamma$  perpendicular to (02 $\bar{1}$ ). Acute bisectrix ( $\alpha$ ) inclined inside the crystal at  $53^\circ 14'$  to the normal of (001), at  $58^\circ$  to normal of (010), at  $16^\circ 52'$  to normal of (11 $\bar{1}$ ).

Decomposed by hydrochloric acid; fusible at 5.

Vesuvius, Japan.

#### INTERMEDIATE PLAGIOCLASES

Oligoclase, andesine, labradorite, and bytownite are usually found as rock constituents in massive or microscopic forms lacking well-defined faces. Chemical or optical tests are necessary to distinguish these plagioclases from each other.

Oligoclase (soda lime feldspar) is found as grayish-white, translucent, somewhat greasy-looking masses with orthoclase in granite (No. 3422). The striations on the basal planes aid in distinguishing it from the orthoclase. Reddish cleavable masses with a golden shimmer due to spangles of hematite or goethite are called "adventurine feldspar" (Nos. 3423, 3424) or "sunstone."

Labradorite (lime soda feldspar), found abundantly in Labrador (Nos. 3425, 3427, 3428), the Adirondacks, and Wichita Mountains,

occurs in dark cleavable masses, commonly iridescent, and showing beautiful red, yellow, and green colors. The colors are due partly to interference effects caused by twinning lamellae and partly to inclusions of hematite, goethite, or diallage. Labradorite is notably absent in rocks containing orthoclase and quartz, but is characteristic of basic rocks like gabbro and dolerite.

Often associated with the labradorite is another iridescent mineral, hypersthene. The rock formed chiefly of these two minerals is called "labrador spar" and is used extensively in decorative work. The finest church in Moscow (St. Savior), with capacity for seven thousand worshipers at one time, is wainscoted with beautiful chatoyant labradorite.

#### CHEMICAL CHARACTERISTICS AND SPECIFIC GRAVITY OF THE FELDSPARS

	Composition	SiO <sub>2</sub>	Al <sub>2</sub> O <sub>3</sub>	K <sub>2</sub> O	Na <sub>2</sub> O	CaO	Specific Gravity
Orthoclase.....	KAlSi <sub>3</sub> O <sub>8</sub>	64.6	18.4	16.9	.....	.....	2.56
Microcline.....	.....	64.6	.....	.....	.....	.....	2.56
Anorthoclase.....	NaKAlSi <sub>3</sub> O <sub>8</sub>	.....	.....	.....	.....	.....	.....
Albite.....	NaAlSi <sub>3</sub> O <sub>8</sub>	68.7	19.5	.....	11.8	0	2.62
Oligoclase.....	Ab <sub>3</sub> An	62.0	24.0	.....	8.7	5.3	2.65
Andesine.....	AbAn	55.6	28.3	.....	5.7	10.4	2.69
Labradorite.....	AbAn <sub>2</sub>	49.3	32.6	.....	2.8	15.3	2.72
Bytownite.....	AbAn <sub>3</sub>	46.6	34.4	.....	1.6	17.4	2.74
Anorthite.....	CaAl <sub>2</sub> Si <sub>2</sub> O <sub>8</sub>	43.2	36.7	.....	0.	20.1	2.75

#### CRYSTALLOGRAPHIC CHARACTERISTICS OF THE FELDSPARS

	Axes	Angle between Axes			Extinction on 001	Extinction on 010	Mean Refraction $\beta$	Rhombic Section Angle
	a: b: c	$\alpha$	$\beta$	$\gamma$				
Orthoclase....	0.658:1:0.555	90°	116°3'	90°	0°0'	5°	1.523	.....
Microcline...	0.658:1:0.555	89°30'	116°3'	90°	15°30'	5°	1.526	.....
Anorthoclase..	.....	.....	.....	.....	.....	.....	.....	.....
Albite.....	0.633:1:0.557	94°3'	116°29'	88°9'	4°30'	19°	1.534	27°
Oligoclase.....	0.632:1:0.552	93°4'	116°23'	90°5'	1°4'	4°36'	1.542	3°
Andesine.....	0.635:1:0.552	93°23'	116°29'	89°59'	5°10'	16°	1.558	1°
Labradorite...	0.637:1:0.554	93°31'	116°3'	89°54'	17°40'	29°28'	1.570	9°
Bytownite.....	.....	.....	.....	.....	27°33'	33°29'	.....	10°
Anorthite....	0.634:1:0.550	93°15'	115°55'	91°12'	37°	36°	1.582	16°

### Leucite

Leucite (λευκός, "white") crystallizes in white, round crystals (Fig. 7) from an inch in diameter to forms microscopic in size in igneous rocks, such as those found in the Leucite Hills, southwestern Wyoming, and at Mount Vesuvius (No. 3244).

When leucite assumes its definite form at temperatures above  $500^{\circ}$  the crystals are trapezohedrons, but at ordinary temperatures, while the external form remains the same, the internal condition is that of the orthorhombic system. The microscope reveals multitudes of fine layers which cross each other at right angles, and occasionally groups diagonal to these (Fig. 185). These layers have the optical properties characteristic of the orthorhombic system. Hence leucite is said to be pseudo-regular. Externally it is always regular, internally under ordinary conditions it is orthorhombic. Crystals microscopic in size are often free from the orthorhombic layers.




FIG. 185.—Microscopic section of leucite between crossed nicols.

Furthermore, inclusions of other minerals, such as augite, olivene, and apatite, are arranged parallel to trapezohedral faces—an additional indication that the mineral swings between the orthorhombic and the regular system. Its instability is further shown by the fact that in nature it is usually white and opaque, owing to its change into analcite, mica, or kaolin.

### SUMMARY

*Leucite*.— $\text{KAl}(\text{SiO}_3)_2$ ;  $\text{K}_2\text{O} = 21.58$  per cent,  $\text{Al}_2\text{O}_3 = 23.40$  per cent,  $\text{SiO}_2 = 55.02$  per cent. Orthorhombic (211). At ordinary temperature all but microscopic crystals contain fine orthorhombic lamellae which are weakly doubly refracting. Cleavage parallel (110) imperfect; brittle; fracture conchoidal.

Hardness = 5.5; gravity = 2.5. Colorless; vitreous; transparent;  $\beta = 1.508$ .

Infusible; soluble in hydrochloric acid.

Vesuvius, Eifel, Wyoming.

### PYROXENE GROUP

Hardly less abundant than the white feldspars as rock constituents are the colored pyroxenes. The name pyroxene ( $\pi\rho\upsilon\rho$ , ξένος,



"a stranger to fire") was given by Haiiy to certain green crystals found in Italian lavas, with the thought that they were only accidentally present, having been caught up while the lava was passing surrounding rock. Now it is known that they are essential constituents of many igneous rocks and are as much misnamed as is a white blackbird! According to their crystallography, they are classified in three sections, the orthorhombic, the monoclinic, and the triclinic. Most prominent in the first section are

### ORTHORHOMBIC PYROXENES

#### Enstatite and Hypersthene

Enstatite is a white, green, or brownish mineral occurring rarely in columnar, orthorhombic crystals, and commonly in fibrous, silky, opaque masses. Measurable crystals were found by von Lang in 1871 in a meteorite which fell in Bohemia, and three years later certain large crystals contained in Norwegian schists were found to be enstatite by Brögger. Chemically enstatite is a magnesium silicate ( $\text{MgSiO}_3$ ). The presence of iron often gives a metallic luster (bronzite). When iron replaces a large part of the magnesium, a somewhat darker, heavier, and more soluble mineral, hypersthene, results.

When a thin section of either of these minerals cut parallel to the prism or brachypinacoid plane is viewed under a microscope between crossed nicols, it becomes dark if the crystal axes are parallel to the cross-hairs of the microscope, as is characteristic of an orthorhombic mineral. The extinction is said to be parallel. Further, these pyroxenes change from reddish-brown to green according to the direction in which a thin section is viewed under the microscope. That is, they are strongly pleochroic, and this property aids in distinguishing them from the pyroxenes next to be studied.

Both enstatite and hypersthene are found in granular eruptive rocks and schists such as are common in most mountain regions.

#### SUMMARY

*Enstatite*.— $\text{MgSiO}_3$ ;  $\text{MgO}$ =40 per cent,  $\text{SiO}_2$ =60 per cent. Orthorhombic;  $a:b:c=0.97:1:0.57$ .  $(110):(1\bar{1}0)=88^\circ 20'$ . Cleavage parallel  $(110)$ ,  $(010)$ . Brittle; fracture even.

Hardness=5.5; gravity=3.2. White, green, brown; vitreous; translucent; mean refraction  $\beta=1.659$ . Double refraction weak, positive;  $\gamma-\alpha=0.009$ . Axial plane  $(010)$ ; acute bisectrix, normal to  $(001)$ ,  $2H=79^\circ$ ;  $\rho < \nu$ .

Infusible, named after ἐναντίας, "opponent," because so refractory. Insoluble; with cobalt turns pink. Decomposes into serpentine and talc. New York, Norway, Germany. Common in meteorites.

*Hyperssthene*.— $(\text{MgFe})\text{SiO}_3$ ; MgO from 26 to 11 per cent, FeO from 10 to 34 per cent. Agrees with enstatite except as follows:

Gravity=3.4. Dark green to black.  $\beta=1.702$ . Double refraction weak, negative;  $\gamma-\alpha=0.013$ . Acute bisectrix normal to (100). Dispersion  $\rho > \nu$ .

Fusible to magnetic mass. Partly soluble in hydrochloric acid.

Common with labradorite in granular eruptive rocks in Labrador, Greenland, Norway, New York.

### MONOCLINIC PYROXENES

The monoclinic section of pyroxenes is more important than either the orthorhombic, already considered, or the triclinic, which will be studied later. The chief monoclinic pyroxenes are

#### Diopside and Augite

These two minerals occur in short, stout, green to black crystals in igneous rocks, and are nearly as abundant as is the feldspar in these rocks (Nos. 3676, 3677, 3680).

Diopside (Nos. 3429 and 3430) is clear pale green in color. Sometimes a crystal is darker at one




FIG. 186a.—Diopside



FIG. 186b.—Photograph of diopside from Cantley, Quebec, Canada: (111), (100), (101), (110), (100), (010); about 12 inches long.

end than at the other, and also differently terminated at the opposite ends. The usual shape of the crystal is illustrated in Figures 186–88. In composition diopside is a silicate of calcium and magnesium,  $\text{CaMg}(\text{SiO}_3)_2$ .




FIG. 187.—Diopside




FIG. 188.—Diopside, showing optic axes, acute bisectrix, axes of elasticity.




FIG. 189.—Augite




FIG. 190.—Augite twin parallel to (100)

Augite is dark green or black (No. 3431). The crystals are usually terminated with pyramidal planes, while prism and pinacoid planes are both well developed. Twins parallel to the orthopinacoid are common (Figs. 189 and 190).

While iron is usually present in both diopside and augite, it is more abundant in the latter. Diopside lacks aluminum. Therefore the pyroxenes are often divided into non-aluminous (diopside) and aluminous (augite) varieties.

Light green or white diopside is abundant in crystalline limestones and dolomites. Green or black augite is common in granite or eruptive rocks (No. 3433). When black basalt decomposes, augite crystals sometimes fall out and may be easily collected in quantities.




FIG. 191.—Augite cross-section perpendicular to prism planes.

A thin section of augite cut perpendicular to the  $c$  axis when viewed under the microscope




FIG. 192.—Enstatite, showing parallel extinction.




FIG. 193.—Diopside, showing oblique extinction angle of  $38^\circ$ .

shows fine cleavage lines parallel to the prism planes. These lines form an angle of nearly  $90^\circ$  with each other and the outline of the figure is eight-sided (Fig. 191). The eight-sided outline, the cleavage angle, and lack of pleochroism aid in distinguishing augite from hornblende. Hornblende presents a six-sided figure with cleavage line forming an angle of  $124^\circ$ , and furthermore is strongly pleochroic.

Fragments or thin sections of the members of the pyroxene group may be recognized by their extinction angle. A section of enstatite

cut parallel to the brachypinacoid and viewed in parallel light between crossed nicols becomes dark when the  $c$  axis is parallel to one of the cross-hairs of the microscope (Fig. 192). Diopside when so examined must be turned clockwise about  $38^\circ$  before it becomes dark (Fig. 193). Augite requires even a greater angle of revolution, sometimes being as large as  $54^\circ$ . In each case the angle of extinction increases with the amount of iron present.

#### SUMMARY

*Diopside*.— $\text{CaMg}(\text{SiO}_3)_2$ ;  $\text{CaO}=25.9$  per cent,  $\text{MgO}=18.5$  per cent,  $\text{SiO}_2=55.6$  per cent. Monoclinic;  $a:b:c=1.0921:1:0.5893$ .  $\beta=105^\circ 50'$ . (110), (100), (010), (001), (10 $\bar{1}$ ), ( $\bar{1}$ 11), ( $\bar{2}$ 21). Twinned on (100). Cleavage perfect parallel (110), imperfect parallel (100), (010). Brittle; fracture conchoidal.

Hardness=5.5; gravity=3.3. Light green; vitreous; transparent. Mean refraction ( $\beta$ )=1.681, maximum ( $\gamma$ )=1.703. Double refraction positive, strong;  $\gamma-a=0.030$ . Axial plane (010). Acute bisectrix inclined  $37^\circ 35'$  to  $c$  and the obtuse bisectrix inclined below  $a$  in front  $21^\circ 45'$  (Fig. 193). Axial angle ( $2E$ )= $68^\circ$ .

Fusible; insoluble.

In crystalline limestones and dolomites, both east and west, and in the drift.

*Augite*.— $\left\{ \begin{array}{l} \text{CaMg}(\text{SiO}_3)_2 \\ \text{MgAl}_2\text{SiO}_6 \end{array} \right\} \text{Fe}_2\text{O}_3$ , is also always present.

Augite has nearly the same character as diopside, but contains  $\text{Al}_2\text{O}_3$  and  $\text{Fe}_2\text{O}_3$  in addition to calcium and magnesium silicates, and is darker in color.

More readily fusible than diopside.

In granitic and eruptive rocks the world over.

#### Jadeite

Jadeite is a compact, tough, alkaline pyroxene having the composition  $\text{NaAl}(\text{SiO}_3)_2$ . It is 7 in hardness and 3.3 in specific gravity, is translucent, and varies in color from blue and green to white. When carved and polished it has a soft waxy luster which is very pleasing, and for that reason it has been used for many hundreds of years as material for carving into ornaments, vases, etc. Being tough and hard, it is very enduring.

## SUMMARY

*Jadeite*.— $\text{NaAl}(\text{SiO}_3)_2$ ;  $\text{Na}_2\text{O}=15.4$  per cent,  $\text{Al}_2\text{O}_3=25.2$  per cent,  $\text{SiO}_2=59.4$  per cent. Monoclinic; massive, sometimes granular, slightly fibrous; fracture splintery; very tough.

Hardness=7; gravity=3.3. Greenish, bluish, white; waxy, dull, translucent;  $2V=72^\circ$ .

Fuses readily; not attacked by acids after fusion; different from saussurite.

Burma, Thibet, Mexico.

## TRICLINIC PYROXENE

The triclinic pyroxene rhodonite (*ρδδον*, “a rose”) is a beautiful red mineral which, because of its hardness (6) and fine color, is used for ornaments such as brooches, cuff buttons, watch charms, inkwells, paper weights, vases, mantelpieces, and table tops. In the Urals masses of such size have been found as to be available for tombstones. In the late Czar’s lapidary shops at Petrograd some years ago, the author saw an oblong block of rhodonite  $7 \times 4 \times 3$  feet in size, being carved for a sarcophagus for royalty, and which was valued at six hundred thousand dollars.




FIG. 194.—Rhodonite

When crystals of rhodonite occur, as is often the case in Norway, England, and New Jersey (No. 3438), they are tabular (Fig. 194) in form or stout like augite.

Rhodonite is a silicate of manganese ( $\text{MnSiO}_3$ ) but usually contains calcium, iron, and in New Jersey zinc.

## SUMMARY

*Rhodonite*.— $\text{MnSiO}_3$ ;  $\text{MnO}=54.1$  per cent,  $\text{SiO}_2=45.9$  per cent. Triclinic;  $a:b:c=1.073:1:0.621$ .  $\alpha=103^\circ 18'$ ,  $\beta=108^\circ 44'$ ,  $\gamma=87^\circ 39'$ . (110), (010), (001), (221). Cleavage (110), (110), perfect; (001), fair. Brittle; fracture uneven.

Hardness=6; gravity=3.6. Red; vitreous; translucent. Mean refraction ( $\beta$ )=1.73. Double refraction negative, weak;  $\gamma-a=0.010$ . Axial plane is inclined at  $63^\circ$  to (110) and  $38^\circ$  to (001). Acute bisectrix inclined  $51^\circ 47'$  to the normal of (110) and  $51^\circ 40'$  to the normal of (001). Axial angle ( $2H$ )= $79^\circ$ ;  $\rho < \nu$ .

Readily fusible (2.5); partly soluble in hydrochloric acid.

Urals, Norway, England, New Jersey.

### AMPHIBOLE GROUP

The minerals in this group resemble pyroxene in composition, color, and form.

Like the pyroxenes they have orthorhombic, monoclinic, and triclinic representatives. The term amphibole ( $\alpha\mu\phi\iota\beta\omicron\lambda\eta$ , "doubtful") was given by Haüy to replace the name "schorl" used by miners for both hornblendes and tourmaline—two minerals which, though resembling each other, are chemically and crystallographically different. The readiest means of distinguishing amphiboles from pyroxenes are the crystal form and cleavage, as may be seen from the descriptions following.

#### ORTHORHOMBIC AMPHIBOLE

Anthophyllite, corresponding to the orthorhombic pyroxene, hypersthene, occurs in brown fibrous or flaky masses in mica schist in Norway, Pennsylvania (No. 3440), North Carolina, etc. Crystals are extremely rare, but under the microscope it may be observed that cleavage parallel to (010) is inferior to that of hypersthene. Anthophyllite is less pleochroitic than hypersthene but has greater double refraction;  $\gamma-a=0.024$ .

The name anthophyllite is derived from the Latin word for clove (*anthophyllum*) because of the clove-brown color of the mineral.




FIG. 195.—Anthophyllite, axial plane and optic axes.

#### SUMMARY

*Anthophyllite*.— $(MgFe)SiO_3$ ;  $MgO \pm 27.8$  per cent,  $FeO = 16.6$  per cent,  $SiO_2 = 55.6$  per cent.

Orthorhombic; crystals rare; fine fibers or flakes common.

Hardness=6; gravity=3.1. Brown to green, at times metalloid; translucent. Mean refraction ( $\beta$ )=1.642. Double refraction positive;

$\gamma - \alpha = 0.024$ . Axial plane parallel (010). Acute bisectrix normal to (001).

Difficultly fusible; insoluble.

In gneisses and schists in Norway, Pennsylvania, North Carolina, etc.

### MONOCLINIC AMPHIBOLES

As the monoclinic pyroxenes are the most abundant and important of the pyroxenes, so the monoclinic amphiboles far surpass in abundance the orthorhombic and triclinic forms. The white tremolite, green actinolite, and black hornblende constitute the chief representatives of the monoclinic forms.

#### Tremolite

Tremolite corresponds to diopside but contains more magnesium and is usually fibrous or columnar and without terminating planes. It is white or pale green in color. As the amount of iron (FeO) increases, it becomes dark and is called actinolite (Nos. 3474, 3475), from the fact that it occurs in blades or rays (*ἀκτίνες*, "ray").

Tremolite is found in granular dolomite and actinolite in schists in the Alps and Appalachians (Nos. 582, 3435, 3444, and 3446) and other mountainous regions.

Actinolite is strongly pleochroic. Light passing through parallel to *a* is greenish yellow, parallel to *b* yellowish green, and parallel to *c* green.

#### SUMMARY

*Tremolite*.— $\text{CaMg}_3(\text{SiO}_3)_4$ ;  $\text{CaO} = 13.45$  per cent,  $\text{MgO} = 28.83$  per cent,  $\text{SiO}_2 = 57.72$  per cent. Monoclinic;  $a:b:c = 0.551:1:0.294$ .  $\beta = 106^\circ 2'$ . (110), (100), (010). Twinning plane 100. Cleavage (110) perfect; (100), (010) imperfect. Brittle; fracture uneven.

Hardness = 5.5; gravity = 3.1. Pale green; vitreous; translucent. Mean refraction ( $\beta$ ) = 1.62; maximum ( $\gamma$ ) = 1.63. Double refraction negative, strong;  $\gamma - \alpha = 0.028$ . Axial plane (010); acute bisectrix almost parallel to (001). Axial angle ( $2N$ ) =  $87^\circ 22'$ ;  $\rho < \nu$ . Extinction angle (between  $\gamma$  and *c*) =  $15^\circ$ .

Fusible; insoluble.

Crystalline dolomites and schists in many mountain regions.

#### Hornblende

Hornblende is the common black amphibole corresponding to augite, the common black pyroxene. It may be distinguished from



augite since the crystals and also their transverse sections are usually six-sided rather than eight-sided (Figs. 196-99) and the cleavage lines parallel to prism planes make an angle of  $56^\circ$ . In hardness, weight, etc., the two minerals are similar.

The chemical composition of hornblende is not as well understood as that of augite, but it may probably be best represented as a




FIG. 196.—Hornblende




FIG. 197.—Hornblende




FIG. 198.—Hornblende section perpendicular to prism.




FIG. 199.—Hornblende twinned on (100).

mixture of the actinolite molecule  $\text{Ca}(\text{Mg}, \text{Fe})_3(\text{SiO}_3)_4$  and a molecule containing aluminum in addition to the calcium and magnesium, thus  $\text{CaMg}_2\text{Al}_2(\text{SiO}_4)_3$ .

Hornblende is found in lava, as at Vesuvius, and in basalt, trachyte, gneiss, and schists in most mountain regions (Nos. 3441, 3442, 3443).

Excellent crystals weathering out of volcanic rocks show abundant twins parallel to the orthopinacoid (100) (Fig. 199).

## SUMMARY

*Hornblende*.— $\left\{ \begin{array}{l} \text{Ca}(\text{MgFe})_3(\text{SiO}_3)_4 \\ \text{CaMg}_2\text{Al}_2(\text{SiO}_3)_3 \end{array} \right\}$ . Crystallography the same as that of tremolite, but end planes are common; (001), (011), (101), (130). Cleavage parallel (110) with angle of  $56^\circ$  perfect. Brittle to tough.

Hardness=5.5; gravity=3.2. Dark green to black; vitreous; translucent to opaque. Color and optical properties vary with the amount of iron present. Mean refraction ( $\beta$ )=1.64 to 1.72. Double refraction positive, strong;  $\gamma-a=0.019$  to 0.072. Axial plane (010). Axial angle




FIG. 200.—Anthophyllite; parallel extinction.




FIG. 201.—Tremolite; extinction angle  $15^\circ$ .




FIG. 202.—Hornblende; extinction angle  $25^\circ$ .

( $2H$ )= $59^\circ$ . Extinction angle (between  $\gamma$  and  $c$ )= $15^\circ$ . Strong interference; strong pleochroism,  $c$  greenish blue,  $b$  emerald green,  $a$  greenish yellow.

Fuses readily and may become magnetic; insoluble.

Mountain ranges, in volcanic and metamorphic rocks.

## Asbestos

Asbestos is a kind of amphibole with fibers (Nos. 479, 480) which are of sufficient length and flexibility to permit of its being woven into cloth. In northern Italy (Lombardy) is found a white asbestos whose silky fibers are often three feet long. Usually the fibers are but an inch or less in length. When too short for weaving, they are used for insulation, fireproof packing, etc., since the mineral is a poor

conductor of heat and is infusible. The greater part of the asbestos used commercially is a fibrous variety of serpentine, a mineral to be described later.

"Mountain cork" and "mountain leather" are matted sheets or nodules of yellowish asbestos.

Georgia produces amphibole asbestos. Arizona, California, and Wyoming yield serpentine asbestos. But our chief supply is the chrysotile (fibrous serpentine) imported from Canada. It is woven into cloth (No. 1876) for theater curtains, gloves, firemen's suits, etc.; made into yarn, rope, paper, and boards; and used for covering steam pipes, lining safes, making paints, filters, etc.

In Griqualand, Africa, is found a fibrous, silky, blue chatoyant amphibole known as crocidolite (Nos. 3476 and 3478). Through oxidation and partial replacement by silica, this becomes converted into a hard, compact, golden-yellow mineral known as "cat's eye" or "tiger eye," and much used for ornaments.

### Nephrite

Nephrite (jade) is another variety of amphibole. It is compact, has a specific gravity of 3, and is harder than other amphiboles, its hardness being 6.5. It is one of the toughest of minerals, and has a splintery fracture. The white variety has the composition of tremolite, and the green variety that of actinolite. On account of its toughness, color, and translucency, like jadeite it has been much prized for centuries throughout the East as a material from which to carve ornaments and weapons. The ancients thought it a cure for kidney diseases (*νεφρός*, "kidney"; jade has the same meaning). It is found as boulders in China, India, New Zealand, Alaska, Mexico, and Germany.

### Beryl

When beryllium aluminium silicate ( $\text{Be}_3\text{Al}_2\text{Si}_6\text{O}_{18}$ ) occurs in greenish masses or in coarse, hexagonal, prismatic crystals, it is called beryl. Transparent, beautiful pale-green crystals are called aquamarine. The dark-green crystals are called emerald, and are among the most prized of gem minerals. Aquamarine means sea water; emerald and beryl are ancient names of unknown significance (Fig. 203a).

Beryl furnishes the best example of holosymmetric hexagonal symmetry found among minerals. Little, bright, pale-green crystals with many faces occur in Siberia and in the Urals (Fig. 203*b*).

Ekaterinburg, Russia, has long been known as a good source for aquamarines and emeralds. They are found in a coarse granite associated with topaz, black tourmaline, and smoky quartz in the neighboring mountains.

A tourmaline granite on the island of Elba and an albitic granite in the Mourne Mountains, Ireland, contain richly terminated beryls which vary in tint from colorless to green, blue, or red.



FIG. 203*a*.—Photograph of beryl from Brazil, (0001) and (10 $\bar{1}$ 0).




FIG. 203*b*.—Beryl

The best emeralds have been procured from Muso, Colombia, in a black, bituminous limestone, where they are accompanied by pyrite, calcite, black dolomite, and quartz. The prism form is strongly developed. The crystals have often been fractured and recemented by calcite.

Coarse beryl of gigantic dimensions has been found in New Hampshire and Massachusetts. Nos. 1667, 3232, and 500 represent the New Hampshire locality. No. 3483 comes from Connecticut. Colorado (No. 3482) and North Carolina have furnished quantities of beryl and a few emeralds.

Beryl crystals are often striated vertically and are usually without termination. They cleave imperfectly parallel to the base. Although without doubt hexagonal, they often show weak double refraction

because of strain. They resemble apatite but may be distinguished by their inferior cleavage and superior hardness.

Chromium doubtless furnishes the color to emerald and aquamarine. Flawless crystals are extremely rare.

#### SUMMARY

*Beryl*.— $\text{Be}_3\text{Al}_2\text{Si}_6\text{O}_{18}$ ;  $\text{BeO}$ =14.11 per cent,  $\text{Al}_2\text{O}_3$ =19.05 per cent,  $\text{SiO}_2$ =66.84 per cent. Hexagonal; holosymmetric;  $a:c=1:0.4989$ .  $(10\bar{1}0)$ ,  $(10\bar{1}1)$ ,  $(20\bar{2}1)$ ,  $(11\bar{2}1)$ ,  $(2131)$ ,  $(0001)$ . Cleavage parallel  $(0001)$  imperfect. Brittle; fracture conchoidal.

Hardness=7.5; gravity=2.7. Bluish green; vitreous; transparent.  $\omega=1.584$ ; double refraction negative, weak;  $\omega-\epsilon=0.006$ .

Fusible with difficulty (5.5). Insoluble.

Urals, Maine, New Hampshire, North Carolina, South Dakota, Colorado.

#### GARNET GROUP

Each of the six varieties of garnet which compose this group crystallizes in remarkably well-formed dodecahedrons, trapezo-




Fig. 204.—Garnet


hedrons, or combinations of these forms (Figs. 204–6). All have imperfect cleavage, conchoidal fracture, high refraction, and greasy luster. In color they are usually red, brown, or black, though some varieties are green or yellow.

They are silicate of calcium, magnesium, manganese, iron, aluminium, and chromium in isomorphous mixtures. Because they are mixtures their color, weight,

fusibility, and solubility are variable, as may be seen in the following table:



PLATE XXVII



*a*, Garnets; dodecahedron from Salida, Colorado, trapezohedron from North Carolina, and combination from Fort Wrangel, Alaska.



*b*, A dodecahedron nearly four inches in diameter from Salida, Colorado.

Grossularite,  $\text{Ca}_3\text{Al}_2(\text{SiO}_4)_3$ ; gravity, 3.5; hardness, 7; fusible (3); yellowish to red or brown.

Pyrope,  $\text{Mg}_3\text{Al}_2(\text{SiO}_4)_3$ ; gravity, 3.7; hardness, 7; fusible (3.5); deep red to black.

Almandite,  $\text{Fe}_3\text{Al}_2(\text{SiO}_4)_3$ ; gravity, 3.5-4.3; hardness, 7; fusible (3); deep red to black.

Spessartite,  $\text{Mn}_3\text{Al}_2(\text{SiO}_4)_3$ ; gravity, 3.8; hardness, 7; fusible (3); purple red to brown.

Uvarovite,  $\text{Ca}_3\text{Cr}_2(\text{SiO}_4)_3$ ; gravity, 3.4; hardness, 7; infusible (6); green.

Andradite,  $\text{Ca}_3\text{Fe}_2(\text{SiO}_4)_3$ ; gravity, 3.8-4.1; hardness, 7; fusible (3); yellow, red, black.

The ordinary varieties of garnet are abundant enough to be mined for use as an abrasive in New York, New Hampshire (No. 616), North Carolina, and Georgia. The only garnets found in Illinois are those occurring in gneiss and schists transported from the north by glaciers. Dentists use garnet disks for polishing, as do also shoe manufacturers, woodworkers, etc. At Morelos, Mexico,




FIG. 205.—Garnet




FIG. 206.—Garnet

a marble containing large pink garnets is polished for ornamental slabs.

Pyrope ( $\pi\rho\rho\omega\pi\acute{o}\varsigma$ , "fiery eyed") (Nos. 3489 and 3177) and almandite are much used in common jewelry, the former often being of a fine, transparent red color resembling ruby, and the latter of a purplish or hyacinth-red color. Almandite is often cut *en cabochon*, i.e., with flat base and rounded top.



Uvarovite (named after a Russian minister), found in small, brilliant green crystals in a serpentine rock in the Urals and in granular limestones in Canada, is cut into gem stones and sold as "olivine," though the latter mineral is no harder nor superior in any respect to uvarovite.

Under the microscope garnet crystals show square or hexagonal outlines and an absence of cleavage. They are isotropic and so strongly refracting ( $n$ =from 1.7 to 1.8) as to present a bold relief and shagreenous surface.

Garnet occurs most commonly as crystals, from the size of a pea to that of a boy's marble, in crystalline schists, gneisses, and granulite, and as grains in phonolite and leucitophyre. Sometimes large masses associated with hornblende and magnetite constitute a garnet rock. Garnet, green augite, and hornblende form eclogite, which is by some regarded as the parent rock of diamonds.

Grossularite (Nos. 3486, 3488, 3490), the lime aluminium garnet, is typically found in metamorphic rocks, such as marbles; pyrope, the magnesium aluminium garnet, in basic rocks containing magnesium; almandite, the iron aluminium garnet, and spessartite, the manganese aluminium garnet, in granitic and gneissic rocks; while andradite, the lime iron garnet, is of widespread occurrence.

The Urals, Alps, Pyrenees, Appalachians, and Cordilleras have all furnished multitudes of various kinds of garnets.

#### SUMMARY

*Garnet*.— $(\text{Ca}, \text{Mg}, \text{Fe}, \text{Mn})_2(\text{Al}, \text{Fe}, \text{Cr})_2(\text{SiO}_4)_3$ . Regular; 110, 211. Cleavage (110) imperfect; brittle; fracture uneven.

Hardness=7 to 7.5; gravity=3.8. Honey-yellow to black; luster vitreous; transparent;  $\omega$ =1.7–1.8.

All except uvarovite soluble with difficulty; andradite most readily. Iron garnets fuse to magnetic globules.

Mountain regions generally.

#### Zircon

Zircon, a brown zirconium silicate, nearly always occurs in crystals, since it is one of the first minerals to solidify out of the molten magma which forms augite syenite, elaeolite syenite, and other igneous rocks. In crystalline schists and gneiss the crystals are usually microscopic in size, but may be readily identified by their

square outlines, brown color, and by their high refraction (1.930) and double refraction (0.062), which give them a bold relief and in an ordinary rock section interference colors of the third order. Because of its hardness and insolubility zircon resists decomposition, and is found with gold, platinum, cassiterite, and magnetite in the heavy sands which result from the destruction of granites and gneisses.

Figures 207 and 208 represent the common forms of the simple crystal. By enlargement of the pyramid (311) acute termination results. With zircon, as with cassiterite and quartz, basal planes are rarely developed.

Clear varieties (No. 3494) are much prized as gems because of their hardness (7.5), high refraction, and great dispersion.




FIG. 207.—Zircon




FIG. 208.—Zircon

In refraction and dispersion they come next to diamonds. In color they vary from colorless through various shades of orange, yellow, red ("jacinth"), pale green, and gray ("jargoon"). When dark brown they become opaque. The color is due to  $\text{Fe}_2\text{O}_3$ , and can be altered by heating in the blowpipe flame.

When finely powdered, zircon is slowly decomposed in hot sulphuric acid.

Ceylon, the Urals, Alps, Norway, North Carolina, Arkansas, and Colorado have all furnished fine specimens of zircon.

#### SUMMARY

*Zircon.*— $\text{ZrSiO}_4$ ;  $\text{ZrO}_2 = 67.2$  per cent;  $\text{SiO}_2 = 32.8$  per cent. Tetragonal; holosymmetric.  $a:c = 1:0.64$ . (111), (311), (100), (110). Cleavage parallel (111), (110), imperfect. Brittle; fracture conchoidal.

Hardness=7.5; gravity=4.7. Brown; luster adamantine; subtranslucent;  $\omega=1.93$ . Double refraction positive, strong;  $\epsilon-\omega=0.062$ .

Infusible; insoluble.

Ceylon, Urals, Norway, North Carolina, Colorado.

### Topaz

The name topaz is given to a mineral which is hard, colorless, transparent, prismatic, often vertically striated, usually terminated at one end by several pyramid planes and a basal plane, easily cleavable parallel to the base, and in chemical composition is a silicate of aluminium and fluorine  $\text{Al}_2(\text{FOH})_2\text{SiO}_4$ .




FIG. 209.—Topaz




FIG. 210.—Topaz, showing optic axes (o) and axial plane.

Many crystals are delicately or deeply colored. Pale-blue crystals are found with orthoclase, smoky quartz, and beryl in granite at Nerchinsk, Siberia (No. 1890); and dark-blue crystals at Mursinsk in the Urals in a similar rock accompanied by lepidolite. Deep brown crystals are found on the Urulga River in the Urals, and in Minas Geraes, Brazil (No. 3493). The brown crystals when heated often become pink. Some regions furnish golden-yellow crystals, but the great majority of occurrences are white or colorless. Many limpid, richly planed crystals are found in Utah and Colorado.

Like cassiterite, topaz was formed under the influence of heat and in the presence of vapors containing fluorine.

Attractive color, together with high refraction and great hardness, places topaz among the gems.

## SUMMARY

*Topaz*.— $\text{Al}_2(\text{FOH})_2\text{SiO}_4$ ;  $\text{Al}_2\text{O}_3 = 55.44$  per cent,  $\text{F} = 20.65$  per cent,  $\text{SiO}_2 = 32.61$  per cent. Orthorhombic;  $a:b:c = 0.528:1:0.477$ . (001), (111), (110), (120), (221), (223), (201), (021), (041), (140). Cleavage parallel (001) perfect. Brittle; fracture sub-conchoidal.

Hardness=8; gravity=3.5. Pale yellow; vitreous; transparent.  $\beta = 1.618$ ;  $\gamma = 1.623$ . Double refraction positive, weak;  $\gamma - \alpha = 0.011$ . Axial plane (010). Acute bisectrix normal to (001);  $2E = 114^\circ$ . Dispersion strong;  $\rho > \nu$ .

Infusible; difficultly soluble.

Urals, Brazil, Japan, Utah, Colorado, California, Missouri.

## Cyanite

This mineral (No. 3496), which easily attracts attention because of its blue color (*κυανός*, "blue"), occurs in long, flat, bladed crystals that show a remarkable difference in hardness in different directions. Across the blades, that is, parallel to the edge made by the macropinacoid (100) and base (001), the hardness is 7, while along the crystal, that is, parallel to the edge formed by the macropinacoid (100) and brachypinacoid (010), the hardness is only 4.5. When Haüy discovered this property, he named the mineral disthene (*δίσ* and *θενος*, "double strength"). When cyanite is heated at  $1350^\circ \text{C}$ . without changing its chemical composition ( $\text{Al}_2\text{SiO}_5$ ) it is transformed into a fibrous mineral of uniform hardness (6.5), lighter specific gravity (3.2; cyanite is 3.6), and straight extinction. The mineral is called sillimanite, and is characteristic of some gneisses and schists. Compact sillimanite (sometimes wrongly called jade) was used in prehistoric times in the manufacture of ornaments and implements.

## SUMMARY

*Cyanite*.— $\text{Al}_2\text{SiO}_5$ ;  $\text{Al}_2\text{O}_3 = 63$  per cent,  $\text{SiO}_2 = 37$  per cent. Triclinic;  $a:b:c = 0.899:1:0.697$ . (001), (100), (010), (110). Cleavage parallel (100) perfect; (010) imperfect. Brittle; fracture fibrous.

Hardness=7 across the crystal, 4.5 parallel to the edge (100); (010); gravity=3.6. Blue to white; vitreous; transparent.  $\beta = 1.72$ . Double refraction negative;  $\gamma - \alpha = 0.016$ . Axial plane inclined  $30^\circ$  to edge (100); (010). Acute bisectrix normal to 100:2  $H = 100^\circ$ .

Infusible; insoluble.

Alps, northern England, Appalachians, Cordilleras.

### Tourmaline

This mineral is worthy of notice for three reasons: first, because it is abundant in igneous and metamorphic rocks; second, because it is used in making optical instruments such as "tourmaline tongs" (see below, p. 166); and third, because the beautiful red, pink, and green varieties are used as gems (Plate XXVIII). Tourmaline is literally found "from Maine to California." Paris, Maine (Nos. 444, 454, 4062), has long been reputed for its magnificent red and green crystals, and more recently San Diego County, California (Nos. 3511, 3788, 3789, 3790), has furnished the museums of the world with handsome groups of red tourmaline (rubellite) in a lavender mica, lepidolite. The tourmalines of Illinois are all emigrants from northern




FIG. 211.—Tourmaline




FIG. 212.—Tourmaline

regions. Black is the prevailing color, and they are usually imbedded in gneisses and granites. Among the most famous foreign localities may be mentioned the region near Ekaterinburg in the Urals, where a coarse granite contains smoky quartz, albite, green and pink mica, and red and fine black tourmaline. At Campolagna, Switzerland, calcite, corundum, diaspore, mica, and green tourmaline are found in a granular dolomite. The granite of the island of Elba consists of quartz, orthoclase, albite, mica, pink beryl, red garnet, and red and black tourmaline. The best gem tourmalines are obtained in Ceylon.

Tourmaline crystals (Nos. 357 and 3505) (Figs. 211 and 212) are usually prismatic, often elongated, sometimes terminated at one end,



PLATE XXVIII



*a*, Tourmaline doubly terminated; variously colored crystal from Mesa Grande, California.



*b*, Black, well-crystallized specimen from Haddam, Connecticut.

rarely at both ends. Occasionally they are flat crystals. Prisms are strongly striated vertically. This striation, the triangular cross-sections, and absence of cleavage serve to distinguish this mineral from black pyroxenes and amphiboles.

The chemical constitution of tourmaline is complex and is still the subject of much discussion. Generally speaking, it is a borosilicate of aluminum, iron, or chromium, of magnesium, and of the alkalis sodium, potassium, and lithium. The following varieties may be distinguished:

Black, iron tourmaline ( $\text{Fe}_4\text{Na}_2\text{B}_6\text{Al}_{14}\text{H}_3\text{Si}_{12}\text{O}_{63}$ ); gravity = 3.2.

Red, green, colorless; alkali tourmaline ( $\text{NaLiK})_4\text{B}_6\text{Al}_{16}\text{H}_3\text{Si}_{12}\text{O}_{63}$ ); gravity = 3.

Brown; colorless; magnesium tourmaline ( $\text{Mg}_{12}\text{B}_6\text{Al}_{10}\text{H}_3\text{Si}_{12}\text{O}_{63}$ ); gravity = 3.

Green; chromium tourmaline (chromium replacing a portion of the aluminium); gravity = 3.1.

Transparent crystals (No. 3788) are often differently colored at the different ends, and some are banded with two or three different shades of color, as may be observed in a section parallel to the base.

Early in the eighteenth century it was discovered that red tourmaline crystals brought from Ceylon when heated became positively electrified at one end and negatively at the other. When any tourmaline crystals after heating are beginning to cool, if they are dusted with finely powdered red lead (+) and sulphur (−), one end—the negative, the “analogous end”—attracts the red lead, while the other—the positive or “antilo-  
gous end”—attracts the sulphur.

The negative end (No. 3791) usually shows a basal plane and the rhombohedron ( $R$ ) over the trigonal prism ( $10\bar{1}0$ ) (Fig. 213). The positive end is usually acute owing to the development of pyramids. All tourmalines absorb the ordinary ray much more completely than they do the extraordinary, consequently black varieties look green or




FIG. 213.—Tourmaline, analogous end.



blue with the ordinary ray ( $\sigma$ ) and brown or red with the extraordinary ( $\epsilon$ ). Brown crystals cut parallel to the optic axis transmit only the extraordinary ray, and can be used as polarizer and analyzer. Two such sections held in wire rings constitute "tourmaline tongs."

#### SUMMARY

*Tourmaline.*— $(\text{FeCrNaKLi})_4\text{Mg}_{12}\text{B}_6\text{Al}_6\text{H}_6\text{Si}_{12}\text{O}_{63}$ . Hexagonal; symmetry ditrigonal polar;  $a:c=1:0.4477$ .  $(0001)$ ,  $(10\bar{1}1)$ ,  $(02\bar{2}1)$ ,  $(10\bar{1}0)$ ,  $(11\bar{2}0)$ ,  $(32\bar{5}1)$ . Cleavage imperfect parallel  $(10\bar{1}1)$ . Brittle; fracture subconchoidal.

Hardness=7; gravity=3.1. Black, brown, red, green, colorless; vitreous; translucent;  $\omega=1.64$ . Double refraction strong, negative;  $\omega-\epsilon=0.017$ . Pleochroic; pyro-electric.

Fusible; insoluble.

Maine to California, Urals, Alps.

#### ZEOLITE GROUP

The zeolites ( $\zeta\epsilon\omega$ , "I boil") are white, pearly, light (gravity, 2), soluble minerals, which boil before the blowpipe because the two to




FIG. 214.—Four-twinning crystals of stilbite.



FIG. 215.—Stilbite sheaf

six molecules of water of crystallization contained in them are loosely held. The different members of the group are so-called "secondary minerals" since they result from the decomposition of other minerals, chiefly feldspar, leucite, etc., in disintegrating igneous rocks. From seventeen representatives three may be chosen to indicate the nature of the group.

#### Stilbite

The most attractive characteristic about this mineral is its pearly sheaflike crystals (Nos. 3255 and 3514), which result from twinning

of monoclinic crystals parallel to the base and orthodome in such a manner as to produce interpenetrating twins. The crystals are flattened parallel to the clinopinacoid, which is also a plane of easy cleavage. The basal cleavage is imperfect. The basaltic rocks found in many places in New Jersey, Michigan, the Cordilleras, Scotland, etc., contain in their cavities fine groups of stilbite crystals.

## SUMMARY

*Stilbite*.— $\text{CaAl}_2\text{Si}_4\text{O}_{16} + 6\text{H}_2\text{O}$ ;  $\text{CaO} = 8.94$  per cent,  $\text{Al}_2\text{O}_3 = 16.31$  per cent,  $\text{SiO}_2 = 57.51$  per cent,  $\text{H}_2\text{O} = 17.24$  per cent. Some  $\text{Na}_2$  usually replaces a portion of the Ca. Monoclinic;  $a:b:c = 0.7623:1:1.1940$ .  $\beta = 50^\circ 50'$  (001), (010), (110). Twinned parallel (001) and ( $\bar{1}01$ ). Cleavage parallel (001). Brittle; fracture uneven.

Hardness = 3.5; gravity = 2.2. White; vitreous; transparent;  $\beta = 1.498$ . Double refraction, strong, negative;  $\gamma - \alpha = 0.006$ . Axial plane (010) acute bisectrix inclined  $85^\circ$  to normal of (001) and  $34^\circ$  to the normal of (100).  $2E = 51.5$ .

Fusible (2.5). Decomposed by hydrochloric acid.

In disintegrating igneous rocks in the Cordilleras, Appalachians Scotland, etc.

## Analcite

The second representative of the zeolites to claim our attention is analcite (No. 3576), one of the best illustrations of the trapezohedral crystals among minerals (Fig. 216). More rarely analcite occurs in cubes with corners truncated by the trapezohedrons. Small crystals are often beautiful and glossy. The larger ones are usually opaque and white or pink.

## SUMMARY

*Analcite*.— $\text{Na}_2\text{Al}_2\text{Si}_4\text{O}_{12} + 2\text{H}_2\text{O}$ ;  $\text{Na}_2\text{O} = 14.1$  per cent,  $\text{Al}_2\text{O}_3 = 23.2$  per cent,  $\text{SiO}_2 = 54.5$  per cent,  $\text{H}_2\text{O} = 8.2$  per cent. Regular; (211), (100). Brittle; fracture uneven.

Hardness = 5.5; gravity = 2.2. Colorless; vitreous; transparent.  $n = 1.487$ .

Fusible (2.5). Gelatinizes in hydrochloric acid.

Same regions as other zeolites.




FIG. 216.—Leucite.

### Natrolite

Natrolite (No. 3517) is closely related to analcite in chemical composition inasmuch as it contains one less molecule of  $\text{SiO}_2$ , but differs markedly in form since it crystallizes in the orthorhombic system and occurs in long prisms that end in very flat pyramids.

It is the commonest of fibrous zeolites, usually constituting masses in cavities. Beautiful tufts of acicular crystals are found in the cavities of basalt.

So fusible is it that it melts in a candle flame, imparting the yellow color characteristic of burning sodium.




FIG. 217.—Natrolite

### SUMMARY

*Natrolite*.— $\text{Na}_2\text{Al}_2\text{Si}_3\text{O}_{10} \cdot 2\text{H}_2\text{O}$ ;  $\text{Na}_2\text{O} = 16.32$  per cent,  $\text{Al}_2\text{O}_3 = 26.86$  per cent,  $\text{SiO}_2 = 47.36$  per cent,  $\text{H}_2\text{O} = 9.46$  per cent. Orthorhombic;  $a:b:c = 0.978:1:0.354$ . (110), (111). Cleavage parallel (110) perfect. Brittle; fracture uneven.

Hardness = 5.5; gravity = 2.2. Colorless; vitreous; transparent.  $\beta = 1.479$ . Double refraction positive, strong;  $\gamma - \alpha = 0.012$ . Axial plane (010). Acute bisectrix normal to (001).  $2E = 99^\circ$ ;  $\rho < \nu$ .

Fuses readily (2). Gelatinizes in hydrochloric acid.

In basalts in the Cordilleras, Michigan, New Jersey, etc.

### MICA GROUP

While zeolites are comparatively rare, the members of the mica group are among the most abundant, well-known, and useful of minerals. The thin, flexible, elastic leaves into which mica may be separated distinguish it so clearly that once seen it is not forgotten.

From the seven members of the group the three most striking and abundant are white mica, muscovite, "isinglass"; black mica, biotite; and lilac mica, lepidolite.

Together with quartz and feldspar, micas are common in the granites, gneisses, and schists. The minerals of this group are always crystallized and never massive. The crystals vary from minute flakes to immense sheets which measure sometimes several feet across. In some localities in Russia, India, Canada, and the United States, deposits are being mined which furnish sheets of large size.

The crystals are flat monoclinic prisms having six sides and often so regularly shaped as to appear to belong to the hexagonal or orthorhombic system (Fig. 218). Accurate measurements and optical investigations, however, reveal their monoclinic symmetry. The angles between the prism planes are always about  $120^\circ$ .

The micas are all silicates of aluminium and of either potassium, sodium, and lithium—the alkali micas—or of iron and magnesium—the ferro-magnesium mica; and contain also fluorine and hydrogen.




FIG. 218.—Muscovite




FIG. 219.—Muscovite: pressure figure dotted; percussion figure solid; optic axes.

Inclusions of other minerals, such as hematite, tourmaline, garnet, etc., are common and are arranged as flat scales along definite lines marked out by the so-called glide planes. When minute they often produce the attractive property known as asterism.

### Muscovite

White mica is called muscovite, from Moscow, where it and a substance resembling it in appearance—gelatin, derived from the sturgeon so abundant in Russian rivers—were long used for window panes and other purposes. It is the potassium mica,  $\text{H}_2\text{KA}_3(\text{SiO}_4)_3$ . Its crystals often attain large size. They are six-sided, rough-faced, and taper because of the dome planes (Nos. 3519, 3520, 378, 1281). Besides the excellent cleavage parallel to the base (001), no other cracks are apt to be found in thin sections. But by pressure with a blunt-pointed instrument three sets of cracks (the dotted line in Fig. 219), the so-called pressure figures, are formed. The cracks

correspond to the glide planes which are parallel to the clinodome ( $\bar{2}05$ ) and the pyramid ( $135$ ). When developed in nature these cracks divide the crystal into trigonal pieces.

The percussion figure produced by striking a cleavage plate with a sharp-pointed instrument consists of a six-rayed star in which the rays intersect at angles of  $60^\circ$  (the solid lines in Fig. 219). The most prominent crack is parallel to the brachypinacoid ( $010$ ), which is the plane of symmetry.

In muscovite the axial plane is perpendicular to the principal crack of the percussion figure and hence parallel to the macropinacoid ( $100$ ). Hence muscovite is called "macrodiagonal mica" or mica of the first class.

Muscovite is formed both in fused magmas and in aqueous solutions.

Its uses are many: by the Russians for windows in war vessels, by the French for decoration and ornamentation, by the Anglo-Saxons for various commercial purposes. It furnishes doors for stoves and furnaces. It is used for electrical purposes. It serves as a non-conductor of heat and electricity. It is an absorbent of nitroglycerine; it is a lubricant. In short, it is a mineral much used by man.

#### SUMMARY

*Muscovite*.— $H_2KAl_3(SiO_4)_3$ ;  $K_2O=11.8$  per cent,  $Al_2O_3=38.5$  per cent;  $SiO_2=45.2$  per cent,  $H_2O=4.5$  per cent. Monoclinic;  $a:b:c=0.5777:1:3.312$ .  $\beta=89^\circ 54'$ . ( $001$ ), ( $111$ ), ( $221$ ), ( $110$ ), ( $010$ ). Twins parallel ( $110$ ) are combined on the base ( $001$ ). Cleavage parallel ( $001$ ). Elastic; fracture uneven; chief percussion figure parallel to ( $010$ ).

Hardness= $2.5$  per cent; gravity= $2.86$ . Axial plane perpendicular to ( $010$ );  $\beta=1.6$ . Double refraction negative, strong;  $\gamma-a=0.039$ ;  $\rho>\nu$ . Pleochroism feeble; transparent, white; vitreous.

Fusible with difficulty; insoluble.

In granites, gneisses, mica schists, in all mountain ranges. New Hampshire, South Carolina, South Dakota, Colorado, New Mexico, and California.

#### Biotite

Black mica, named biotite after Biot, the celebrated French mineralogist, is the magnesian mica  $(H,K)_2(Mg,Fe)_2Al_2(SiO_4)_2$  (Nos. 3535, 3536, 1240, 1768, 453).

In this mica the axial plane is usually parallel to the chief percussion figure, which, as noted above, is parallel to the brachypinacoid (010). Such biotite is said to be a brachydiagonal mica, or mica of the second class.

Whether muscovite or biotite is the more abundant mica is difficult to say, since both abound in nearly all kinds of igneous rocks. Biotite decomposes more readily than muscovite and forms such minerals as chlorite, epidote, quartz, and iron oxide. Its




FIG. 220.—Biotite; axial plane parallel to (010).




FIG. 221.—Basal section of biotite, showing position of axial plane and percussion figure.

characteristic color is black, but while undergoing decomposition it assumes red and green shades. It is strongly pleochroic and apparently uniaxial. Fine crystals are found at Vesuvius.

#### SUMMARY

**Biotite.**— $(\text{H,K})_2(\text{Mg,Fe})_2\text{Al}_2(\text{SiO}_4)_3$ ;  $\text{K}_2\text{O}=7.64$  per cent,  $\text{MgO}=21.89$  per cent,  $\text{H}_2\text{O}=4.02$  per cent,  $\text{F}=0.89$  per cent,  $\text{Fe}_2\text{O}_3=7.86$  per cent,  $\text{Al}_2\text{O}_3=16.95$  per cent,  $\text{SiO}_2=39.30$  per cent. A little  $\text{FeO}$ ,  $\text{MnO}$ ,  $\text{CaO}$ , and  $\text{Na}_2\text{O}$  are usually present. Monoclinic;  $a:b:c=0.577:1:3.274$ .  $\beta=90^\circ$ . (110), (111), (001), (010), (221). Twinning plane parallel to (110), combination face (001). Cleavage parallel to (001) perfect; laminae elastic; fracture uneven.

Hardness=2.5; gravity=2.86. Black, pleochroic;  $\alpha$  green,  $\beta$  and  $\gamma$  dark brown. Translucent; luster vitreous; streak colorless. Plane of optic axes parallel to (010). Acute bisectrix inclined  $30'$  to the perpendicular of (001).  $\beta=1.6$ . Double refraction negative, strong;  $\gamma-\alpha=0.04$ ;  $\rho < \nu$ ;  $2E=12^\circ 48'$ .

Fusible with difficulty; soluble in sulphuric acid.

Maine, North Carolina, Colorado.

### Lepidolite

This beautiful violet mica (No. 553) occurs in small flakes only (*λεπίς*, "scale"). It is usually accompanied with other lithium minerals such as tourmaline. The violet color is probably due to manganese. San Diego, California, has within the last few years supplied most of the museums of the country with fine specimens of lepidolite.

#### SUMMARY

*Lepidolite*.— $\text{Li}_2\text{K}_2\text{F}_2\text{Al}_3\text{H}_2\text{Si}_6\text{O}_{20}$ ;  $\text{Li}_2\text{O}$ =1 to 6 per cent,  $\text{K}_2\text{O}$ =4 to 10 per cent,  $\text{F}$ =2 to 8 per cent,  $\text{Al}_2\text{O}_3$ =26 to 33 per cent,  $\text{SiO}_2$ =49 to 52 per cent. Monoclinic; crystallographic and optical properties similar to those of the other micas, but the axial plane is sometimes parallel and sometimes perpendicular to the plane of symmetry.

Hardness=2.5; gravity=2.8. Lilac or rose-colored.

Partly soluble in hydrochloric acid.

Maine, Massachusetts, Connecticut, California.

### SERPENTINE AND TALC GROUP

These two magnesian silicates are common decomposition products of other ferro-magnesian silicates. They are basic salts and not hydrated, inasmuch as they part with their water at high temperatures only. Like most secondary minerals they are soft. Crystallization is inconspicuous. They occur most commonly in compact masses in veins or beds.

#### Serpentine

Serpentine is a green, red, or yellow mineral often more or less fibrous in structure, as may be seen under the microscope in those masses which fill veins. At times the fibers are well pronounced and silky in luster, and the mineral is then called chrysotile—the most abundant asbestos (No. 3540). The fibers are longer, more tenacious and silky than those of the amphibole asbestos. The most productive asbestos mines in North America, those in Megantic and Beauce counties, Quebec, furnish the chrysotile variety of asbestos. Arizona, California, and Wyoming are furnishing small quantities.

Some serpentines supply ornamental stones of great beauty (Nos. 4336, 4337). The permanent dark-green color is rendered even more attractive by white particles of magnesite and talc, and

by splashes of blood-red iron stains. "Verde antique" is a brecciated serpentine.

## SUMMARY

*Serpentine.*— $\text{H}_4\text{Mg}_3\text{Si}_2\text{O}_9$ ;  $\text{MgO}=43.5$  per cent,  $\text{SiO}_2=43.5$  per cent,  $\text{H}_2\text{O}=13$  per cent. Massive; fracture splintery.

Hardness=3; gravity=2.6. Green; luster greasy; translucent;  $\beta=1.57$ . Double refraction negative, weak;  $\gamma-\alpha=0.010$ .

Fusible with difficulty (6). Soluble in hydrochloric acid.

Appalachians, Cordilleras, Alps, etc.

## Talc

Talc is a foliated, silvery, soft, greasy mineral. Though the flakes show hexagonal outline, yet, like mica, talc is monoclinic. The angle between the optic axes is very small. In many regards talc resembles one of the micas, phlogopite, but is softer and not elastic (Nos. 353, 368). Talc usually contains from 1 to 4 per cent of iron oxide.

The compact varieties, such as steatite or soapstone, were used by the Chinese in ancient times for ornaments and images, and by savage and civilized peoples today. A clearer conception of the value of the mineral industry in the United States may be gained by the knowledge that while talc is a mineral rarely mentioned, more than a million dollars' worth of it are produced in the United States annually and manufactured into such articles as bath and laundry tubs, sinks, mantles, hearthstones, fire brick, griddles, slate pencils, gas tips, crayons, French chalk for tailors, adulterant for sugars, lubricators for dressing skins and leather, toilet powders, dynamite, paper-making, pigments in high-grade paints, etc. New York is the leading state in the production of talc and Virginia in soapstone.

## SUMMARY

*Talc.*— $\text{H}_2\text{Mg}_3(\text{SiO}_3)_4$ ;  $\text{MgO}=31.7$  per cent,  $\text{SiO}_2=63.5$  per cent,  $\text{H}_2\text{O}=4.8$  per cent. Monoclinic. Cleavage parallel (001) perfect; sectile, pliable.

Hardness=1; gravity=2.7. Color silver white; luster pearly; translucent.  $\beta=1.55$ . Double refraction negative, strong;  $\gamma-\alpha=0.040$ . Acute bisectrix normal to the cleavage.  $2E=19^\circ$ .

Almost infusible. Insoluble.

The Alps, Appalachians, Cordilleras, etc.



### Kaolinite

Decomposition of orthoclase, albite, leucite, beryl, and other minerals often gives rise to a secondary mineral, kaolinite, which is a hydrated aluminous silica,  $H_4Al_2Si_2O_9$ . The pure form is kaolinite (from the Chinese *kaoling*, "high ridge"). As various impurities such as iron, calcium, and magnesium increase, various clays result. In structure and other physical characters kaolinite resembles the hydrated magnesian silicate serpentine. It is white, scaly, flexible, inelastic, soft (hardness, 2), light, (gravity, 2.6), unctuous, and plastic.

Some clays have absorbent properties which render them valuable as fuller's earth. Others fuse at such temperatures and yield a product of such character as to be valuable for porcelain, china, tile, brick, etc. Others are plastic because of the elongated particles which constitute them, and are useful in clay-modeling.

Various varieties of clays are among the most abundant mineral constituents of the regolith covering Illinois. They form soil for agriculture and plastic material for the manufacture of tile, porcelain, brick, etc. Their contribution thus to the wealth of the state is difficult to estimate. The value of clay products sold annually in this state amounts to more than fifteen million dollars, but this takes no account of the return through soil fertility. Clay soils are useful chiefly in furnishing a binder for more porous soil, as retaining moisture, and as being a container for other plant food.

### SUMMARY

*Kaolinite*.— $H_4Al_2Si_2O_9$ ;  $Al_2O_3=39.5$  per cent,  $SiO_2=46.5$  per cent,  $H_2O=14$  per cent. Water is driven off at  $330^\circ$ . Monoclinic. Scales flexible, inelastic; friable to compact; unctuous, plastic.

Hardness=2.5; gravity=2.6. White, blue, yellow, red, green; luster pearly to earthy; translucent. Biaxial, negative.

Infusible; insoluble. Blue color with cobalt solution.

Many eastern and middle states, such as Massachusetts, Delaware, Georgia, Illinois, etc.



PLATE XXIX



*a*, Apatite, Renfrew, Canada



*b*, Barite, Alston Moor, England

## CLASS VIII. NIOBATES, TANTALATES

## CLASS IX. PHOSPHATES, ARSENATES, VANADATES, ANTIMONATES, NITRATES

### Apatite

Apatite is a mineral of great commercial importance, occurring in metamorphic limestones and in granites often as well-formed crystals (No. 3551) varying from microscopic size to dimensions of a foot or more (Plate XXIXa). As usual, the more nearly perfect crystals with well-terminated ends occur in cavities. Their prevailing color is blue, green (No. 3550), yellow, or brown. Among the most beautiful apatite crystals found are little limpid hexagonal prisms contained in crystalline schists in the St. Gothard and Untersulzbachthal. Microscopic crystals are found in a variety of igneous rocks. The planes most commonly appearing (Fig. 222) are the following:  $(10\bar{1}0)$ ,  $(11\bar{2}0)$ ,  $(10\bar{1}1)$ ,  $(11\bar{2}1)$ ,  $(0001)$ . The prisms are usually vertically striated. Apatite resembles beryl in appearance, but is softer, has imperfect cleavage parallel to the base, and a high, refractive index.

Chemically there are two varieties of apatite: the ordinary, which contains fluorine; and the less common, in which fluorine is replaced by chlorine. According to physical condition there are two kinds which are even more markedly different than are the two chemical varieties. The first is pure crystallized apatite, which is found filling veins and as inclusions in metamorphic rocks (Nos. 3666, 3709, 3712). The second is phosphorite (No. 4307), the white, structureless variety, organic in origin and occurring in extensive beds in the Carolinas and Tennessee. It has resulted by the concentration of phosphatic material which was previously disseminated through sands and sandstone.




FIG. 222.—Apatite

The crystallized apatite is found in most of the Appalachian states and in the drifts over the middle states in granular limestone and in granites, gneisses, and schists, and in veins in iron ores.

Apatite is one of the most important of all minerals to man, inasmuch as it is the chief source of phosphorus, a chemical substance indispensable to plant growth.

#### SUMMARY

*Apatite*.— $\text{Ca}_5\text{F}(\text{PO}_4)_3$ ;  $\text{CaO}$ =55.5 per cent,  $\text{P}_2\text{O}_5$ =42.3 per cent,  $\text{F}$ =3.8 per cent. Hexagonal; symmetry hexagonal equatorial;  $a:c=1:0.7346$ .  $(10\bar{1})$ ,  $(10\bar{1}1)$ ,  $(11\bar{2}1)$ ,  $(21\bar{3}1)$ . Cleavage parallel to  $(0001)$ ,  $(10\bar{1}0)$ . Brittle; fracture conchoidal.

Hardness=5; gravity=3.2. Colorless; luster vitreous; transparent;  $\omega=1.646$ . Double refraction negative, weak;  $\omega-\epsilon=0.004$ .

Fusible with difficulty; soluble in hydrochloric acid.

Ottawa County, Quebec, Canada; Bolton, Massachusetts; Crown Point, New York; New Jersey.

#### Pyromorphite

Pyromorphite (No. 507) is a lead chloro-phosphate found in quantities in upper levels of lead mines, where it has been forming during the decomposition of lead sulphide. It was named pyromorphite because, when fused before the blowpipe, upon cooling it solidifies with many facets ( $\pi\upsilon\rho$ , "fire";  $\mu\omicron\rho\phi\eta$ , "form"). These facets are not true crystal faces. The true crystals are composed of prisms and basal planes which produce barrel-shaped forms because of aggregation and curvature of the prisms. A violet color is shown by large hexagonal prisms occasionally, but the prevailing color is green or brown.

#### SUMMARY

*Pyromorphite*.— $\text{Pb}_5\text{Cl}(\text{PO}_4)_3$ ;  $\text{PbO}$ =82.3 per cent,  $\text{P}_2\text{O}_5$ =15.7 per cent,  $\text{Cl}$ =2.6 per cent.

Hexagonal; symmetry hexagonal equatorial.  $(1010)$ ,  $(10\bar{1}1)$ ,  $(0001)$ . Cleavage parallel  $(10\bar{1}0)$ ,  $(10\bar{1}1)$  imperfect. Brittle; fracture sub-conchoidal.

Hardness=3.5; gravity=7. Green; luster resinous. Translucent;  $\omega=1.50$ . Double refraction negative, weak;  $\omega-\epsilon=0.006$ .

Easily fusible (1.5). Soluble in nitric acid.

Missouri, Wisconsin, Colorado, New Mexico, and Australia.

## CLASS X. BORATES, URANATES

### Borax, Colemanite, Boracite

These three borates are of importance as the source of boron compounds which are useful as antiseptics, medicines, cosmetics, and welding compounds. All of them are most commonly met with in arid regions in connection with salt lakes, past or present. Crystals of colemanite and boracite are often beautiful because of their transparent character and lustrous surfaces.

#### SUMMARY

*Borax*.— $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$ ;  $\text{Na}_2\text{O}$ =16.2 per cent,  $\text{B}_2\text{O}_3$ =36.6 per cent,  $\text{H}_2\text{O}$ =47.2 per cent. Monoclinic, prismatic class;  $a:b:c=1.09:1:0.562$ ;  $\beta=73^\circ$ ; (100), (110), (001), (111). Cleavage perfect (100); fracture conchoidal.

Hardness=2; gravity=1.7. White, vitreous, translucent;  $\beta=1.47$ . Double refraction negative;  $\gamma-a=0.004$ ; acute bisectrix normal to (010);  $2E=59^\circ$ ;  $\rho>\nu$ .

Fusible, swells up; soluble in water; sweetish.

Thibet, Peru, California, Nevada.

*Colemanite*.— $\text{Ca}_2\text{B}_6\text{O}_{11} \cdot 5\text{H}_2\text{O}$ ;  $\text{CaO}$ =27.2 per cent,  $\text{B}_2\text{O}_3$ =50.9 per cent,  $\text{H}_2\text{O}$ =21.9 per cent. Monoclinic, prismatic class;  $a:b:c=0.77:1:0.541$ ;  $\beta=70^\circ$ ; (110), (301), (100), (010), (001), (111), (021), (221). Cleavage (010); fracture uneven.

Hardness=4; gravity=2.4. Colorless, white; translucent, vitreous;  $\beta=1.5902$ .

Fusible, exfoliates; soluble in hot hydrochloric acid; insoluble in water.

California, Chile.

*Boracite*.— $\text{Mg}_3(\text{MgCl})_2\text{B}_{10}\text{O}_{30}$ ;  $\text{MgO}$ =31.4 per cent,  $\text{Cl}$ =7.9 per cent,  $\text{B}_2\text{O}_3$ =62.5 per cent. Dimorphous; crystals formed above  $265^\circ\text{C}$ ., regular; below that, orthorhombic; (110), (100), (111), (211). Cleavage (111) imperfect; brittle; fracture conchoidal.

Hardness=7; gravity=3. Colorless, vitreous, translucent. Double refraction;  $\beta=1.667$ ;  $\gamma-a=0.011$ ;  $2V=90^\circ$ .

Fusible, swells up; soluble in hydrochloric acid.

Stassfurt, Prussia.

### Uraninite

The uranate uraninite is of interest because it is a source of uranium, of radium, and of helium. Its composition is doubtful, inasmuch as a large number of rare elements are present. In addition to oxides of uranium, thorium, lead, iron, and calcium, small quantities of the following have been found: zirconium, cerium, lanthanum, didymium, yttrium, erbium, helium, manganese, sodium, potassium, silicon, phosphorus, and hydrogen. Its composition may be expressed by the formula  $U_3O_8$ . Uranium compounds are used in the laboratory for the determination of phosphorus and zinc, in the manufacture of pigments, glazes, and special steels.

### SUMMARY

*Uraninite*.— $U_3O_8$ . Regular, (111), (110), (100). Crystals rare, crystalline masses, botryoidal groups. Brittle; fracture conchoidal.

Hardness = 5.5; gravity = 9.5. Brown, black; luster dull.

Infusible; soluble in nitric and sulphuric acids; radio-active.

Colorado, Cornwall, Austria.

## CLASS XI. SULPHATES, CHROMATES, TELLURATES

Class XI, containing the sulphates, chromates, and tellurates, is an outstanding class because of at least four commercially and scientifically interesting minerals, namely, barite, celestite, anglesite, and gypsum.

### Barite

Barite, or heavy spar (*βαρύς*, "heavy"), so named since it is nearly twice as heavy as other white minerals like calcite or gypsum, is important because of its fine crystals, its great masses, and its usefulness.




FIG. 223.—Barite



FIG. 224.—Barite




FIG. 225.—Barite

The crystals are usually flat (Nos. 3562 and 3556), and consist of large basal planes with short prisms, as in Figure 223. Forms composed of dome planes elongated parallel to the *a* axis (Fig. 224) are not uncommon (No. 4060). Cleavage pieces take the form of Figure 225, and the position of the axes is indicated by the cleavage, which usually shows pearly cracks. Prismatic cleavage is good. Aggregates of crystals produce rounded masses from which acute prism edges protrude. Radiated, columnar, and massive (No. 3559) forms are common, though a white, earthy, massive condition is most characteristic. Discoloration by iron is usual.

Inorganic phosphorescence was first discovered when an Italian investigator in the early part of the seventeenth century heated



barite on charcoal and noticed that in the dark it continued to emit a glow, due to the reduction of the sulphate to sulphide.

Barite is found in veins and masses with ores of lead, antimony, and iron in limestones, especially in Georgia, Missouri, and Tennessee.

It is used in the manufacture of white paint, filler for paper, barium for chemical and medicinal uses, etc. Nearly four hundred thousand dollars' worth of barite was produced in the United States in 1915.

#### SUMMARY

*Barite*.— $\text{BaSO}_4$ ;  $\text{BaO}$ =65.7 per cent,  $\text{SO}_3$ =34.3 per cent. Orthorhombic;  $a:b:c=0.815:1:1.314$ . (001), (110), (102), (011), (122), (111). Cleavage parallel (110) and (001) perfect; brittle; fracture uneven.

Hardness=3; gravity=4.5. Colorless; luster vitreous; transparent;  $\beta=1.637$ . Double refraction positive, strong;  $\gamma-\alpha=0.012$ . Axial plane (010); acute bisectrix perpendicular to (100);  $2E=64^\circ$ ;  $\rho < v$ .

Fusible (3) with decrepitation; insoluble in acid.

Georgia, Missouri, Tennessee, Kentucky.

#### Celestite

The next member of the group is the strontium sulphate, celestite, so named (coelestinus, "blue") since the first crystals discovered (in

Pennsylvania) exhibited delicate blue shades, due, no doubt, to the presence of traces of iron phosphate. The crystals are inclosed by a variety of planes. While barite is more often elongated along the  $a$  axis, the elongation of celestite takes place parallel to the  $b$  axis. The massive forms are common in limestone, marl, sandstone, and beds of gypsum.




FIG. 226.—Celestite (001), (104), (102), (110), and (011).

Sicily; Strontian, Scotland; North Bass Island, Lake Erie; Pennsylvania, Kansas, Texas, West Virginia, and Tennessee contain supplies of this mineral, which together with strontianite are the chief sources of strontium nitrate, a compound much used to produce the crimson colors in fireworks.

## SUMMARY

*Celestite*.— $\text{SrSO}_4$ ;  $\text{SrO}=56.4$  per cent,  $\text{SO}_3=43.6$  per cent.

Orthorhombic;  $a:b:c=0.779:1:1.280$ . (001), (110), (011), (102), (104). Cleavage parallel (001) perfect; parallel (110) good. Brittle; fracture uneven.

Hardness=3; gravity=3.9. Colorless; luster vitreous; transparent;  $\beta=1.624$ . Double refraction positive, weak;  $\gamma-a=0.009$ . Axial plane parallel (010). Acute bisectrix perpendicular to (100).  $2E=88^\circ 38'$ .

Fusible (3) with decrepitation. Insoluble in acids.

Lake Erie, Pennsylvania, New York, Kansas, Texas, West Virginia, Tennessee.

## Anglesite

Lead sulphate ( $\text{PbSO}_4$ ) crystals resemble barite and celestite in being flat (No. 3563). They are elongated not only parallel to the  $b$  axis but also quite commonly parallel to the  $c$ . They were first




FIG. 227.—Anglesite




FIG. 228.—Anglesite

found on the island of Anglesy, and many localities now furnish fine, lustrous, transparent, colorless crystals which line the cavities in glistening, granular galena. The lead mines of Missouri, Wisconsin, Kansas, Colorado, Mexico, and Australia furnish this mineral in massive varieties and in such quantities as to render it an important ore of lead.

Its easy fusibility (1.5), adamantine luster, and great weight (gravity, 6.3) render it easy of determination.

## SUMMARY

*Anglesite*.— $\text{PbSO}_4$ ;  $\text{PbO}=73.6$  per cent,  $\text{SO}_3=26.4$  per cent. Orthorhombic;  $a:b:c=0.785:1:1.289$ . (110), (001), (011), (102), (104), (122), (111), cleavage parallel (110) and (001) fair. Brittle; fracture conchoidal.

Hardness=3; gravity=3.6. Colorless; luster adamantine; transparent;  $\beta=1.883$ . Double refraction positive, strong;  $\gamma-a=0.016$ . Axial plane parallel (010). Acute bisectrix perpendicular to (100).  $2H=89^\circ 52'$ ;  $\rho < \nu$ .

Easily fusible (1.5). Soluble in nitric acid with difficulty. Missouri, Wisconsin, California, Mexico, and Australia.

Barite, celestite, and anglesite constitute a fine example of an isomorphous group, with simple, bright, glassy, tabular or prismatic orthorhombic crystals which cleave parallel to the base (001) and prism (110). The optical characteristics are all similar.

### Gypsum

The next sulphate of importance is the hydrated calcium sulphate gypsum. This is a mineral vastly more abundant than all the other members of the group combined. The name gypsum was used by the Greeks ( $\gamma\upsilon\psi\upsilon\sigma$ ). Dioscorides and Pliny called it "selenites," from which our word "selenite" is derived, which is now restricted




FIG. 229.—Gypsum




FIG. 230.—Gypsum




FIG. 231.—Gypsum

to lustrous, satiny, crystallized gypsum. Near Paris (Montmartre) gypsum was early quarried, ground, and burned for plaster and hence was named plaster of Paris. When used upon the fields as a fertilizer it is called land plaster. When translucent, compact, and suitable for carving, it is called alabaster. Satin spar is composed of compact fibers, with the luster of satin, and is used for cheap jewelry. It is



PLATE XXX



Gypsum, showing fishtail twin and curled form

PLATE XXXI



Gypsum, "selenite," Wayne County, Utah



easy to work but even more easy to destroy, since it is so soft that it has slight value.

The chief use of gypsum, however, is as plaster of Paris. When ground and burnt, it loses its water of crystallization, then upon being mixed with water again it takes up the lost molecules, and recrystallizes or "sets" ( $\text{CaSO}_4 \cdot \text{H}_2\text{O}$ ).

The greatest quantities are now mined in Michigan (No. 3523), New York, Virginia, Ohio, Iowa, Alabama, Arkansas. For hundreds




FIG. 232.—Gypsum, twinned by juxtaposition parallel (100).




FIG. 233.—Gypsum, twinned by interpenetration parallel (100).

of miles beds of gypsum may be seen stretching like a white ribbon over the country in Wyoming, Colorado, and many other Cordilleran states.

Crystals sometimes several feet long are found (No. 4058). In Utah a few years ago a huge geode was discovered whose walls were covered with gigantic transparent crystals. Many of them now adorn the museums of this and other countries (Nos. 3890, 4308, 4498) (Plate XXXI). Multitudes of excellent crystals have been obtained at Girgenti, Sicily (No. 3567); Bex, Switzerland; Montmartre, France; Oxford, England (No. 3535). See Figures 229 to 233. Large transparent crystals from Kansas and Colorado, imperfect in outline, yield beautiful cleavage pieces. The cleavage parallel to the clinopinacoid (010) is so perfect that plates of any desired thickness may be obtained and used under a microscope to detect



the weak double refraction of some minerals. For example, if a plate of such thickness as to yield red of the first order between crossed nicols is used, it will become blue when the thin section under examination is positive (since gypsum is positive). In this case the color is raised. It becomes yellow, that is, depressed, when the mineral is negative.

There are two other cleavages also, one with a fibrous surface parallel to an orthodome (101), and one with conchoidal surface parallel to the orthopinacoid (100). Cleavage lines aid in orienting crystals that are without crystal faces.

Twinning is parallel to the orthopinacoid (100) both by juxtaposition (Fig. 232) and by interpenetration (Fig. 233). The fibrous cleavage cracks parallel to the orthodome do not run uniformly across the crystal, but meet at an angle of  $47^{\circ}50'$  on the line of contact.

The crystals are often curved or lenticular. (See Plate XXX.)

#### SUMMARY

*Gypsum*.— $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ;  $\text{CaO}=32.5$  per cent,  $\text{SO}_3=46.6$  per cent,  $\text{H}_2\text{O}=20.9$  per cent. Monoclinic;  $a:b:c=0.690:1:0.412$ .  $\beta=80^{\circ}42'$ . (010), (111), (110), (130), ( $\bar{1}03$ ). Twinned on (100), also on (101). Cleavage parallel (010) perfect; parallel (100) and (101) imperfect; parallel (100) and ( $10\bar{1}$ ) imperfect. Sectile; flexible.

Hardness=2; gravity=2.3. Colorless; luster vitreous; transparent;  $\beta=1.522$ . Double refraction positive;  $\gamma-\alpha=0.010$ . Axial plane parallel (010). Acute bisectrix inclined  $37^{\circ}30'$  to the normal of (100), and  $43^{\circ}12'$  to the normal of (001).  $2E=95^{\circ}$ .  $\rho > \nu$ . Inclined dispersion.

Easily fusible (3). Soluble in hydrochloric acid.

Michigan, New York, Virginia, Ohio, Iowa, Alabama, Arkansas, and the Cordilleran states.

## CLASS XII. TUNGSTATES, MOLYBDATES

The class of tungstates and molybdates contains but few minerals and those few are of slight importance. One example of each may be considered: the tungstate, wolframite; and the molybdate, wulfenite.

### Wolframite

Wolframite (No. 3533), a black mineral accompanying cassiterite in tin-bearing regions, and greatly resembling cassiterite in appearance, is the chief source of tungsten, an element being used in an increasing degree in manufactures. Well-formed monoclinic crystals resembling those shown in Figure 234 are common, but bladed, lamellar, or granular forms are more abundant. Its perfect cleavage parallel to (010) and its stibnite-like luster aid in distinguishing it, although otherwise it is a somewhat difficult mineral to identify, since blowpipe reactions for tungsten are masked by the presence of the iron and manganese in its formula,  $(\text{FeMn})\text{WO}_4$ . After wolframite is boiled in aqua regia, tungstic oxide appears as a yellow residue.




FIG. 234.—Wolframite

Tungsten steel is especially valuable for permanent magnets, cutting tools, wires for electric purposes, etc. Tungsten is also used as a dye which renders cotton less inflammable.

Wolframite is found in veins in Cornwall, Zinnwald, Black Hills, North Carolina, and Missouri.

### SUMMARY

*Wolframite*.— $(\text{FeMn})\text{WO}_4$ ; FeO varying from 2 to 19 per cent, MnO from 6 to 22 per cent,  $\text{WO}_3=76$  per cent. Monoclinic; holosymmetric. Cleavage parallel (010) perfect, parallel (100) imperfect. Brittle; fracture uneven.

Hardness=5.5; gravity=7.3. Black; streak reddish brown; luster metallic; opaque.

Fusible (3) to magnetic bead. Decomposed by hydrochloric acid. North Carolina, Missouri, South Dakota.

**Wulfenite**

This molybdate (Nos. 3528 and 3531),  $\text{PbMoO}_4$ , is a heavy, red, resinuous mineral which occurs in granular masses, and often in thin, tabular, square crystals (Fig. 235), or less commonly in acute pyramids (Fig. 236). Were it not so brittle and soft, it would be one of the most



FIG. 235.—Wulfenite




FIG. 236.—Wulfenite

prized of gems, since it is beautiful in color and has a high luster. Commercially it is of small value because of its rarity.

**SUMMARY**

*Wulfenite*.— $\text{PbMoO}_4$ ;  $\text{PbO}$ =60.7 per cent,  $\text{MoO}_3$ =39.3 per cent. Tetragonal; symmetry, tetragonal polar.  $a:c=1:1.577$ . (001), (102), (111), (320). Cleavage parallel (111) good. Brittle; fracture sub-conchoidal.

Hardness=3; gravity=6.7. Red; streak white; luster adamantine; translucent;  $\omega=2.402$ . Double refraction negative, strong;  $\omega-\epsilon=0.098$ .

Easily fusible (2); soluble in hydrochloric acid.

Arizona, New Mexico, California, Missouri, Pennsylvania.

### CLASS XIII. ORGANIC ACID SALTS

The organic acid salts, oxalates and mellates, which constitute Class XIII are rare and unimportant, hence we may pass at once to the next class.

### CLASS XIV. HYDROCARBONS

Though the members of this class are all of organic origin, yet they have been so changed by the loss of some constituent as to rank as mineral substances. Several of them are amorphous. Others retain the structure of the substance from which they were derived. Some of them may be most properly classified as rocks, but since they constitute part of a series they are here included. The most abundant representatives are the fossil resins, asphalt, heavy and light oils, gas, and coal.

#### Fossil Resins

Amber is a fossil resin occurring in amorphous masses which vary in size from small grains or droplets to chunks a foot or more in diameter. It was exuded from ancient conifers or leguminous trees, and buried by drifting sands in recent geological formations in Spain, Sicily, Germany, etc. It is characterized by conchoidal fracture, softness (hardness, 2), low specific gravity (gravity, 1.1), yellowish to brown color, greasy luster, and translucency. It shows fluorescence, is negatively electric, melts at about  $287^{\circ}$  and burns with a bright flame and an agreeable odor. It is composed of carbon, hydrogen, and oxygen ( $C_{10}H_8O_4$ ); C=78.93 per cent, H=10.55 per cent, O=10.52 per cent. It is used in the manufacture of buttons, beads, pipestems, varnish, amber oil, and acid.

Copal is a kind of amber which contains a larger proportion of hydrogen and melts at a lower temperature ( $210^{\circ}$ ). It is slightly harder than amber (hardness, 2.5). It is the dried sap of leguminous and coniferous trees which are found in many parts of the world, such as New Zealand, Australia, Madagascar, the east and west coasts of

Africa, and various places in South America. Nine-tenths of the copal used is obtained from deposits buried sometimes as deeply as twenty feet and often no doubt thousands of years old.

All varieties of amber are used chiefly as material from which to manufacture varnish. Insects imprisoned in the gum as it was exuding from trees have been preserved with remarkable fidelity, so that not only are their most delicate membranes intact, but in many cases an idea of the original color can be obtained.

### Asphalt

This hydrocarbon is of indefinite composition. It is black, burns with a pitchy odor and is slightly heavier than water (gravity, 1.1; hardness, 2). It melts at 100° C. and ignites readily with black smoke and bright flame. At a sufficiently low temperature it shows conchoidal fracture. It is soluble in ether.

It occurs in beds or lakes in the island of Trinidad and in veins or disseminated through sandstone or limestone in Kentucky, California, etc.

It forms one of the most satisfactory paving materials when mixed with sand and broken limestone. It is used for roofing, for calking material on ships, for paint on metal and woodwork, and as an adulterant of rubber goods.

### Petroleum

Petroleum is one of the most important of mineral substances, being second only to coal and iron in the contribution which it yearly makes to the wealth of mankind. It has been found in many countries, but nowhere so extensively as in the United States. It occurs in strata from the Ordovician to the Pleistocene. It has been produced by the distillation under great pressure of both animal and vegetable substances.

Petroleum, or "rock oil," is composed of a variety of oils, forming a series from the volatile and easily flowing oils to viscous oils, lubricating oils, and greases. It consists chiefly of the paraffines ( $C_nH_{2n+2}$ ) in Pennsylvania and of the naphthenes ( $C_6H_{12}$ ) in the Caucasus. The color varies from dark brown to greenish, and the gravity from 0.7 to 0.9. Petroleum shows a distinct fluorescence.

Benzine, naphtha, gasoline, kerosene, lubricating oil, vaseline, dye-stuffs, and other chemicals are derived from petroleum.

Pennsylvania long held chief place in the production of petroleum, but recently California has surpassed her, and Ohio, Indiana, Illinois, Kansas, Oklahoma, and Texas have all shown remarkable pools. Mexico and the Caucasus are increasing in productiveness.

#### Natural Gas

Closely associated with petroleum and having the same origin is natural gas. It too consists mainly of the lower paraffine methane ( $\text{CH}_4$ ) and ethane ( $\text{CH}_3$ ), and also carbon monoxides, carbon dioxide, hydrogen, neon, and the new gas helium, so necessary for war balloons.

#### Coal

Coal consists of solid hydrocarbons derived from vegetable growths of former geological ages. Trees, shrubs, weeds, mosses, and especially spores of cryptogams contributed to its formation. More than five hundred different species of plants have been identified among those concerned in the production of coal. Among them are six species of algae; two hundred and fifty species of ferns; eighty-three species of lycopods, that is, club mosses whose powder is used for fireworks, medicines, etc.; thirteen species of equisetites, that is, rushes, horsetails, etc.; sixty sigillarids, whose trunks were ribbed and scarred like giant cacti; twelve species of *noeggerathia* with overlapping scales on their trunks and pinnate leaves; forty-four *astero-phyllites*; and three species of cycads, the sago palms.

All of the above were acotyledons, the lowest form of vegetable life. They comprise five-sixths of all the plants which have been identified in coal. The remaining one-sixth were fifteen species of coniferous trees and fifteen of the palms. All of the above-named species are similar to vegetation which thrives in a warm, moist climate today. These plants, grown in swamps or near lakes and rivers, were deposited in beds, buried under mud that later turned to stone, and by the loss of hydrogen and oxygen were converted into coal.

Cross-sections of coal fields in all parts of the country point to such a history. At the bottom of a coal field occurs a conglomerate such as would form on a new shore line. This is covered by sandstone that indicates long action of the waves and gradual decrease in their severity. Next comes shale, and then clay, "fire clay," such as would be formed in the shallow waters of ponds into which sluggish streams

carry silt. These are followed by the coal from a few inches to several feet in thickness, such as might be formed in a peat swamp. The lowest coal bed in Illinois is called coal No. 1. This is covered by gray shale showing subsidence of the swamp and burial under mud. Further subsidence brought conditions favorable to formation of sandstone. Re-elevation introduced other shale and fire-clay formations which, in their turn, were followed by swamps in which coal No. 2 was formed. This shifting of the shore line was repeated many times in some localities, as is indicated not only by the different coal seams but also by their containing rocks. There are a dozen different beds in Illinois.

Mollusks, fishes, and amphibians buried in these deposits and changed to stone give further light upon the history of coal.

Coal is found in five geological systems, from the middle Tertiary down through the upper Cretaceous, the lower Jurassic (Oölite), and the Triassic to the Carboniferous. Of these systems the Carboniferous far surpasses all others in production.

In America there are seven extensive coal regions. The first is that included in Acadia, Nova Scotia, New Brunswick, and Rhode Island. The coal measures of Nova Scotia are 7,000 feet thick and contain 76 seams. In Rhode Island and Massachusetts a graphitic anthracite is found.

The second region, covering 70,000 square miles along the Appalachians, includes the famous coal fields of Pennsylvania, Ohio, Maryland, Virginia, West Virginia, Kentucky, Tennessee, and Alabama. In some portions of this field the coal measures are 4,000 feet thick. From no region of the world has more or better anthracite and bituminous coal been obtained.

The third field occupies about 7,000 square miles in Michigan, where the productive Carboniferous is but about 300 feet thick. Indiana and Illinois, with approximately 1,000 feet of Carboniferous strata covering 58,000 square miles, comprise the fourth field—one of the most actively worked and most remunerative.

At one time, continuous with this field was that which now constitutes the fifth field, covering 94,000 square miles. It is found in Iowa, Missouri, Arkansas, and Texas. Here the Carboniferous rocks are thicker than in any other portion of America but not for that reason more promising.

The sixth region is one of scattered character, occurring chiefly in Montana, Wyoming, Colorado, Utah, and Arizona. On the Pacific Coast is the seventh region, embracing Washington, Oregon, and California. Altogether there are more than 335,000 square miles of known coal-bearing territory in North America.

The anthracite area covers less than 1,000 square miles. Half of this is in Massachusetts and Rhode Island, where the anthracite is almost without fuel value because of its graphitic character and consequently no production has been reported in recent years. Colorado contains 15 square miles. Pennsylvania has a field covering 470 square miles. From this latter region practically all the anthracite produced in the United States is obtained. The total coal production in the United States in 1915 was valued at six hundred and eighty-six million dollars. More than a thousand million tons of coal are used in the world each year, and of this amount the United States furnishes the greatest part. Before the world-war Great Britain came next in production, followed in descending order by Germany, Austro-Hungary, France, Belgium, Russia, Japan, India, Canada, New South Wales, Spain, South African Republic, and New Zealand.

Coal was first used in London in 1240. After people had been using it for sixty-six years a law was passed against it on account of the smoke, which was declared to spoil ladies' complexions and clothes! As early as 1552 men began to fear all the coal in the world would soon be exhausted! In 1698 the first mention of coal in the United States was made by Father Hennepin as occurring near Fort Creve Coeur on the Illinois River near the place where Peoria now stands. Anthracite was discovered in Rhode Island in 1760. Being graphitic in character it was not used, and even the excellent variety which occurs in Pennsylvania lay unutilized for forty years after its presence was known. All early use of coal was very local owing to lack of transportation, but with the advent of coal, transportation and the growth of cities became a possibility.

As society is now constituted, no mineral substance could be spared with greater difficulty, and, in fact, without coal modern civilization would be impossible. Railroads, steamships, and great manufacturing plants would disappear. Men would miserably perish in winter's cold or all be driven to the tropics.



TABLE OF VARIOUS COALS

Name	Physical Condition	Hard- ness	Grav- ity	Color	Remarks
Peat.....	Feltlike aggregates	1.5	1.05	Light brown	Heat value low. Found in swamps in northern countries.
Lignite.....	Earthy; slightly conchoidal fracture	1.5	1.2	Black; powder brown	Sooty flame; does not fuse nor coke in closed vessel; partly soluble in warm potassium hydrate yielding brown solution. In younger stratified rocks. Source of gasoline, benzene, petroleum, paraffine, carbolic acid, vinegar, etc. Chief fuel in portions of western United States.
Bituminous coal, classified as follows:	Compact; conchoidal fracture	2	1.4	Black; powder brown	Long yellow flame; some varieties fuse; others coke; soluble in potassium hydrate without leaving brown solution; vegetable structure shown after bleaching in nitric acid, potassium chlorate, and washing in alcohol. In older geological formations (Pennsylvanian). Underlying many hundred square miles of country in the United States. Chief fuel, source of gas, tar, oils, and other derivations.
1. Caking coal (fat).....	Softens while burning				
2. Coking coal (lean).....	Retains shape while burning				
3. Cherry coal.....	Bright, lustrous				
4. Splint coal.....	Slaty and heavy				
5. Pitch coal.....	Very black, coarse conchoidal fracture				
6. Glance coal.....	Satiny, brittle				
7. Cannel coal (candle).....	Fine grained, bright flame, sufficiently tough to be used for boxes, buttons, chandeliers, etc.				
Anthracite.....	Compact; brittle; conchoidal fracture; absence of cleavage	2.5	1.7	Black to grayish	Difficult to ignite, burns with weak flame, does not fuse. Soluble in potassium hydrate without brown solution. Plant cells may be seen in ashes on microscopic slides. In limited areas in the oldest sedimentary rocks and crystalline schists.

One pound of coal in a good engine will produce six-horse-power for one hour. One ton will produce thirteen-thousand-horse-power; and since some railroads use ten thousand tons per day, they have the equivalent of the work of one hundred and thirty million horses for one hour—without the necessity of feeding the horses. It is estimated that one pound of coal can produce as many foot pounds of energy as one man in one day. Three hundred pounds will furnish as much power as one man per year. Then if half the coal produced in the United States in 1915 were used as a source of power, it could do the work of sixteen hundred million men. This furnishes one explanation of the remarkable growth in wealth of the United States in the last fifty years—a growth which has not been equaled before in the history of the world. In using coal so lavishly we are drawing on the energy stored in the earth by the slow growth and transformation of a succession of swamps and forests requiring the sunshine of millions of years. The disappearance of the coal supply is but a question of time. In less than 300 years workable coal seams will probably be exhausted in Europe. Those in other parts of the world will last longer. But coal producers and users should seek to avoid the wasteful methods which at present prevail in: (1) mining, (2) removal, and (3) in use in furnace, stove, and fireplace.

The following table shows the typical proportions of carbon, hydrogen, oxygen, and nitrogen in the transformation of wood to anthracite.

CHEMICAL CHANGES IN TRANSFORMATION OF WOOD TO COAL

	Carbon	Hydrogen	Oxygen and Nitrogen
Wood.....	50	6	43
Peat.....	59	6	33
Lignite.....	69	5.2	25
Bituminous coal.....	82	5	12.2
Anthracite coal.....	95	2.5	2.5

## SUMMARY

Having proceeded thus far, the visitor to the museum has made the acquaintance of about 100 different minerals. Many more are worthy of his interest and attention; yet this number is sufficient to give an idea of the minerals which constitute the world, and which are used by man for ornament, for medicine, as the source of metal, for building material, and in many other ways.

Such a study furnishes results similar to those which would be obtained by a student of human society who went into a community of some 1,200 inhabitants and made the acquaintance of 100 different people engaged in different occupations, holding different responsibilities, and showing varied attainments. One who has finished such a study would have a fair idea of the whole community.

So one who has passed through the museum, noting carefully the minerals shown and giving attention to their physical and chemical laws, their geography, geology, and relation to human activities, has a good idea of the whole mineral world. It is not necessary for him to study all the thirteen hundred different species and varieties of minerals. However, for one who wishes to go farther, a comprehensive list of minerals is given on pages 202 to 275; and for further study he is referred to the books listed on page 200.

No country is better supplied with minerals than the United States, and few countries make as good use of their resources in this regard as we do. The world-war resulted in stimulating mineral production in this country. For a number of minerals we had been accustomed to go to foreign countries; for antimony we had gone to China, for chromium to New Caledonia, for graphite to Ceylon, for magnesite to Greece, for manganese and platinum to Russia, for sulphur to Sicily, for tin to Singapore, for vanadium to Peru. But with increased difficulty of ocean transportation, prospectors and producers became increasingly active in the search for and the mining of these minerals. So the time is near at hand when the United States may be nearly independent in regard to the minerals necessary for the activities of its people.

In 1915 the total wealth added to the country by our minerals was two billion three hundred and ninety-three million dollars.

New minerals and chemical substances are being constantly discovered, and with their discovery new ideas and inspiration is gained by advanced workers in various departments of science. Most prominent among recent advances are those which have been made by men studying the ultimate constitution of matter.

## NAMES OF MINERALS

Among the minerals which we have seen, the name of one at least, kaolinite, is of Chinese origin; two are of Singhalese origin: corundum and tourmaline; three are of Arabic origin: marcasite, amber, and talc. Bismuth, zincite, and hornblende are taken directly from the German; while gold, silver, and iron are old Anglo-Saxon words.

Many minerals are named after some geographical locality, such as aragonite, anglesite, labradorite, muscovite, strontianite, tremolite.

Others are named after men distinguished in the science of mineralogy or otherwise—biotite, dolomite, goethite, franklinite, magnesite, magnetite, proustite, smithsonite, tennantite, witherite.

A still larger number were derived from the Latin language: sulphur, antimony, platinum, mercury, stibnite, argentite, erubescite, tetrahedrite, sylvite, fluorite, cassiterite, rutile, spinel, cerussite, manganite, albite, enstatite, actinolite, garnet, celestite, asphalt.

And finally, a still larger number of mineral names originate from the Greek: diamond, graphite, copper, molybdenite, galena, chalcocite, sphalerite, cinnabar, pyrrhotite, chalcopyrite, pyrite, arsenopyrite, pyrrargyrite, halite, cryolite, chalcedony, cuprite, hematite, chromite, pyrolusite, limonite, calcite, siderite, rhodochrosite, malachite, azurite, orthoclase, microcline, oligoclase, anorthite, hypersthene, pyroxene, diopside, augite, rhodonite, barite, gypsum.

## THE USES OF MINERALS

Minerals contribute toward the welfare of mankind in manifold ways. Many of the harder, more brightly colored, or highly refracting minerals since earliest times have been used as objects of personal adornment, and today among the most prized of all material objects are such minerals as diamonds, rubies, sapphires, emeralds, aquamarine, amethyst, agates, turquoise, tourmaline, olivene, rhodonite, and malachite.

The metals, together with their sulphides, oxides, carbonates, and silicates, play a weighty rôle in the life of men of all races and all stages of development. The condition of society would be materially different were there no gold, silver, platinum, copper, iron, tin, zinc, lead, paladium, chromium, aluminium, manganese, magnesium, mercury, antimony, or bismuth.

Some minerals form foods without which it would be well-nigh impossible for men to exist. For example, salt and the minerals which are the source of the alkalies are almost indispensable to life.

The number of minerals which are used in the arts and manufactures is large and important. Sulphur, phosphorus, soda, potash, chlorine, fluorine, and calcium contribute largely to the wealth of men.

Attention has already been called to the indispensable character of the hydrocarbon compounds. Without them modern civilization would be an impossibility.

Minerals as rock constituents form mountains and plains, and by their decomposition furnish the ultimate food supply of mankind. The study of minerals in their capacity of soil-formers is one of surpassing interest.

## HISTORY OF THE STUDY OF MINERALS

The science of mineralogy, depending as it does upon physics, chemistry, and other well-developed sciences, has been one of the latest to be pursued, although from very early times minerals were used for ornaments, weapons, and domestic utensils. While ancient literature abounds in references to minerals, little more was known about them in early times than their external form. Hebrew literature mentions the use of clay, niter, salt, sand, and sulphur, as well as of gold, silver, copper, emerald, agate, chalcedony, carnelian, jasper, onyx, sardonyx, topaz, ruby, and sapphire.

Aristotle, 322 B.C., who is reputed by his admirers "to have known something of every science," has given no evidence of acquaintance with minerals. Several references to the subject in his writings are thought to have been interpolations made later by other writers. Pliny, 79 A.D., was the first Latin writer to describe minerals, and his accounts are usually so incomplete as to leave doubt as to the minerals to which they apply! Avicenna (d. 1036), an Arabian doctor of medicine born near Bokhara, distinguished salts, metals, minerals, and stones. Agricola (d. 1555), a German doctor born in Joachimsthal, used the terms quartz and spar, and noted the hardness, cleavage, form, and luster of certain minerals. This is a short list to cover all the years to the seventeenth century. But during the latter part of the seventeenth century a number of men began to be interested in the subject, Robert Boyle (1691) investigating their chemistry, Niccolus Steno discovering the constancy of crystal angles, and Bartholinus noting the double refraction of calcite. During the eighteenth century Linnaeus, the great classifier, attempted to classify minerals according to their form, while Cronstedt attempted a chemical classification. Two Frenchmen, Romé de l'Isle and René Just Haüy, were enthusiastic investigators in crystallography. De l'Isle described and pictured many forms. Haüy discovered laws underlying them. Jealousy of each other's work made them enemies. Haüy enjoyed recording de l'Isle's errors while correcting them. But their work formed the basis of later work in crystallography.

During the nineteenth century the study of minerals was pursued by many workers and the advancement in many lines assumed admirable proportions. In Germany, Weiss developed the idea of systems of crystallization, Mohs investigated the natural history of minerals, and Werner studied chemical classification and developed determination of minerals by simple physical characteristics. In Germany, France, England, and America the number of workers increased, some pursuing the subject of crystal formation, as did Bravais, Sohncke, Naumann, Miller, and Liebisch; others working at the chemical side of the subject, as, for example, Berzelius, Rose, Bunsen, Mitscherlich, Plattner, and Rammelsberg; still others studying optical mineralogy, noting particularly the relation of form to physical properties—Brewster, Senarmont, Des Cloizeaux, Zirkel, Sorby, Wollaston. The systematic side of the subject was developed by Beudant, Breithaupt, Groth, and Dana. Fouqué, Michel Levy, and Daubrée gave attention to the artificial formation of minerals.

The increase in interest in the subject of mineral study has come largely from advance in mining and in the use of minerals in arts and manufactures. The study has been and is naturally one of materials rather than of laws; but as the science has progressed, the principles and laws have been gradually perceived and formulated. In the development of the science, contributions have been made to other sciences: to physics, knowledge of optical and electrical phenomena; to chemistry, knowledge of new substances; to geology, light on the origin, composition, and decay of rocks; to metallography, facts concerning the contents and treatment of ores. Within the last fifty years the number of workers in mineralogy has increased to such an extent that the list is an extensive one and the science has been brought to great perfection in various lines.

In the United States many excellent books on the subject have been written. No work has ever surpassed in completeness that of James Dwight Dana. His *System of Mineralogy*, which first appeared in 1837, has passed through six editions. After the elder Dana's death the *Manual of Mineralogy*, which first appeared in 1848 and has passed through thirteen editions, and the *Textbook of Mineralogy*, which first appeared in 1877, were rewritten, enlarged, and kept up to date, first by Edward Salisbury Dana and later by William E. Ford. Among many books which have appeared during the last



dozen years, the following may be noted especially. In them there has been a general endeavor to present this rather difficult science in such a manner as to render it more attractive to the general student. Increasing use is made of diagrams, of models, and of photographs of minerals. Anyone wishing to pursue the subject further should consult the following excellent works:

- Bayley, W. S., *Descriptive Mineralogy*. D. Appleton & Co., 1917.  
 Brush, G. J., and Penfield, S. L., *Determinative Mineralogy and Blowpipe Analysis*. John Wiley, 1907.  
 Butler, G. M., *Handbook of Minerals*. John Wiley, 1908.  
 Erni, H., and Brown, A. P., *Mineralogy Simplified*. Philadelphia, 1901.  
 Farrington, O. C., *Meteorites*. Lakeside Press, 1915.  
 ———, *Gems and Gem Minerals*. Lakeside Press, 1902.  
 Fulton, A. E. H., *Crystallography and Practical Crystal Measurement*. Macmillan Co., 1911.  
 Gratacap, L. P., *Popular Guide to Minerals*. D. Van Nostrand Co., 1912.  
 Groth, P., and Jackson, B. H., *Optical Properties of Crystals*. John Wiley, 1910.  
 Iddings, J. P., *Rock Minerals*. John Wiley, 1906.  
 Johannsen, Albert, *Determination of Rock-Forming Minerals*. John Wiley, 1908.  
 Kraus, E. H., *Descriptive Mineralogy*. George Wahr, 1911.  
 Kunz, G. F., *Gems and Precious Stones of North America*. New York.  
 Lewis, J. V., *Determinative Mineralogy*. John Wiley, 1915.  
 Merrill, G. P., *Rocks, Rock Weathering, and Soils*. John Wiley, 1906.  
 ———, *Non-Metallic Minerals*. Macmillan Co., 1904.  
 Moses, A. J., and Parsons, C. L., *Mineralogy, Crystallography, and Blowpipe Analysis*. D. Van Nostrand Co., 1909.  
 Phillips, A. H., *Mineralogy*. Macmillan Co., 1912.  
 Pirsson, L. V., *Rocks and Rock Minerals*. John Wiley, 1908.  
 Van Horn, F. R., *Lecture Notes on Mineralogy*. Cleveland: 1903.  
 Winchell, N. H. and A. N., *Elements of Optical Mineralogy*. D. Van Nostrand Co., 1909.

An attractive little monthly magazine, *The American Mineralogist*, edited by Edgar T. Wherry, to be obtained of H. W. Trudell, Philadelphia, describes new minerals and records events of interest to mineralogists. *Economic Geology*, *Mineral Industry*, *American Journal of Science*, *Science*, and other journals contain many articles on mineralogy. Various state geological reports and those of the United States Geological Survey are full of interesting and valuable information concerning occurrence and production of minerals.

## **COMPREHENSIVE LIST OF MINERALS**

## COMPREHENSIVE

	Composition	Form
<b>I. ELEMENTS</b>		
1. Diamond.....	C	Regular
2. Bort.....	C	Regular
3. Carbonado.....	C	Massive
4. Cliftonite.....	C	Massive
5. Graphite.....	C	Hexag.
6. Schungite.....	C	Amorph.
7. Sulphur.....	S	Ortho.
8. Selen sulphur.....	SeS	Ortho.
9. Arsenic.....	As	Hexag.
10. Allemonite.....	SbAs <sub>3</sub>	Hexag.
11. Tellurium.....	Te	Hexag.
12. Antimony.....	Sb	Hexag.
13. Bismuth.....	Bi	Hexag.
14. Zinc.....	Zn	Hexag.
15. Gold.....	Au	Regular
16. Electrum.....	Au·Ag	Regular
17. Silver.....	Ag	Regular
18. Copper.....	Cu	Regular
19. Mercury.....	Hg	Amorph.
20. Lead.....	Pb	Regular
21. Amalgam.....	(Ag,Hg)	Regular
22. Arquerite.....	(Ag <sub>11</sub> Hg)	Regular
23. Kongsbergite.....	(Ag <sub>13</sub> Hg)	Regular
24. Tin.....	Sn	Tetrag.
25. Platinum.....	Pt	Regular
26. Iridium.....	Ir	Regular
27. Iridosmine.....	IrOs	Hexag.
28. Nevyanskite.....	IrOs	Hexag.
29. Siserskite.....	IrOs	Hexag.
30. Palladium.....	Pd	Regular
31. Allopalladium.....	Pd	Hexag.
32. Iron.....	Fe	Regular
33. Awaruite.....	FeNi <sub>2</sub>	Regular
34. Josephinite.....	Fe <sub>2</sub> Ni <sub>3</sub>	Regular
35. Meteoric Iron.....	Fe	Regular
36. Kamacite.....	Fe <sub>14</sub> Ni	Regular
37. Taenite.....	Fe <sub>9</sub> Ni <sub>10</sub>	Regular
38. Plessite.....	Fe <sub>9</sub> Ni <sub>10</sub>	Regular
39. Cohenite.....	(Fe,Ni,Co) <sub>3</sub> C	Regular
39a. Schreibersite.....	(Fe,Ni,Co) <sub>3</sub> P	Tetrag.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
1.	Colorless	10	3.5	Kimberley	Gem
2.	Dark	10	3.5	Kimberley	} Drills
3.	Black	10	3.5	Kimberley	
4.	Black	2.5	2.1	Meteorites	Carbon
5.	Black	1	2	Ceylon	Pencils
6.	Black	1	1.9	Russia	Carbon
7.	Yellow	2	2	Sicily	Drugs
8.	Reddish	.....	.....	Sicily	Selenium
9.	Tin white	3.5	5.6	Freiberg	} Drugs
10.	Tin white	3.5	6.2	Andreasberg	
11.	Tin white	2	6.1	Colorado	} Alloys
12.	White	3	6.6	Japan	
13.	Reddish	2.5	9.8	Western U.S.	Zinc
14.	White	2	6.9	Australia	Coin
15.	Yellow	2.5	19	Western U.S.	Gold
16.	Amber	2.5	15	Urals	Coin
17.	White	2.5	11	Western U.S.	Wire
18.	Red	2.5	8.9	Michigan	Amalgamation
19.	White	.....	13.5	California	Lead
20.	Gray	1.5	11.3	Colorado	} Silver
21.	White	3	14	Sweden, S. America	
22.	White	3	10	Sweden, S. America	} Tin
23.	White	3	14	Sweden, S. America	
24.	White	2	7	Siberia, N.S. Wales	Dentistry
25.	White	5	21	Urals	Pen points
26.	Regular	6	22	Urals	} Iridium and osmium
27.	White	6	21	Urals, S. America	
28.	White	6	19	Urals, S. America	} Palladium
29.	White	6	21	Urals, S. America	
30.	Steel gray	4	11	Brazil	} Iron and nickel
31.	Steel gray	.....	.....	Hartz	
32.	Iron black	4	7	Meteorites	}
33.	Gray	5	8	New Zealand	
34.	Gray	5	8	Oregon	}
35.	Gray	4	7	Meteorites	
36.	Gray	4	7	Meteorites	}
37.	Gray	4	7	Meteorites	
38.	Gray	4	7	Meteorites	}
39.	Tin white	6	7	Meteorites	
39a.	Tin white	6	7	Meteorites	

	Composition	Form
<b>II. SULPHIDES</b>		
<b>1. Sulphides of Semi-Metals</b>		
40. Realgar .....	AsS	Mono.
41. Orpiment .....	As <sub>2</sub> S <sub>3</sub>	Mono.
42. Stibnite .....	Sb <sub>2</sub> S <sub>3</sub>	Ortho.
43. Metastibnite .....	Sb <sub>2</sub> S <sub>3</sub>	Amorph.
44. Bismuthinite .....	Bi <sub>2</sub> S <sub>3</sub>	Ortho.
45. Guanajuatite .....	Bi <sub>2</sub> Se <sub>3</sub>	Ortho.
46. Tetradymite .....	Bi <sub>2</sub> (Te,S) <sub>3</sub>	Hexag.
47. Joseite .....	Bi <sub>2</sub> Te <sub>2</sub> Se	Hexag.
48. Wehrlite .....	Bi <sub>2</sub> Te <sub>2</sub>	Hexag.
49. Molybdenite .....	MoS <sub>2</sub>	Hexag.
<b>2. Sulphides of Metals</b>		
<b>a. Basic</b>		
50. Dyscrasite .....	Ag <sub>3</sub> Sb · Ag <sub>3</sub> Sb	Ortho.
51. Horsfordite .....	Cu <sub>6</sub> Sb	Ortho.
52. Hunttilite .....	Ag <sub>3</sub> As	Ortho.
53. Animikite .....	Ag <sub>3</sub> Sb	Ortho.
54. Domeykite .....	Cu <sub>3</sub> As	Ortho.
55. Algodonite .....	Cu <sub>4</sub> As	Ortho.
56. Whitneyite .....	Cu <sub>3</sub> As	Ortho.
57. Chilenite .....	Ag <sub>6</sub> Bi	Amorph.
58. Stützite .....	Ag <sub>4</sub> Te	Hexag.
<b>b. Monosulphides</b>		
59. Galena .....	PbS	Regular
60. Cuproplumbite .....	Cu <sub>2</sub> S · 2PbS	Massive
61. Alisonite .....	3Cu <sub>2</sub> S · PbS	Massive
62. Altaite .....	AgTe	Regular
63. Clausthalite .....	PbSe	Regular
64. Tilkerodite .....	(PbCo)Se	Regular
65. Naumannite .....	(Ag,Pb)Se	Regular
66. Argentite .....	Ag <sub>2</sub> S	Regular
67. Jalpaite .....	(Ag,Cu) <sub>2</sub> S	Regular
68. Hessite .....	Ag <sub>2</sub> Te	Regular
69. Petzite .....	(Ag,Au) <sub>2</sub> Te	Massive
70. Aguilarite .....	Ag <sub>2</sub> S · Ag <sub>2</sub> Se	Regular
71. Berzelianite .....	Cu <sub>2</sub> Se	Dend.
72. Lehrbachite .....	PbSe · HgSe	Massive
73. Eucairite .....	Cu <sub>2</sub> Se · Ag <sub>2</sub> Se	Regular
74. Zorgite .....	(PbCu <sub>2</sub> Ag <sub>2</sub> )Se	Massive
75. Crookesite .....	(CuTlAg) <sub>2</sub> Se	Massive
76. Umangite .....	CuSe · Cu <sub>2</sub> Se	Massive
77. Chalcocite .....	Cu <sub>2</sub> S	Ortho.
78. Stromeyerite .....	(Ag,Cu) <sub>2</sub> S	Ortho.
79. Sternbergite .....	AgFeS <sub>3</sub>	Ortho.
80. Friesite .....	Ag <sub>2</sub> Fe <sub>2</sub> S <sub>3</sub>	Ortho.
81. Acanthite .....	Ag <sub>2</sub> S	Ortho.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
40.	Aurora red	1.5	3.5	California	} Arsenic
41.	Yellow	1.5	3.5	Utah, Wyoming	
42.	Lead gray	2	5	Japan	} Antimony
43.	Brick red	.....	.....	Nevada	
44.	White, iridescent	2	6	England, N.C.	} Bismuth
45.	Bluish gray	2	6	Mexico	
46.	Steel gray	1.5	7	North Carolina	} Bismuth
47.	Steel gray	2	7.9	Brazil	
48.	Steel gray	2	8.4	Hungary	} Molybdenum
49.	Lead gray	1.5	4.6	Washington	
50.	Tin white	3.5	9.5	Hartz Mts.	Silver
51.	Tin white	.....	8.8	Mytilene	Copper
52.	Tin white	.....	7	Lake Superior	} Silver
53.	White	.....	9	Lake Superior	
54.	Tin white	.....	7.5	Lake Superior	} Copper
55.	Tin white	.....	7.6	Lake Superior	
56.	Reddish white	.....	8.4	Houghton, Mich.	} Silver
57.	Silver white	2	.....	Chile	
58.	Lead gray	.....	.....	Nagyag	
59.	Lead gray	2.5	7	Missouri, Colorado	Lead
60.	Dark blue	.....	6	Chile	} Copper
61.	Indigo blue	.....	6	Chile	
62.	Tin white	3	8.1	Chile, Colorado	Lead
63.	Lead gray	3	8	Hartz Mts.	} Lead
64.	Lead gray	.....	8	Hartz Mts.	
65.	Iron black	2.5	8	Hartz Mts.	} Silver
66.	Lead gray	2.5	7	Western U.S.	
67.	Lead gray	.....	6	Mexico	} Gold
68.	Lead gray	2.5	8	Boulder, Colo.	
69.	Iron black	2.5	9	California	Silver
70.	Iron black	2.5	7.5	Mexico	Copper
71.	Silver white	2	6.7	Sweden	Copper
72.	Iron black	.....	7.8	Hartz Mts.	Mercury
73.	Lead gray	.....	7.5	Chile	} Copper
74.	Lead gray	2	7	Hartz Mts.	
75.	Lead gray	3	6.9	Sweden	Silver
76.	Cherry red	3	5.6	Argentina	Copper
77.	Lead gray	2	5	Montana	Copper
78.	Steel gray	2.5	6	Siberia, Colorado	} Silver
79.	Pinchbeck brown	1	4.2	Saxony	
80.	Dark gray	2.5	4	Joachimsthal	
81.	Iron black	2.5	7	Joachimsthal	

	Composition	Form
<b>II. SULPHIDES—continued</b>		
82. Sphalerite.....	ZnS	Regular
83. Marmatite.....	ZnS·FeS	Regular
84. Przibramite.....	ZnS·CdS	Regular
85. Metacinnabarite.....	HgS	Regular
86. Guadalcazarite.....	HgS·ZnS	Regular
87. Tiemannite.....	HgSe	Regular
88. Onofrite.....	Hg(S·Se)	Regular
89. Coloradoite.....	HgTe	Regular
90. Alabandite.....	MnS	Regular
91. Oldhamite.....	CaS	Regular
92. Pentlandite.....	(Fe,Ni)S	Regular
93. Troilite.....	FeS	Hexag.
94. Cinnabar.....	HgS	Hexag.
95. Covellite.....	CuS	Hexag.
96. Greenockite.....	CdS	Hexag.
97. Wurtzite.....	ZnS	Hexag.
98. Erythrozincite.....	ZnS·MnS	
99. Millerite.....	NiS	Hexag.
100. Beyrichite.....	NiS	Hexag.
101. Hauchecornite.....	(Ni,Co) <sub>7</sub> (S,Sb,Bi) <sub>8</sub>	Tetrag.
102. Niccolite.....	NiAs	Hexag.
103. Breithauptite.....	NiSb	Hexag.
104. Pyrrhotite.....	Fe <sub>n</sub> S <sub>12</sub>	Hexag.
<b>c. Intermediate</b>		
105. Horbachite.....	4Fe <sub>2</sub> S <sub>3</sub> Ni <sub>2</sub> S <sub>3</sub>	Hexag.
106. Polydymite.....	Ni <sub>4</sub> S <sub>5</sub>	Regular
107. Grünauite.....	Ni <sub>4</sub> S <sub>5</sub> ·Bi <sub>2</sub> S <sub>3</sub>	Regular
108. Sychnodymite.....	(Co,Cu) <sub>4</sub> S <sub>5</sub>	Regular
109. Melonite.....	Ni <sub>4</sub> Te <sub>3</sub>	Hexag.
110. Bornite (= Erubescite)	3Cu <sub>2</sub> S·Fe <sub>2</sub> S <sub>3</sub>	Regular
111. Linnaeite.....	(Ni,Co,Fe) <sub>3</sub> S <sub>4</sub>	Regular
112. Daubreehlite.....	FeS·Cr <sub>2</sub> S <sub>3</sub>	
113. Cubanite.....	CuFe <sub>2</sub> S <sub>4</sub>	Regular
114. Carrollite.....	CuCo <sub>2</sub> S <sub>4</sub>	Regular
115. Chalcopyrite.....	CuFeS <sub>2</sub>	Regular
<b>d. Disulphides</b>		
116. Pyrite.....	FeS <sub>2</sub>	Regular
117. Hauerite.....	MnS <sub>2</sub>	Regular
118. Smaltite-chloanthite	CoAs <sub>2</sub> ·NiAs <sub>2</sub>	Regular
119. Cobaltite.....	CoAsS	Regular
120. Gersdorffite.....	NiAsS	Regular
121. Corynite.....	Ni(As,Sb)S	Regular
122. Willyamite.....	CoS <sub>2</sub> ·NiS <sub>2</sub> ·CoSb <sub>2</sub> ·NiSb <sub>2</sub>	Regular
123. Ullmannite.....	NiSbS	Regular
124. Kallilite.....	Ni(Sb,Bi)S	Massive
125. Sperrylite.....	PtAs <sub>2</sub>	Regular
126. Laurite.....	RuS <sub>2</sub>	Regular

## LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
82.	Yellow	3.5	3.9	Missouri	} Zinc
83.	Dark brown	.....	3.9	Cornwall	
84.	Dark brown	3.5	3.9	Hungary	
85.	Grayish black	3	7	California	} Mercury
86.	.....	2	7	Gaudalcazar, Mex.	
87.	Steel gray	2.5	8	Hartz Mts.	
88.	Blackish gray	2.5	8	Mexico	} Manganese
89.	Iron black	3	8	Colorado	
90.	Iron black	3.5	3.9	Colorado	
91.	Pale brown	4	2.5	S.C. Meteorites	Calcium
92.	Bronze yellow	3.5	4.6	Norway	Nickel
93.	Tombac brown	4.7	4.7	Meteorites	Iron
94.	Reddish brown	2	8	Spain, California	Mercury
95.	Indigo blue	1.5	4.5	Chile	Copper
96.	Yellow	3.5	4.9	Scotland	Cadmium
97.	Brownish black	3.5	3.9	Peru	} Zinc
98.	Red	2	3.9	Siberia	
99.	Brass yellow	3.5	5	Saxony	
100.	Lead gray	3	4.7	Westerwald	} Nickel
101.	Bronze yellow	5	6	Westphalia	
102.	Copper red	5	7	Sweden	
103.	Red	5	7.5	Andreasberg	} Sulphur
104.	Bronze	3.5	4.5	Pennsylvania	
105.	Steel gray	4.5	4	Horback	} Nickel
106.	Gray	4.5	4.5	Grünau	
107.	Steel gray	4.5	5	Grünau	
108.	Steel gray	.....	4.7	Siegen	Cobalt
109.	Reddish white	.....	.....	California	Nickel
110.	Copper	3	5	Chile	Copper
111.	Steel gray	5.5	4.8	Sweden	Cobalt
112.	Black	.....	5	Meteoric iron	Chromium
113.	Bronze	4	4	Cuba	} Copper
114.	Steel gray	5.5	4.8	Maryland	
115.	Yellow	3.5	4	Western U.S.	Copper
116.	Yellow	6	5	Everywhere	Sulphur
117.	Brown	4	3	Hungary	Manganese
118.	Tin white	5.5	6	Saxony	} Cobalt
119.	Silver white	5.5	6	Sweden	
120.	Silver white	5.5	6	Sweden	
121.	Silver white	4.5	5.9	Olso	} Nickel
122.	Silver white	5	7	New South Wales	
123.	Steel gray	5.5	6	Germany	} Nickel
124.	Bluish gray	.....	.....	Germany	
125.	Tin white	6	10	Canada	Platinum
126.	Iron black	7.5	6.9	Borneo	Ruthenium



## COMPREHENSIVE

	Composition	Form
<b>II. SULPHIDES—continued</b>		
127. Skutterudite.....	$\text{CoAs}_3$	Regular
128. Nickel-skutterudite...	$(\text{Ni}, \text{Co}, \text{Fe})\text{As}_3$	Massive
129. Bismuto-smaltite.....	$\text{Co}(\text{As}, \text{Bi})_3$	Massive
130. Marcasite.....	$\text{FeS}_2$	Ortho.
131. Löllingite.....	$\text{FeAs}_2$	Ortho.
132. Leucopyrite.....	$\text{Fe}_2\text{As}_4$	Ortho.
133. Geyerite.....	$\text{Fe}(\text{AsS})_2$	Ortho.
134. Arsenopyrite.....	$\text{FeAsS}$	Ortho.
135. Danaite.....	$\text{FeCoAsS}$	Ortho.
136. Safflorite.....	$\text{CoAs}_2$	Ortho.
137. Rammelsbergite.....	$\text{NiAs}_2$	Ortho.
138. Glaucodot.....	$(\text{Co}, \text{Fe})\text{AsS}$	Ortho.
139. Alloclasite.....	$\text{Co}(\text{As}, \text{Bi})\text{S}$	Ortho.
140. Wolfachite.....	$\text{Ni}(\text{As} \cdot \text{Sb})\text{S}$	Ortho.
140a. Maucherite.....	$\text{Ni}_3\text{As}_2$	Tetrag.
<b>e. Tellurides</b>		
141. Sylvanite.....	$(\text{Au}, \text{Ag})\text{Te}_2$	Mono.
142. Krennerite.....	$(\text{Au}, \text{Ag})\text{Te}_2$	Ortho.
143. Calaverite.....	$(\text{Au}, \text{Ag})\text{Te}_2$	Mono.
144. Nagyagite.....	$\text{Au}_2\text{Pb}_{12}\text{Sb}_2(\text{S}, \text{Te})_{24}$	Ortho.
<b>f. Oxysulphides</b>		
145. Kermesite.....	$\text{Sb}_2\text{S}_2\text{O}$	Mono.
146. Voltzite.....	$\text{Zn}_3\text{S}_4\text{O}$	Globules

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
127.	Tin white	6	6.7	Norway	Cobalt
128.	Gray	.....	.....	New Mexico	
129.	Tin white	.....	6.9	Zschorlau	Sulphur
130.	Yellow	6.5	4.8	Bohemia	
131.	Silver white	5.5	7	Lolling-Huttenberg	Arsenic
132.	Silver white	5.5	7	Lolling-Huttenberg	
133.	.....	.....	6.8	Saxony	Arsenic
134.	Silver white	5.5	6	Freiberg	
135.	Gray	5.5	6	Franconia	Cobalt
136.	Tin white	4.5	7	Saxony	
137.	Tin white	5.5	7	Saxony	Nickel
138.	Tin white	5	5.9	Chile	
139.	Steel gray	4.5	6.6	Orawitza	Cobalt
140.	Silver white	4.5	6	Wolfach	
140½.	Reddish	5	7	Thuringen	Nickel
141.	Steel gray	1.5	7.9	Nagyag	
142.	Silver white	.....	8	Nagyag, Colo.	Gold
143.	Yellow	2.5	9	California, Colo.	
144.	Lead gray	1	6.3	Nagyag	Antimony
145.	Red	1	4.5	Hungary	
146.	Red	4	3.6	Joachimsthal	Zinc

	Composition	Form
<b>III. SULPHO-SALTS</b>		
<b>I. Sulpharsenites, etc.</b>		
<b>a. Acidic</b>		
147. Livingstonite.....	$\text{HgS} \cdot 2\text{Sb}_2\text{S}_3$	Ortho.
148. Chiviatite.....	$2\text{PbS} \cdot 3\text{Bi}_2\text{S}_3$	Ortho.
149. Cuprobismutite.....	$3\text{Cu}_2\text{S} \cdot 4\text{Bi}_2\text{S}_3$	Ortho.
150. Rezbanyite.....	$4\text{PbS} \cdot 5\text{Bi}_2\text{S}_3$	Ortho.
<b>b. Meta</b>		
151. Zinkenite.....	$\text{PbSb}_2\text{S}_4$	Ortho.
152. Andorite.....	$2(\text{Pb}, \text{Ag}, \text{Sb})_2\text{S}_6$	Ortho.
153. Webnerite.....		
154. Sundtite.....		
155. Sartorite.....	$\text{PbS} \cdot \text{As}_2\text{S}_3$	Ortho.
156. Emplectite.....	$\text{Cu}_2\text{S} \cdot \text{Bi}_2\text{S}_3$	Ortho.
157. Chalcostibite.....	$\text{Cu}_2\text{S} \cdot \text{Sb}_2\text{S}_3$	Ortho.
158. Galenobismutite.....	$\text{PbS} \cdot \text{Bi}_2\text{S}_3$	Ortho.
159. Berthierite.....	$\text{FeS} \cdot \text{Sb}_2\text{S}_3$	Ortho.
160. Matildite.....	$\text{Ag}_2\text{S} \cdot \text{Bi}_2\text{S}_3$	Ortho.
161. Miargyrite.....	$\text{Ag}_2\text{S} \cdot \text{Sb}_2\text{S}_3$	Mono.
162. Lorandite.....	$\text{TiAsS}_2$	Mono.
<b>c. Intermediate</b>		
163. Plagionite.....	$5\text{PbS} \cdot 4\text{Sb}_2\text{S}_3$	Mono.
164. Schirmerite.....	$3(\text{Ag}, \text{Pb})\text{S} \cdot 2\text{Bi}_2\text{S}_3$	Ortho.
165. Klaprotholite.....	$3\text{Cu}_2\text{S} \cdot 2\text{Bi}_2\text{S}_3$	Ortho.
166. Binnite.....	$3\text{Cu}_2\text{S} \cdot 2\text{As}_2\text{S}_3$	Regular
167. Warrenite.....	$3\text{PbS} \cdot 2\text{Sb}_2\text{S}_3$	Ortho.
168. Jamesonite.....	$\text{Pb}_2\text{Sb}_2\text{S}_5$	Ortho.
169. Dufrenoyite.....	$2\text{PbS} \cdot \text{As}_2\text{S}_3$	Ortho.
170. Rathite.....	$2\text{PbS} \cdot \text{As}_2\text{S}_3$	Ortho.
171. Cosalite.....	$2\text{PbS} \cdot \text{Bi}_2\text{S}_3$	Ortho.
172. Kobellite.....	$2\text{PbS} \cdot (\text{Bi}, \text{Sb})_2\text{S}_3$	Ortho.
173. Brongniardite.....	$\text{PbS} \cdot \text{Ag}_2\text{S} \cdot \text{Sb}_2\text{S}_3$	Regular
174. Semseyite.....	$7\text{PbS} \cdot 3\text{Sb}_2\text{S}_3$	Mono.
175. Schapbachite.....	$\text{PbS} \cdot \text{Ag}_2\text{S} \cdot \text{Bi}_2\text{S}_3$	Ortho.
176. Freieslebenite.....	$(\text{Pb}, \text{Ag})_5 \cdot \text{Sb}_4\text{S}_{11}$	Mono.
177. Diaphorite.....	$(\text{Pb}, \text{Ag})_5 \cdot \text{Sb}_4\text{S}_{11}$	Ortho.
180. Boulangerite.....	$\text{Pb}_7\text{Sb}_4\text{S}_{15}$	Ortho.
181. Embrithite.....	$10\text{PbS} \cdot 3\text{Sb}_2\text{S}_3$	Ortho.
<b>d. Ortho</b>		
182. Bournonite.....	$(\text{Pb}, \text{Cu})_3\text{Sb}_2\text{S}_6$	Ortho.
183. Aikinite.....	$3(\text{Pb}, \text{Cu})\text{S} \cdot \text{Bi}_2\text{S}_3$	Ortho.
184. Wittichenite.....	$3\text{Cu}_2\text{S} \cdot \text{Bi}_2\text{S}_3$	Ortho.
185. Stylopyrite.....	$3(\text{Cu}_2, \text{Ag}_2, \text{Fe})\text{S} \cdot \text{Sb}_2\text{S}_3$	Ortho.
186. Lillianite.....	$3\text{PbS} \cdot \text{BiSbS}_3$	Ortho.
187. Guitermanite.....	$3\text{PbS} \cdot \text{As}_2\text{S}_3$	Ortho.
188. Tapalpite.....	$3\text{Ag}_2(\text{S}, \text{Te}) \cdot \text{Bi}_2(\text{S}, \text{Te})_3$	Ortho.
189. Pyrargyrite.....	$\text{Ag}_3\text{SbS}_4$	Hexag.

LIST OF MINERALS

No.	Color	Hardness	Gravity	Locality	Chief Constituent or Use
147.	Lead gray	2	4.8	Mexico	Mercury
148.	Lead gray	.....	6.9	Chiviato	Lead
149.	Bluish black	.....	6	Colorado	Copper
150.	Lead gray	2.5	6	Hungary	Lead
151.	Steel gray	3	5	Hartz	Lead
152.	Dark gray	3	5	Felsobanya	Silver
153.					
154.	Dark gray	3	5	Binnenthal	Lead
155.					
156.	Tin white	2	6	Saxony	Lead
157.	Gray	3	4.75	Hartz	
158.	Lead gray	3	6.9	Sweden	Antimony
159.	Steel gray	2	4	Saxony	
160.	Gray	2	6.9	Peru	Silver
161.	Iron black	2	5	Saxony	
162.	Red	2	5	Allchar	Thallium
163.	Lead gray	2.5	5	Wolfsberg	Lead
164.	Lead gray	2	6	Colorado	Silver
165.	Steel gray	2	4.6	Wittichen	Copper
166.	Steel gray	2.5	4	Tyrol	
167.	Grayish black	.....	.....	Colorado	Lead
168.	Steel gray	2	5.5	Cornwall	
169.	Lead gray	3	5.5	Tyrol	Lead
170.	Lead gray	3	5.5	Tyrol	
171.	Steel gray	2.5	6	Mexico	Silver
172.	Steel gray	.....	6	Sweden, Colorado	
173.	Black	3.5	5.9	Mexico	Lead
174.	Gray	.....	5.9	Hungary	Bismuth
175.	Lead gray	3.5	6	Schapbach	Silver
176.	Steel gray	2	6	Frieberg	
177.	Gray	2.5	5.9	Bohemia	Lead
180.	Lead gray	2.5	5.7	France	
181.	Lead gray	2.5	2.5	Nerchinsk	
182.	Gray	2.5	5.7	Hartz	Lead
183.	Lead gray	2	6	Urals	Bismuth
184.	Steel gray	3	4.5	Baden	
185.	Iron black	3	4.7	Chile	Antimony
186.	Steel gray	.....	.....	Sweden	Lead
187.	Bluish gray	3	5.9	Colorado	
188.	Gray	3	7.8	Mexico	Bismuth
189.	Black	2.5	5.7	Andreasberg	Silver

	Composition	Form
III. SULPHO-SALTS— <i>continued</i>		
190. Proustite. . . . .	$\text{Ag}_3\text{AsS}_3$	Hexag.
191. Sanguinite. . . . .	$\text{Ag}_3\text{AsS}_3$	Hexag.
192. Falkenhaynite. . . . .	$3\text{Cu}_2\text{S} \cdot \text{Sb}_2\text{S}_3$	Regular
193. Pyrostilpnite. . . . .	$3\text{Ag}_2\text{S} \cdot \text{Sb}_2\text{S}_3$	Mono.
194. Rittingerite. . . . .	$\text{Ag}_{10}\text{As}_2\text{Se}_3$	Mono.
<i>e. Basic</i>		
195. Tetrahedrite. . . . .	$\text{Cu}_6\text{Sb}_2\text{S}_7$	Regular
196. Freibergite. . . . .	$\text{Cu}_6\text{Sb}_2\text{S}_7 \cdot \text{Ag}_2\text{S}$	Regular
197. Schwartzite. . . . .	$\text{Cu}_6\text{Sb}_2\text{S}_7 \cdot \text{HgS}$	Regular
198. Tennantite. . . . .	$\text{Cu}_3\text{As}_2\text{S}_7$	Regular
199. Jordanite. . . . .	$4\text{PbS} \cdot \text{As}_2\text{S}_3$	Mono.
200. Meneghinite. . . . .	$4\text{PbS} \cdot \text{Sb}_2\text{S}_3$	Ortho.
201. Stephanite. . . . .	$\text{Ag}_3\text{SbS}_4$	Ortho.
202. Geocronite. . . . .	$5\text{PbS} \cdot \text{Sb}_2\text{S}_3$	Ortho.
203. Beegerite. . . . .	$6\text{PbS} \cdot \text{Bi}_2\text{S}_3$	Regular
204. Kilbrickenite. . . . .	$6\text{PbS} \cdot \text{Sb}_2\text{S}_3$	Massive
205. Polybasite. . . . .	$\text{Ag}_3\text{SbS}_4$	Mono.
206. Pearceite. . . . .	$9\text{Ag}_2\text{S} \cdot \text{As}_2\text{S}_3$	Mono.
207. Polyargyrite. . . . .	$12\text{Ag}_2\text{S} \cdot \text{Sb}_2\text{S}_3$	Regular
<i>2. Sulpharsenates, etc.</i>		
208. Enargite. . . . .	$\text{Cu}_3\text{AsS}_4$	Ortho.
209. Clarite. . . . .	$\text{Cu}_3\text{AsS}_4$	Mono.
210. Luzonite. . . . .	$\text{Cu}_3\text{AsS}_4$	Massive
211. Famatinite. . . . .	$3\text{Cu}_2\text{S} \cdot \text{Sb}_2\text{S}_3$	Ortho.
212. Xanthoconite. . . . .	$3\text{Ag}_2\text{S} \cdot \text{As}_2\text{S}_3$	Hexag.
213. Epiboulangerite. . . . .	$3\text{PbS} \cdot \text{Sb}_2\text{S}_3$	Ortho.
214. Epigenite. . . . .	$4\text{Cu}_2\text{S} \cdot 3\text{FeS} \cdot \text{As}_2\text{S}_3$	Ortho.
215. Stannite. . . . .	$\text{Cu}_3\text{FeSnS}_4$	Regular
216. Argyrodite. . . . .	$\text{Ag}_3\text{GeS}_4$	Regular
217. Canfieldite. . . . .	$\text{Ag}_3\text{SnS}_4$	Regular
218. Franckeite. . . . .	$\text{Pb}_3\text{Sb}_2\text{Sn}_2\text{S}_{12}$	Massive
219. Cyllindrite. . . . .	$\text{Pb}_3\text{Sb}_2\text{Sn}_2\text{S}_{12}$	Massive
220. Sulvanite. . . . .	$3\text{Cu}_2\text{S} \cdot \text{V}_2\text{S}_5$	Massive

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
190.	Scarlet	2	5.5	Freiberg	} Silver
191.	Red	2	.....	Chile	
192.	Gray black	.....	4.8	Joachimsthal	} Copper
193.	Red	2	4	Andreasberg	
194.	Iron black	2	5.6	Chile	} Silver
195.	Iron black	3	4	Hartz	} Copper
196.	Steel gray	.....	4.8	Hartz	
197.	Iron black	.....	5	Hartz	} Copper
198.	Iron black	3	4	Freiberg	
199.	Lead gray	3	6	Tyrol	} Lead
200.	Lead gray	2.5	6	Tuscany	
201.	Iron black	2	6	Freiberg	} Silver
202.	Lead gray	2.5	6	Sweden	
203.	Gray	.....	6	Colorado	} Lead
204.	Lead gray	.....	6	Ireland	
205.	Iron black	2	6	Mexico	} Silver
206.	Iron black	3	6	Colorado, Montana	
207.	Iron Black	2.5	6.9	Wolfach	
208.	Black	3	4	Peru	} Copper
209.	Gray	3.5	4	Baden	
210.	Steel gray	3.5	4	Luzon	
211.	Gray	3.5	4.5	Argentina	} Silver
212.	Orange yellow	2	5	Freiberg	
213.	Gray	.....	6	Altenberg	} Lead
214.	Steel gray	3.5	.....	Baden	
215.	Steel gray	4	4	South Dakota	} Copper
216.	Steel gray	2.5	6	Freiberg	
217.	Black	.....	5.5	Bolivia	} Silver
218.	Blackish gray	.....	5.5	Bolivia	
219.	Blackish gray	2.5	5	Bolivia	} Lead
220.	Bronze	3	4	Australia	

	Composition	Form
<b>IV. HALOIDS</b>		
<b>1. Anhydrous</b>		
221. Calomel.....	Hg <sub>2</sub> Cl <sub>2</sub>	Tetrag.
222. Nantokite.....	Cu <sub>2</sub> Cl <sub>2</sub>	Regular
223. Marshite.....	Cu <sub>2</sub> I <sub>2</sub>	Regular
224. Halite.....	NaCl	Regular
225. Huantajayite.....	20NaCl·AgCl	Regular
226. Sylvite.....	KCl	Regular
227. Sal ammoniac.....	NH <sub>4</sub> Cl	Regular
228. Cerargyrite.....	AgCl	Regular
229. Embolite.....	Ag(Br,Cl)	Regular
230. Bromyrite.....	AgBr	Regular
231. Iodobromite.....	2AgCl·2AgBr·AgI	Regular
232. Miersite.....	Ag <sub>2</sub> I <sub>2</sub>	Regular
233. Cuproiodargyrite.....	CuI·AgI	Incrust.
234. Iodyrite.....	AgI	Hexag.
235. Fluorite.....	CaF <sub>2</sub>	Regular
235a. Yttrofluorite.....	(Ca <sub>2</sub> ,Y <sub>2</sub> )F <sub>6</sub>	Regular
236. Hydrophilite.....	CaCl <sub>2</sub>	Regular
237. Chloromagnosite.....	MgCl <sub>2</sub>	Regular
238. Scacchite.....	MnCl <sub>2</sub>	Regular
239. Chloralluminite.....	AlCl <sub>3</sub> ·XH <sub>2</sub> O	Regular
240. Molysite.....	FeCl <sub>2</sub>	Incrust.
241. Sellaite.....	MgF <sub>2</sub>	Tetrag.
242. Lawrencite.....	FeCl <sub>2</sub>	Hexag.
243. Cotunnite.....	PbCl <sub>2</sub>	Ortho.
244. Tysonite.....	(Ce,L <sub>a</sub> ,Di)F <sub>3</sub>	Hexag.
245. Cryolite.....	Na <sub>3</sub> AlF <sub>6</sub>	Mono.
246. Chiolite.....	5NaF·3AlF <sub>3</sub>	Tetrag.
247. Hieratite.....	2KF·SiF <sub>4</sub>	Regular
<b>2. Oxychlorides, etc.</b>		
248. Atacamite.....	Cu <sub>2</sub> ClH <sub>3</sub> O <sub>3</sub>	Ortho.
249. Percylite.....	PbCuO <sub>2</sub> H <sub>2</sub> Cl <sub>2</sub>	Regular
249a. Boleite.....	PbCuCl <sub>2</sub> (OH) <sub>2</sub> · $\frac{1}{2}$ AgCl	Regular
249b. Cumengite.....	PbCuCl <sub>2</sub> (OH) <sub>2</sub> · $\frac{1}{2}$ AgCl	Tetrag.
250. Matlockite.....	Pb <sub>2</sub> OCl <sub>2</sub>	Tetrag.
251. Mendipite.....	Pb <sub>2</sub> O <sub>2</sub> Cl <sub>2</sub>	Ortho.
252. Laurionite.....	PbClOH	Ortho.
253. Fiedlerite.....	PbClOH	Mono.
254. Penfieldite.....	Pb <sub>2</sub> OCl <sub>2</sub>	Hexag.
255. Daviesite.....	Pb·O·Cl	Ortho.
256. Schwartzembergite.....	Pb(I,Cl) <sub>2</sub> ·2PbO	Hexag.
257. Fluocerite.....	(Ce,L <sub>a</sub> ,Di) <sub>2</sub> OF <sub>4</sub>	Hexag.
258. Nocerite.....	2(Ca,Mg)Fe·(Ca,Mg)O	Hexag.
259. Daubreeite.....	2Bi <sub>2</sub> O <sub>3</sub> ·BiCl <sub>3</sub> ·3H <sub>2</sub> O	Amorph.
<b>3. Hydrous</b>		
260. Carnallite.....	KMgCl <sub>2</sub> ·6H <sub>2</sub> O	Ortho.
261. Douglasite.....	2KCl·FeCl <sub>2</sub> ·2H <sub>2</sub> O	Mono.
262. Bischofite.....	MgCl <sub>2</sub> ·6H <sub>2</sub> O	Mono.
263. Kremersite.....	KCl·NH <sub>4</sub> Cl·FeCl <sub>2</sub> ·H <sub>2</sub> O	Regular

LIST OF MINERALS

No.	Color	Hardness	Gravity	Locality	Chief Constituent or Use
221.	Gray	1	6	Spain	Medicine
222.	Colorless	2.5	3.9	Chile	Copper
223.	Oil brown	.....	.....	New South Wales	
224.	Colorless	2.5	2	Kansas, Louisiana	Salt
225.	White	2	.....	Chile	
226.	Colorless	2	1.9	Stassfurt	Potassium
227.	White	1.5	1.5	Vesuvius	Medicine
228.	Pearl gray	1	5.5	Colorado, Nevada	Silver
229.	Grayish green	1	5	Chile	
230.	Yellow	2	5.8	Mexico	Iodine
231.	Greenish	1.5	5.7	Nassau	
232.	Yellow	.....	.....	New South Wales	Flux
233.	Yellow	2	5.6	Peru	
234.	Yellow	1.5	5.6	New Mexico	Yttrium, fluorine
235.	Blue	4	3	Illinois	
235.	Yellow	4	3	Norway	Chlorine
236.	White	.....	2.5	Vesuvius	
237.	White	.....	.....	Vesuvius	Magnesium
238.	White	.....	.....	Vesuvius	
239.	White	.....	.....	Vesuvius	Chlorine
240.	Red	.....	.....	Vesuvius	
241.	Colorless	5	2.9	Savoy	Fluorine
242.	Green	.....	.....	Meteorites	Iron
243.	White	.....	5	Vesuvius	Lead
244.	Wax yellow	4.5	6	Pike's Peak	Cerium
245.	Colorless	2.5	2.9	Western Greenland	Aluminum
246.	Snow white	3.5	2.8	Ilmen Mts.	
247.	Gray	.....	.....	Vulcano	Potassium
248.	Green	3	3.7	Arizona	Copper
249.	Blue	2.5	.....	Mexico	
249a.	Indigo blue	3	5	Lower California	Lead and copper
249b.	.....	.....	.....	.....	
250.	Yellowish	2.5	7	Cromford	Lead
251.	White	2.5	7	England	
252.	Colorless	3	.....	Greece	Lead
253.	Colorless	.....	.....	Greece	
254.	White	.....	.....	Greece	Cerium
255.	Colorless	.....	.....	Sierra Gorda	
256.	Honey yellow	2	6	Atacama	Fluorine
257.	Yellow	4	5.7	Sweden	
258.	White	.....	.....	Italy	Bismuth
259.	Yellow	2	6	Bolivia	
260.	White	1	1.6	Stassfurt	Magnesium
261.	Colorless	.....	2	Stassfurt	
262.	Colorless	1	1.6	Prussia	Chlorine
263.	Red	.....	.....	Vesuvius	



## COMPREHENSIVE

	Composition	Form
IV. HALOIDS— <i>continued</i>		
264. Erythrosiderite.....	$2\text{KCl} \cdot \text{FeCl}_3 \cdot \text{H}_2\text{O}$	Ortho.
265. Tachydrate.....	$\text{CaCl}_2 \cdot 2\text{MgCl}_2 \cdot 12\text{H}_2\text{O}$	Hexag.
266. Fluellite.....	$\text{AlF}_3 \cdot \text{H}_2\text{O}$	Hexag.
267. Prosopite.....	$\text{CaF}_2 \cdot 2\text{Al}(\text{F},\text{OH})_3$	Mono.
268. Pachnolite.....	$\text{NaF} \cdot \text{CaF}_2 \cdot \text{AlF}_3 \cdot \text{H}_2\text{O}$	Mono.
269. Thomsenelite.....	$\text{NaCaAlF}_6 \cdot \text{H}_2\text{O}$	Mono.
270. Gearsutite.....	$\text{CaF}_2 \cdot \text{Al}(\text{F},\text{OH})_3 \cdot \text{H}_2\text{O}$	Earthy
271. Ralstonite.....	$(\text{Na}_2\text{Mg})\text{F}_2 \cdot 3\text{Al}(\text{F},\text{OH})_3 \cdot 2\text{H}_2\text{O}$	Regular
272. Tallingite.....	$\text{Cu}_5(\text{OH})_8\text{Cl}_2 \cdot 4\text{H}_2\text{O}$	Botry.
273. Footeite.....	$8\text{Cu}(\text{OH})_2 \cdot \text{CuCl}_2 \cdot 4\text{H}_2\text{O}$	Mono.
274. Yttrocerite.....	$(\text{Y},\text{Er},\text{Ce})\text{F}_3 \cdot 5\text{CaF}_2 \cdot \text{H}_2\text{O}$	Earthy

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
264.	Red	.....	.....	Vesuvius	} Chlorine
265.	Yellow	.....	.....	Stassfurt	
266.	Colorless	3	2	Cornwall	} Fluorine
267.	Colorless	4.5	2.8	Colorado	
268.	Colorless	3	2.9	Colorado	
269.	Colorless	2	2.9	Colorado	
270.	White	2	.....	Colorado	} Copper
271.	Colorless	4.5	2.5	Greenland	
272.	Blue	3	3.5	Cornwall	} Yttrium
273.	Blue	.....	.....	Arizona	
274.	Blue	4	3	Sweden	

## COMPREHENSIVE

	Composition	Form
V. OXIDES		
1. <i>Oxides of Silicon</i>		
275. Quartz .....	SiO <sub>2</sub>	Hexag.
276. Star quartz .....	SiO <sub>2</sub>	Hexag.
277. Amethyst .....	SiO <sub>2</sub>	Hexag.
278. Rose quartz .....	SiO <sub>2</sub>	Hexag.
279. Citrine .....	SiO <sub>2</sub>	Hexag.
280. Cairngorm .....	SiO <sub>2</sub>	Hexag.
281. Milky quartz .....	SiO <sub>2</sub>	Hexag.
282. Sapphire quartz .....	SiO <sub>2</sub>	Hexag.
283. Sagenitic .....	SiO <sub>2</sub>	Hexag.
284. Cat's eye .....	SiO <sub>2</sub>	Hexag.
285. Aventurine .....	SiO <sub>2</sub>	Hexag.
286. Chalcedony .....	SiO <sub>2</sub>	Crypto.
287. Carnelian .....	SiO <sub>2</sub>	Crypto.
288. Chrysoprase .....	SiO <sub>2</sub>	Crypto.
289. Prase .....	SiO <sub>2</sub>	Crypto.
290. Plasma .....	SiO <sub>2</sub>	Crypto.
291. Agate .....	SiO <sub>2</sub>	Crypto.
292. Onyx .....	SiO <sub>2</sub>	Crypto.
293. Sardonyx .....	SiO <sub>2</sub>	Crypto.
294. Siliceous sinter .....	SiO <sub>2</sub>	Crypto.
295. Flint .....	SiO <sub>2</sub>	Crypto.
296. Hornstone .....	SiO <sub>2</sub>	Crypto.
297. Basanite .....	SiO <sub>2</sub>	Crypto.
298. Jasper .....	SiO <sub>2</sub>	Crypto.
299. Quartzite .....	SiO <sub>2</sub>	Crypto.
300. Itacolunite .....	SiO <sub>2</sub>	Crypto.
301. Buhrstone .....	SiO <sub>2</sub>	Crypto.
302. Silicified wood .....	SiO <sub>2</sub>	Crypto.
303. Quartzine .....	SiO <sub>2</sub>	Triclinic
304. Tridymite .....	SiO <sub>2</sub>	Hexag.
305. Asmanite .....	SiO <sub>2</sub>	Ortho.
306. Cristobalite .....	SiO <sub>2</sub>	Regular
307. Melanophlogite .....	SiO <sub>2</sub> · O <sub>2</sub>	Regular
308. Opal .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
309. Precious opal .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
310. Fire opal .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
311. Girasol .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
312. Resin opal .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
313. Hydrophane .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
314. Milk opal .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
315. Cacholong .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
316. Opal agate .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
317. Menillite .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
318. Wood opal .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
319. Hyalite .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
320. Pearl sinter .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.
321. Geyserite .....	SiO <sub>2</sub> · H <sub>2</sub> O	Amorph.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
275.	Colorless	7	2.6	Ubiquitous	Abrasives
276.	Whitish	7	2.6	South America	
277.	Purple	7	2.6	Colorado	
278.	Pink	7	2.6	Black Hills	Ornaments
279.	Yellow	7	2.6	Colorado	
280.	Brown	7	2.6	Scotland	
281.	Milk white	7	2.6	Alleghanies	
282.	Indigo	7	2.6	Brazil	
283.	Colorless	7	2.6	Brazil	Various Gems Gems
284.	Milky	7	2.6	Ceylon	
285.	Red	7	2.6	Colorado	
286.	Various	7	2.6	Ubiquitous	Ornaments
287.	Red.	7	2.6	Brazil	
288.	Green	7	2.6	Colorado	
289.	Leek green	7	2.6	Saxony	Ornaments
290.	Green	7	2.6	India	
291.	Banded	7	2.6	Colorado	
292.	Black and white	7	2.6	Colorado	Rock forming Arrow points
293.	Banded	7	2.6	Colorado	
294.	White	7	2.6	Yellowstone Park	
295.	Brown	7	2.6	Chalk Cliffs	Abrasives
296.	Brown	7	2.6	Wyoming	
297.	Black	7	2.6	California	
298.	Red	7	2.6	Colorado	Ornaments Rock forming Curios
299.	Various	7	2.6	Wisconsin	
300.	Gray	7	2.6	North Carolina	
301.	Gray	7	2.6	North Carolina	Millstones
302.	Various	7	2.6	Arizona	
303.	Various	7	2.6	Colorado	
304.	Colorless	7	2	Yellowstone Park	Ornaments
305.	Colorless	.....	.....	Meteorites	
306.	White	6.5	2	Mexico	
307.	Brown	6.5	2	Sicily	Gems
308.	Various	5.5	1.9	Hungary	
309.	Various	5.5	1.9	Hungary	
310.	Red to yellow	5.5	.....	Mexico	Gems
311.	Bluish white	5.5	.....	Mexico	
312.	Yellow	5.5	.....	Mexico	
313.	White	5.5	1.9	Mountains	Gems
314.	Milk white	5.5	1.9	Mountains	
315.	White	5.5	1.9	Hungary	
316.	Light	5.5	1.9	Mountains	.....
317.	Amber	5.5	1.9	France	
318.	Dull grayish	5.5	1.9	Colorado	
319.	Colorless	5.5	1.9	New Jersey, Conn.	.....
320.	White	5.5	1.9	Yellowstone Park	.....
321.	White	5.5	1.9	Yellowstone Park	.....

	Composition	Form
<b>V. OXIDES—continued</b>		
322. Float stone.....	$\text{SiO}_2 \cdot \text{H}_2\text{O}$	Amorph.
323. Tripolite.....	$\text{SiO}_2 \cdot \text{H}_2\text{O}$	Amorph.
324. Infusorial earth.....	$\text{SiO}_2 \cdot \text{H}_2\text{O}$	Amorph.
<b>2. Semi-Metals</b>		
325. Arsenolite.....	$\text{As}_2\text{O}_3$	Regular
326. Claudetite.....	$\text{As}_2\text{O}_3$	Mono.
327. Senarmontite.....	$\text{Sb}_2\text{O}_3$	Regular
328. Valentinite.....	$\text{Sb}_2\text{O}_3$	Ortho.
329. Bismite.....	$\text{Bi}_2\text{O}_3$	Ortho.
330. Tellurite.....	$\text{TeO}_2$	Ortho.
331. Molybdite.....	$\text{MoO}_3$	Ortho.
332. Tungstite.....	$\text{WO}_3$	Ortho.
333. Cervantite.....	$\text{Sb}_2\text{O}_3 \cdot \text{Sb}_2\text{O}_5$	Ortho.
334. Stibiconite.....	$\text{H}_2\text{Sb}_2\text{O}_5$	Massive
<b>3. Metals</b>		
<b>a. Protoxides</b>		
335. Cuprite.....	$\text{Cu}_2\text{O}$	Hexag.
336. Chalcotrichite.....	$\text{Cu}_2\text{O}$	Hexag.
337. Tile ore.....	$\text{Cu}_2\text{O}$	Hexag.
338. Ice.....	$\text{H}_2\text{O}$	Hexag.
339. Periclase.....	$\text{MgO}$	Regular
340. Manganosite.....	$\text{MnO}$	Regular
341. Bunsenite.....	$\text{NiO}$	Regular
342. Zincite.....	$\text{ZnO}$	Hexag.
343. Massicot.....	$\text{PbO}$	Massive
344. Tenorite.....	$\text{CuO}$	Mono.
345. Paramelaconite.....	$\text{CuO}$	Tetrag.
<b>b. Sesquioxides</b>		
346. Corundum.....	$\text{Al}_2\text{O}_3$	Hexag.
347. Sapphire.....	$\text{Al}_2\text{O}_3$	Hexag.
348. Ruby.....	$\text{Al}_2\text{O}_3$	Hexag.
349. Emery.....	$\text{Al}_2\text{O}_3$	Hexag.
350. Hematite.....	$\text{Fe}_2\text{O}_3$	Hexag.
351. Specular hematite....	$\text{Fe}_2\text{O}_3$	Hexag.
352. Columnar hematite...	$\text{Fe}_2\text{O}_3$	Hexag.
353. Red ochrous hematite	$\text{Fe}_2\text{O}_3$	Hexag.
354. Clay ironstone.....	$\text{Fe}_2\text{O}_3$	Hexag.
355. Martite.....	$\text{Fe}_2\text{O}_3$	Regular
356. Ilmenite.....	$\text{FeTiO}_3$	Hexag.
357. Pyrophanite.....	$\text{MnTiO}_3$	Hexag.
<b>c. Intermediate Oxides</b>		
358. Spinel.....	$\text{MgAl}_2\text{O}_4$	Regular
359. Ruby spinel.....	$\text{MgAl}_2\text{O}_4$	Regular
360. Ceylonite-pleonaste...	$(\text{Mg}, \text{Fe})\text{O} \cdot \text{Al}_2\text{O}_3$	Regular

## LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
322.	White	5.5	1.9	Yellowstone Park	} Abrasives
323.	White	5.5	1.9	Virginia	
324.	Gray	5.5	1.9	Missouri	
325.	Colorless	.....	3.7	California	Drugs
326.	White	2.5	3.8	Portugal	Arsenic
327.	White	2	5.3	Quebec	} Antimony
328.	White	2.5	5	New Brunswick	
329.	Straw yellow	.....	4.3	Cornwall	Bismuth
330.	White	2	5.9	Boulder, Colo.	Tellurium
331.	Straw yellow	1	4.5	Pennsylvania	Molybdenum
332.	Yellow	.....	.....	North Carolina	Tungsten
333.	Yellow	4	4	Spain	} Antimony
334.	Yellow	4	5	Arkansas	
335.	Red	3.5	5.8	W. United States	} Copper
336.	Red	3.5	5.8	Arizona	
337.	Brown	3.5	5.8	Arizona	
338.	Colorless	1	0.9	Cold regions	Ice
339.	Grayish	6	3.6	Sweden	Magnesium
340.	Green	5	5	Sweden	Manganese
341.	Green	5	6	Johanngeorgen- stadt	Nickel
342.	Red	4	5	New Jersey	Zinc
343.	Yellow	2	8	Mexico	Lead
344.	Black	3	5.8	Tennessee	} Copper
345.	Black	5	5.8	Arizona	
346.	Various	9	4	Appalachians	Abrasives
347.	Blue	9	4	Ceylon	} Gems
348.	Red	9	4	Upper Burma	
349.	Black	9	4	New York	Abrasives
350.	Red	6	5	New York	} Iron
351.	Black	6	5	Elba	
352.	Brownish red	6	5	New York	
353.	Red	3	5.4	Minnesota	} Iron
354.	Brownish black	3	3	Minnesota	
355.	Iron black	6	4.8	East U.S.	} Iron
356.	Iron black	5	4.5	East U.S.	
357.	Red	5	4.5	Sweden	Manganese
358.	Red	8	3.5	New York	} Gems
359.	Red	8	3.6	New York	
360.	Brown	8	3.5	New York	

	Composition	Form
<b>V. OXIDES—continued</b>		
361. Chlorospinel.....	$\text{MgO} \cdot (\text{Al, Fe})_2\text{O}_3$	Regular
362. Picotite-chrome spinel	$(\text{Mg, Fe})\text{O} \cdot (\text{Al, Fe, Cr})_2\text{O}_3$	Regular
363. Hercynite.....	$\text{FeAl}_2\text{O}_4$	Regular
364. Gahnite.....	$\text{ZnAl}_2\text{O}_4$	Regular
365. Automolite.....	$\text{ZnAl}_2\text{O}_4$	Regular
366. Dysluite.....	$(\text{Zn, Fe, Mn})\text{O} \cdot (\text{Al, Fe})_2\text{O}_3$	Regular
367. Kreittonite.....	$(\text{Zn, Fe, Mg})\text{O} \cdot (\text{Al, Fe})_2\text{O}_3$	Regular
368. Magnetite.....	$\text{FeO} \cdot \text{Fe}_2\text{O}_3$	Regular
369. Franklinite.....	$(\text{Fe, Zn, Mn})\text{O} \cdot (\text{Fe, Mn})_2\text{O}_3$	Regular
370. Magnesioferrite.....	$\text{MgFeO}_4$	Regular
371. Jacobsite.....	$(\text{Mn, Mg})\text{O} \cdot (\text{Fe, Mn})_2\text{O}_3$	Regular
372. Chromite.....	$\text{FeO} \cdot \text{Cr}_2\text{O}_3$	Regular
373. Chrysoberyl.....	$\text{BeAl}_2\text{O}_4$	Hexag.
374. Alexandrite.....	$\text{BeAl}_2\text{O}_4$	Hexag.
375. Cat's eye.....	$\text{BeAl}_2\text{O}_4$	Hexag.
376. Hausmannite.....	$\text{Mn}_3\text{O}_4$	Tetrag.
377. Minium.....	$\text{Pb}_3\text{O}_4$	Powder.
378. Crednerite.....	$\text{Cu}_2\text{Mn}_2\text{O}_5$	Mono.
379. Pseudobrookite.....	$\text{Fe}_4(\text{TiO}_4)_3$	Ortho.
380. Braunitz.....	$3\text{Mn}_2\text{O}_3 \cdot \text{MnSiO}_3$	Tetrag.
381. Bixbyite.....	$\text{FeO} \cdot \text{MnO}_2$	Regular
<b>d. Dioxides</b>		
382. Cassiterite.....	$\text{SnO}_2$	Tetrag.
383. Stream tin.....	$\text{SnO}_2$	Tetrag.
384. Polianite.....	$\text{MnO}_2$	Tetrag.
385. Rutile.....	$\text{TiO}_2$	Tetrag.
386. Nigrine.....	$\text{TiO}_2 (+ 2\% \text{Fe}_2\text{O}_3)$	Tetrag.
387. Ilmenorutile.....	$\text{TiO}_2 (+ 10\% \text{Fe}_2\text{O}_3)$	Tetrag.
388. Plattnerite.....	$\text{PbO}_2$	Tetrag.
389. Baddeleyite.....	$\text{ZrO}_2$	Mono.
390. Octahedrite.....	$\text{TiO}_2$	Tetrag.
391. Brookite.....	$\text{TiO}_2$	Ortho.
392. Pyrolusite.....	$\text{MnO}_2$	Amorph.
<b>e. Hydrrous Oxides</b>		
393. Diaspore.....	$\text{Al}_2\text{O}_3 \cdot \text{H}_2\text{O}$	Ortho.
394. Goethite.....	$\text{Fe}_2\text{O}_3 \cdot \text{H}_2\text{O}$	Ortho.
395. Manganite.....	$\text{Mn}_2\text{O}_3 \cdot \text{H}_2\text{O}$	Ortho.
396. Limonite.....	$2\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$	Amorph.
397. Bog ore.....	$2\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$	Amorph.
398. Clay ironstone.....	$2\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$	Amorph.
399. Turgite.....	$2\text{Fe}_2\text{O}_3 \cdot \text{H}_2\text{O}$	Amorph.
400. Xanthosiderite.....	$\text{Fe}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$	Amorph.
401. Bauxite.....	$\text{Al}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$	Grains
402. Wocheinite.....	$\text{Al}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$	Grains
403. Brucite.....	$\text{MgO} \cdot \text{H}_2\text{O}$	Hexag.
404. Pyrochroite.....	$\text{MnO} \cdot \text{H}_2\text{O}$	Hexag.
405. Gibbsite.....	$\text{Al}_2\text{O}_3 \cdot \text{H}_2\text{O}$	Mono.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
361.	Green	8	3.5	New York	} Gems
362.	Brown	8	4	N.Y. and N.J.	
363.	Black	7.5	3.9	Ronsberg	} Iron
364.	Green	7.5	4	New Jersey	
365.	Green	7.5	4	Sweden	} Zinc
366.	Brown	7.5	4	Pennsylvania	
367.	Black	7	4	Brazil	} Iron
368.	Iron black	5.5	5	Adirondacks	
369.	Black	6	5	New Jersey	} Zinc
370.	Iron black	6	4.5	Vesuvius	
371.	Black	6	4.7	Sweden	} Magnesium
372.	Black	5.5	4.5	W. United States	
373.	Green	8.5	3.5	Urals	} Manganese
374.	Green	8.5	3.6	Urals	
375.	Greenish	8.5	3.6	Ceylon	} Chromium
376.	Black	5	4.8	Sweden	
377.	Red	2	4.6	Baden	} Lead
378.	Iron black	4.5	4.9	Friedrichsrode	
379.	Dark brown	6	4	Transylvania	} Titanium
380.	Dark brown	6	4.7	Hartz	
381.	Black	6	4.9	Utah	} Manganese
382.	Black	7	7	Malay Peninsula	} Tin
383.	Black	7	7	Malay Peninsula	
384.	Steel gray	6	4.9	Bohemia	} Manganese
385.	Red	6.5	4	Arkansas	
386.	Black	6.5	4	Arkansas	} Titanium
387.	Black	.....	5	Ilmen Mts.	
388.	Iron black	5.5	8.5	Idaho	} Lead
389.	Colorless	6.5	5.5	Ceylon	
390.	Brown	5	3.8	Rhode Island	} Zirconium
391.	Brown	5.5	3.8	Arkansas	
392.	Black	2	4.7	Alabama	} Titanium
393.	White	6.5	3	North Carolina	} Aluminum
394.	Brown	5	4	Pa., Colorado	
395.	Black	4	4	Colorado	} Iron
396.	Brown	5.5	3.8	Minnesota	
397.	Brown	2	3.8	Minnesota	} Iron
398.	Brown	2	3.8	Widespread	
399.	Brown	2	4.1	Connecticut	} Iron
400.	Yellow	2.5	4.1	Hartz Mts.	
401.	Gray	3	2.5	Arkansas	} Aluminum
402.	Gray	3	2.5	Carniola	
403.	White	2.5	2	New York	} Magnesium
404.	White	2.5	3.2	New Jersey	
405.	White	2.5	2	New York	} Manganese
					} Aluminum



## COMPREHENSIVE

	Composition	Form
V. OXIDES— <i>continued</i>		
406. Sassolite . . . . .	$B_2O_3 \cdot 3H_2O$	Triclin.
407. Hydrotalcite . . . . .	$Al_2O_3 \cdot 6MgO \cdot 15H_2O$	Hexag.
408. Pyroaurite . . . . .	$Fe_2O_3 \cdot 6MgO \cdot 15H_2O$	Hexag.
409. Chalcophanite . . . . .	$(Mn,Zn)O \cdot 2MnO_2 \cdot 2H_2O$	Hexag.
410. Psilomelane . . . . .	$H_4MnO_5$	Massive
411. Wad . . . . .	$H_4MnO_5 \cdot H_2O$	Amorph.
412. Bog manganese . . . . .	$H_4MnO_5 \cdot H_2O$	Amorph.
413. Asbolite . . . . .	$H_4MnO_5 \cdot H_2O \cdot CoO$	Amorph.
414. Lampadite . . . . .	$H_4MnO_5 \cdot H_2O \cdot (Co \cdot Cu)O$	Amorph.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
406.	White	1	1.4	California	Boric acid
407.	White	2	2	Norway	
408.	White	.....	.....	Sweden	
409.	Iron black	2.5	3.9	New Jersey	Magnesium
410.	Iron black	5	4	Arkansas	
411.	Black	6	3	Germany	Manganese
412.	Black	6	3	Germany	
413.	Black	6	3	Germany	
414.	Black	.....	.....	Germany	

	Composition	Form
VI. CARBONATES		
1. <i>Anhydrous</i>		
415. Calcite.....	$\text{CaCO}_3$	Hexag.
416. Dog-tooth spar.....	$\text{CaCO}_3$	Hexag.
417. Nail-head spar.....	$\text{CaCO}_3$	Hexag.
418. Iceland spar.....	$\text{CaCO}_3$	Hexag.
419. Fontainebleau lime-stone.....	$\text{CaCO}_3$	Hexag.
420. Satin spar.....	$\text{CaCO}_3$	Hexag.
421. Argentine.....	$\text{CaCO}_3$	Lamellar
422. Aphrite.....	$\text{CaCO}_3$	Lamellar
423. Saccharoid. limestone.	$\text{CaCO}_3$	Crypt.
424. Shell marble.....	$\text{CaCO}_3$	Shelly
425. Lumachelle.....	$\text{CaCO}_3$	Chatoy.
426. Ruin marble.....	$\text{CaCO}_3$	Brecci.
427. Lithographic stone...	$\text{CaCO}_3$	Massive
428. Hydraulic limestone..	$\text{CaCO}_3$ , also $\text{SiO}_2, \text{Al}_2\text{O}_3$ , etc.	Massive
429. Chalk.....	$\text{CaCO}_3$	Massive
430. Oölite.....	$\text{CaCO}_3$	Granular
431. Pisolite.....	$\text{CaCO}_3$	Grains
432. Stalactite.....	$\text{CaCO}_3$	Cylind.
433. Stalagmite.....	$\text{CaCO}_3$	Cylind.
434. Calc sinter.....	$\text{CaCO}_3$	Incrust.
435. Travertine.....	$\text{CaCO}_3$	Incrust.
436. Agaric mineral.....	$\text{CaCO}_3$	Grains
437. Rock meal.....	$\text{CaCO}_3$	Grains
438. Thinolite.....	$\text{CaCO}_3$	Grains
439. Dolomite.....	$(\text{Ca}, \text{Mg})\text{CO}_3$	Hexag.
440. Magnesite.....	$\text{MgCO}_3$	Hexag.
441. Breunnerite.....	$\text{MgCO}_3 \cdot \text{H}_2\text{O}$	Hexag.
442. Mesitite.....	$2\text{MgCO}_3 \cdot \text{FeCO}_3$	Hexag.
443. Pistomesite.....	$\text{MgCO}_3 \cdot \text{FeCO}_3$	Hexag.
445. Siderite.....	$\text{FeCO}_3$	Hexag.
446. Sphaerosiderite.....	$\text{FeCO}_3$	Concret.
447. Rhodochrosite.....	$\text{MnCO}_3$	Hexag.
448. Smithsonite.....	$\text{ZnCO}_3$	Hexag.
449. Sphaerocobaltite.....	$\text{CoCO}_3$	Hexag.
450. Aragonite.....	$\text{CaCO}_3$	Ortho.
451. Flos ferri.....	$\text{CaCO}_3$	Stalact.
452. Tarnowitzite.....	$\text{CaCO}_3 \cdot \text{PbCO}_3$	Stalcat.
453. Witherite.....	$\text{BaCO}_3$	Ortho.
454. Bromlite.....	$(\text{Ba}, \text{Ca})\text{CO}_3$	Ortho.
455. Strontianite.....	$\text{SrCO}_3$	Ortho.
456. Cerussite.....	$\text{PbCO}_3$	Ortho.
457. Barytocalcite.....	$\text{BaCO}_3 \cdot \text{CaCO}_3$	Mono.
458. Bismutospharite.....	$\text{Bi}_2(\text{CO}_3)_3 \cdot 2\text{Bi}_2\text{O}_3$	Spherical
459. Parisite.....	$(\text{CaF})(\text{CeF})\text{Ce}(\text{CO}_3)_2$	Hexag.
460. Bastnäsite.....	$(\text{Ce}, \text{La}, \text{Di})_2\text{C}_2\text{O}_7 \cdot (\text{Ce}, \text{La}, \text{Di})\text{F}$	Hexag.
461. Phosgenite.....	$(\text{PbCl})_2\text{CO}_3$	Tetrag.
462. Northupite.....	$\text{MgCO}_3 \cdot \text{Na}_2\text{CO}_3 \cdot \text{NaCl}$	Regular

## LIST OF MINERALS

No	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
415.	Colorless	3	2.7	Ubiquitous	} Calcium Prisms
416.	Colorless	3	2.7	Missouri	
417.	Colorless	3	2.7	Missouri	
418.	Colorless	3	2.7	Iceland	
419.	Colorless	3	2.7	France	
420.	Colorless	3	2.7	France	} Calcium
421.	White	3	2.7	France	
422.	White	3.5	2.7	France	
423.	Yellow	3	2.7	France	
424.	Yellow	3	2.7	Carinthia	
425.	Dark brown	3	2.7	France	
426.	Brown	3	2.7	Italy	
427.	Buff	3	2.7	Solenhofen	
428.	Buff	3	2.7	Virginia	
429.	White	3	2.7	England	
430.	White	3	2.7	Missouri	
431.	White	3	2.7	Missouri	
432.	White	3	2.7	Kentucky	
433.	White	3	2.7	Kentucky	
434.	White	3	2.7	Yellowstone Park	
435.	White	3	2.7	Tivoli	
436.	White	3	2.7	Caverns	
437.	White	3	2.7	Paris	
438.	Yellow	3	2.7	Nevada	
439.	White	3.5	2.8	Illinois	} Building
440.	Colorless	4	3	Greece	
441.	White	4	3	Massachusetts	} Magnesium
442.	Yellowish	3.5	3	Traversella	
443.	Yellowish	3.5	3	Traversella	} Iron
445.	Gray	3.5	3.8	Germany	
446.	Brown	3.5	3.8	E. United States	
447.	Red	4	3	Colorado	} Manganese
448.	White	5	4	Pennsylvania	
449.	Red	4	4	Saxony	} Zinc
450.	Colorless	3.5	2.9	New York, Illinois	
451.	White	3.5	2.9	New York, Illinois	} Calcium
452.	White	.....	2.9	Silesia	
453.	Colorless	3.5	4.2	England	} Barium
454.	White	4	3.7	England	
455.	Colorless	3.5	3.7	New York	} Strontium
456.	Colorless	3.5	6	Cordilleras	
457.	White	4	3.6	Cumberland	} Lead
458.	Yellow	3	7	Saxony	
459.	Yellow	4.5	4	Colombia	} Cerium
460.	Yellow	4	4.9	Colorado	
461.	White	2.7	6	England	} Lead
462.	White	.....	.....	California	

	Composition	Form
<b>VI. CARBONATES—continued</b>		
<i>2. Hydrous</i>		
463. Teschemacherite.....	$\text{HNH}_4\text{CO}_3$	Ortho.
464. Malachite.....	$\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$	Mono.
465. Azurite.....	$2\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$	Mono.
466. Chessylite.....	$2\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$	Mono.
467. Aurichalcite.....	$2(\text{Zn}, \text{Cu})\text{CO}_3 \cdot 3(\text{Zn}, \text{Cu})(\text{OH})_2$	Mono.
468. Hydrozincite.....	$\text{ZnCO}_3 \cdot 2\text{Zn}(\text{OH})_2$	Earthy
469. Hydrocerussite.....	$2\text{PbCO}_3 \cdot \text{Pb}(\text{OH})_2$	Hexag.
470. Dawsonite.....	$\text{Na}_3\text{Al}(\text{CO}_3)_2 \cdot 2\text{Al}(\text{OH})_3$	Mono.
471. Thermonatrite.....	$\text{Na}_2\text{CO}_3 \cdot \text{H}_2\text{O}$	Ortho.
472. Nesquehonite.....	$\text{MgCO}_3 \cdot 3\text{H}_2\text{O}$	Ortho.
473. Natron.....	$\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$	Mono.
474. Pirssonite.....	$\text{CaCO}_3 \cdot \text{Na}_2\text{CO}_3 \cdot 2\text{H}_2\text{O}$	Ortho.
475. Gaylussite.....	$\text{CaCO}_3 \cdot \text{Na}_2\text{CO}_3 \cdot 5\text{H}_2\text{O}$	Mono.
476. Lanthanite.....	$\text{La}_2(\text{CO}_3)_3 \cdot 9\text{H}_2\text{O}$	Ortho.
477. Trona.....	$\text{Na}_2\text{CO}_3 \cdot \text{HNaCO}_3 \cdot 2\text{H}_2\text{O}$	Mono.
478. Hydromagnesite.....	$3\text{MgCO}_3 \cdot \text{Mg}(\text{OH})_2 \cdot 3\text{H}_2\text{O}$	Amorph.
479. Hydrogioberite.....	$\text{MgCO}_3 \cdot \text{Mg}(\text{OH})_2 \cdot 2\text{H}_2\text{O}$	Compact
480. Lansfordite.....	$3\text{MgCO}_3 \cdot \text{Mg}(\text{OH})_2 \cdot 21\text{H}_2\text{O}$	Triclin.
481. Zaratite.....	$\text{NiCO}_3 \cdot 2\text{Ni}(\text{OH})_2 \cdot 4\text{H}_2\text{O}$	Stalact.
482. Remingtonite.....	$\text{CaCO}_3 \cdot \text{H}_2\text{O}$	Incrust.
483. Tengerite.....	$\text{YCO}_3 \cdot \text{H}_2\text{O}$	Pulver.
484. Bismutite.....	$\text{Bi}_2\text{O}_3 \cdot \text{CO}_2 \cdot \text{H}_2\text{O}$	Amorph.
485. Uranothallite.....	$2\text{CaCO}_3 \cdot \text{U}(\text{CO}_3)_2 \cdot 10\text{H}_2\text{O}$	Ortho.
486. Liebigite.....	$\text{CaCO}_3 \cdot (\text{UO}_2)\text{CO}_3 \cdot 20\text{H}_2\text{O}$	Concret.
487. Voglite.....	$(\text{U}, \text{Ca}, \text{Cu})\text{CO}_3 \cdot \text{H}_2\text{O}$	Scales

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
463.	Yellow	1.5	1.4	Africa	Lead
464.	Green	3.5	4	Arizona	} Copper
465.	Blue	3.5	3.7	Arizona	
466.	Blue	3.5	3.7	France	
467.	Green	2	3.5	France	
468.	White	2	3.5	Pennsylvania	} Zinc
469.	Colorless	2	6	Sweden	Lead
470.	White	3	2	Tuscany	} Aluminum
471.	White	1	1.5	Nevada	
472.	Colorless	2.5	1.8	Pennsylvania	Magnesium
473.	Gray	1	1.4	Egypt	} Sodium
474.	Colorless	3.5	2.3	California	
475.	White	2.3	1.9	Utah	} Lanthanum
476.	White	2.5	2.6	Pennsylvania	
477.	Gray	2.5	2	Nevada	Sodium
478.	White	3.5	2	New Jersey	} Magnesium
479.	Gray	.....	2	Italy	
480.	White	2.5	1.5	Pennsylvania	} Nickel
481.	Green	3	2	Texas	
482.	Rose	2	.....	Maryland	Cobalt
483.	White	.....	.....	Texas	Ytterium
484.	White	4	6.8	South Carolina	Bismuth
485.	Green	2	.....	Bohemia	} Uranium
486.	Green	2	.....	Joachimsthal	
487.	Green	.....	.....	Joachimsthal	

	Composition	Form
<b>VII. SILICATES</b>		
<b>1. Anhydrous</b>		
<b>a. Disilicates</b>		
488. Petalite.....	$\text{LiAl}(\text{Si}_2\text{O}_6)_2$	Mono.
489. Milarite.....	$\text{HKCa}_3\text{Al}_2(\text{Si}_2\text{O}_6)_6$	Hexag.
490. Eudidymite.....	$\text{HNaBeSi}_2\text{O}_6$	Mono.
491. Epididymite.....	$\text{HNaBeSi}_2\text{O}_6$	Ortho.
492. Orthoclase.....	$\text{KAISi}_3\text{O}_8$	Mono.
493. Adularia.....	$\text{KAISi}_3\text{O}_8$	Mono.
494. Valencianite.....	$\text{KAISi}_3\text{O}_8$	Mono.
495. Sanidine.....	$\text{KAISi}_3\text{O}_8$	Mono.
496. Rhyacolite.....	$\text{KAISi}_3\text{O}_8$	Mono.
497. Loxoclase.....	$\text{KAISi}_3\text{O}_8 \cdot 7\text{Na}_2\text{O}$	Mono.
498. Murchisonite.....	$\text{KAISi}_3\text{O}_8$	Mono.
499. Perthite.....	$\text{KAISi}_3\text{O}_8$	Mono.
500. Hyalophane.....	$(\text{K}, \text{Ba})\text{Al}_2(\text{SiO}_3)_4$	Mono.
501. Microcline.....	$\text{KAISi}_3\text{O}_8$	Triclinic
502. Amazonstone.....	$\text{KAISi}_3\text{O}_8$	Triclinic
503. Chesterlite.....	$\text{KAISi}_3\text{O}_8$	Triclinic
504. Anorthoclase.....	$\text{KAISi}_3\text{O}_8$	Triclinic
505. Albite.....	$\text{NaAlSi}_3\text{O}_8$	Triclinic
506. Peristerite.....	$\text{NaAlSi}_3\text{O}_8$	Triclinic
507. Pericline.....	$\text{NaAlSi}_3\text{O}_8$	Triclinic
508. Cleavelandite.....	$\text{NaAlSi}_3\text{O}_8$	Triclinic
509. Oligoclase.....	* $\text{Ab}_2\text{An}_1$	Triclinic
510. Sunstone.....	* $\text{Ab}_2\text{An}_1$	Triclinic
511. Andesine.....	* $\text{Ab}_2\text{An}_1$	Triclinic
512. Labradorite.....	* $\text{Ab}_2\text{An}_1$	Triclinic
513. Maskelynite.....	* $\text{Ab}_2\text{An}_1$	Grains
514. Anorthite.....	$\text{CaAl}_2\text{Si}_2\text{O}_8$	Triclinic
515. Indianite.....	$\text{CaAl}_2\text{Si}_2\text{O}_8$	Triclinic
516. Cyclopite.....	$\text{CaAl}_2\text{Si}_2\text{O}_8$	Triclinic
517. Celsian.....	$\text{BaAl}_2\text{Si}_2\text{O}_8$	Triclinic
<b>b. Metasilicates</b>		
518. Leucite.....	$\text{KAl}(\text{SiO}_3)_2$	Ortho.
519. Pollucite.....	$\text{H}_2\text{Cs}_3\text{Al}_4(\text{SiO}_3)_9$	Regular
520. Enstatite.....	$\text{MgSiO}_3$	Ortho.
521. Chladnite.....	$\text{MgSiO}_3$	Ortho.
522. Bronzite.....	$\text{MgSiO}_3$	Ortho.
523. Hypersthene.....	$(\text{Fe}, \text{Mg})\text{SiO}_3$	Ortho.
524. Bastite.....	$(\text{Fe}, \text{Mg})\text{SiO}_3$	Ortho.
525. Peckhamite.....	$2(\text{Mg}, \text{Fe})\text{SiO}_3 \cdot (\text{Mg}, \text{Fe})\text{SiO}_4$	Ortho.
526. Pyroxene.....	$\text{Ca}(\text{Mg}, \text{Fe})\text{Si}_2\text{O}_6 \cdot (\text{Mg}, \text{Fe})(\text{AlFe})_2\text{Si}_2\text{O}_6$	Mono.
527. Diopside.....	$\text{CaMg}(\text{SiO}_3)_2$	Mono.
528. Malacolite.....	$\text{CaMg}(\text{SiO}_3)_2$	Mono.
529. Alalite.....	$\text{CaMg}(\text{SiO}_3)_2$	Mono.
530. Traversellite.....	$\text{CaMg}(\text{SiO}_3)_2$	Mono.
531. Violan.....	$\text{CaMg}(\text{SiO}_3)_2$	Mono.

\* Ab = Albite; An = Anorthite.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
488.	Colorless	6	2	Massachusetts	Lithium Potassium
489.	Colorless	5	2.5	Switzerland	
490.	White	6	2.5	Norway	
491.	White	5.5	3.5	South Greenland	Rock forming
492.	Colorless	6	2.5	California	
493.	Colorless	6	2.5	Switzerland	
494.	Colorless	6	2.5	Valencia	
495.	Colorless	6	2.5	Valencia	
496.	Colorless	6	2.5	Monte Somma	
497.	Colorless	6	2.5	New York	
498.	Red	6	2.5	England	
499.	Red	6	2.5	Ontario	
500.	Red	6	2.8	Sweden	
501.	White	6	2.5	Pike's Peak	
502.	White	6	2.5	Pike's Peak	
503.	White	6	2.5	Pennsylvania	
504.	White	6	2.5	Pennsylvania	
505.	White	6	2.6	E. United States	
506.	White	6	2.6	E. United States	
507.	White	6	2.6	E. United States	
508.	Bluish	6	2.6	New Hampshire	
509.	White	6	2.6	New York	
510.	White	6	2.6	Norway	
511.	White	5	2.6	Rocky Mts.	
512.	Gray	5	2.7	New York	
513.	Colorless	5	2.7	Meteorites	
514.	White	6	2.7	Mt. Vesuvius	
515.	White	6	2.7	India	
516.	White	6	2.7	Cyclopean Island	
517.	Colorless	6	3	Sweden	
518.	Colorless	5.5	2.5	Vesuvius	Rock forming
519.	White	6.5	2.9	Maine	
520.	Gray	5.5	3	New York	
521.	Gray	5.5	3	Meteorites	
522.	Green	5.5	3	New York	
523.	Brownish green	5	3	New York	
524.	Green	3.5	2.5	Hartz	
525.	Yellow	.....	3.2	Meteorites	
526.	White	5	3	Igneous rocks	
527.	Light green	5.5	3.3	Igneous rocks	
528.	Light green	5.5	3.3	Sweden	
529.	Green	5.5	3.3	Piedmont	
530.	Green	5.5	3.3	Traversella	
531.	Blue	5.5	3.3	Italy	



	Composition	Form
VII. SILICATES— <i>continued</i>		
532. Canaanite.....	$\text{CaMg}(\text{SiO}_3)_2$	Mono.
533. Lavrovite.....	$\text{CaMg}(\text{SiO}_3)_2$	Mono.
534. Hedenbergite.....	$\text{CaFe}(\text{SiO}_3)_2$	Mono.
535. Sahlite.....	$\text{CaFe}(\text{SiO}_3)_2$	Mono.
536. Baikalite.....	$\text{CaFe}(\text{SiO}_3)_2$	Mono.
537. Coccoilite.....	$\text{CaFe}(\text{SiO}_3)_2$	Mono.
538. Diallage.....	$\text{CaFe}(\text{SiO}_3)_2$	Mono.
539. Omphacite.....	$\text{CaFe}(\text{SiO}_3)_2$	Mono.
540. Schefferite.....	$\text{CaMg}(\text{Fe, Mn})(\text{SiO}_3)_2$	Mono.
541. Jeffersonite.....	Like schefferite + Zn	Mono.
542. Augite.....	$\text{CaMg}(\text{SiO}_3)_2$	Mono.
543. Leucaugite.....	$\text{CaMg}(\text{SiO}_3)_2$	Mono.
544. Fassaite.....	$\text{CaMg}(\text{SiO}_3)_2$	Mono.
545. Acmite.....	$\text{NaFe}(\text{SiO}_3)_2$	Mono.
546. Spodumene.....	$\text{LiAl}(\text{SiO}_3)_2$	Mono.
547. Hiddenite.....	$\text{LiAl}(\text{SiO}_3)_2$	Mono.
548. Jadeite.....	$\text{NaAl}(\text{SiO}_3)_2$	Mono.
549. Chloromelanite.....	$\text{NaAl}(\text{SiO}_3)_2$	Mono.
550. Nephrite.....	$\text{NaAl}(\text{SiO}_3)_2$	Mono.
551. Wollastonite.....	$\text{CaSiO}_3$	Mono.
552. Pectolite.....	$\text{HNaCa}_2(\text{SiO}_3)_3$	Mono.
553. Rosenbuschite.....	$6\text{CaSiO}_3 \cdot 2\text{Na}_2\text{ZrO}_2\text{F}_2 \cdot (\text{TiSiO}_3\text{TiO}_3)$	Mono.
554. Wöhlerite.....	$\text{Ca}_{10}\text{Na}_5\text{Fe}_3\text{Nb}_2\text{Zr}_3\text{Si}_{10}\text{O}_{42}$	Mono.
555. Låvenite.....	$(\text{Na, Ca, Mn, Fe})(\text{F, Zr, O})\text{Si}_2\text{O}_6$	Mono.
556. Rhodonite.....	$\text{MnSiO}_3$	Triclinic
557. Bustamite.....	Like rhodonite + Fe, Ca	Triclinic
558. Fowlerite.....	Like rhodonite + Fe, Ca, Zn	Triclinic
559. Babingtonite.....	$(\text{Ca, Fe, Mn})\text{SiO}_3$	Triclinic
560. Hiortdahlite.....	$(\text{Na}_2, \text{Ca})(\text{Si, Zr})\text{O}_3$	Triclinic
561. Anthophyllite.....	$(\text{Mg, Fe})\text{SiO}_3$	Ortho.
562. Gedrite.....	Like anthophyllite + Al	Ortho.
563. Amphibole.....	$\text{CaMgFe}[\text{MnNa}_2\text{K}_2\text{H}_2(\text{SiO}_3)_4]$	Mono.
564. Tremolite.....	$\text{CaMg}_3(\text{SiO}_3)_4$	Mono.
565. Actinolite.....	Like tremolite + Fe	Mono.
566. Nephrite.....	$\text{CaMg}_3(\text{SiO}_3)_4$	Compact
567. Asbestos, amianthus.....	$\text{CaMg}_3(\text{SiO}_3)_4$	Fibrous
568. Mountain leather.....	$\text{CaMg}_3(\text{SiO}_3)_4$	Fibrous
569. Mountain cork.....	$\text{CaMg}_3(\text{SiO}_3)_4$	Fibrous
570. Smaragdite.....	$\text{CaMg}_3(\text{SiO}_3)_4$	Fibrous
571. Uralite.....	$\text{CaMg}_3(\text{SiO}_3)_4$	Fibrous
572. Cummingtonite.....	Like actinolite + Mg	Fibrous
573. Dannemorite.....	Like actinolite + Mn	Fibrous
574. Grunerite.....	$\text{FeSiO}_3$	Fibrous
575. Richterite.....	$(\text{K}_2, \text{Na}_2, \text{Mg, Ca, Mn, Fe})_4(\text{SiO}_3)_4$	Fibrous
576. Breislakite.....	$(\text{K}_2, \text{Na}_2, \text{Mg, Ca, Mn})_4(\text{SiO}_3)_4$	Fibrous
577. Hornblende.....	$\text{Ca}(\text{MgFe})_2\text{SiO}_3 \cdot \text{CaMg}_2\text{Al}_2(\text{SiO}_4)_3$	Mono.
578. Edenite.....	$\text{Ca}(\text{MgFe})_2\text{SiO}_3 \cdot \text{CaMg}_2\text{Al}_2(\text{SiO}_4)_3$	Mono.
579. Koksharovite.....	$\text{Ca}(\text{MgFe})_2\text{SiO}_3 \cdot \text{CaMg}_2\text{Al}_2(\text{SiO}_4)_3$	Mono.
580. Pargasite.....	$\text{Ca}(\text{MgFe})_2\text{SiO}_3 \cdot \text{CaMg}_2\text{Al}_2(\text{SiO}_4)_3$	Mono.

## LIST OF MINERALS

No.	Color	Hardness	Gravity	Locality	Chief Constituent or Use
532.	Gray	5.5	3.3	Connecticut	Rock forming
533.	Green	5.5	3.3	East Siberia	
534.	Black	5.5	3.5	Sweden	
535.	Green	5.5	3.5	Sweden	
536.	Green	5.5	3.5	Siberia	
537.	Dark green	4	3	Mountains	
538.	Green	4	3	Mountains	
539.	Brown	4	3	Mountains	
540.	Brown	4	3	Mountains	
541.	Dark	6	3	New Jersey	
542.	Green	5.5	3.3	E. United States	
543.	White	6.5	3	E. United States	
544.	Green	6.5	3	Vesuvius	
545.	Gray	6	3.5	Colorado	
546.	Green	6.5	3	Massachusetts	Ornaments
547.	Green	6.5	3	Massachusetts	
548.	Green	6.5	3	Asia	
549.	Dark green	6.5	3	Asia	Rock forming
550.	Green	6.5	3	Asia	
551.	White	4.5	2.8	New York	Ornaments
552.	White	5	2.6	New Jersey	
553.	White	5	2.6	Norway	
554.	Yellow	5.5	3.4	Norway	Rock forming
555.	Yellow	6	3.5	Norway	
556.	Red	6	3.5	Russia	
557.	Red	6	3.5	Mexico	Ornaments
558.	Red	6	3.5	New Jersey	
559.	Black	5.5	3	Norway	
560.	Yellow	5.5	3	South Norway	Rock forming
561.	Brown	5.5	3	North Carolina	
562.	Brown	5.5	3	North Carolina	
563.	Green	5	2.9	Mountains	Rock forming
564.	Gray	5	2.9	Mountains	
565.	Green	5	3	Mountains	
566.	Green	6	2.9	Mexico	Ornaments
567.	Gray	3	2.9	Mountains	
568.	Gray	3	2.9	Mountains	Cloth
569.	Gray	3	2.9	Mountains	
570.	Green	3	2.9	Alps	
571.	Green	3	2.9	Alps	Rock forming
572.	Gray	3	3	Massachusetts	
573.	Brown	3	3	Sweden	
574.	Brown	3	3.7	Sweden	Rock forming
575.	Brown	3	3.7	Sweden	
576.	Brown	3	3.7	Vesuvius	
577.	Black	5.5	3	Vesuvius	
578.	Gray	5.5	3	New York	
579.	Gray	5.5	3	New York	
580.	Green	5.5	3	Finland	

	Composition	Form
<b>VII. SILICATES—continued</b>		
581. Kataforite . . . . .	$\text{Ca}(\text{MgFe})_3(\text{SiO}_3)_4 \cdot \text{CaMg}_2\text{Al}_2(\text{SiO}_3)_3$	Mono.
582. Kupfferite . . . . .	$\text{Ca}(\text{MgFe})_3(\text{SiO}_3)_4 \cdot \text{CaMg}_2\text{Al}_2(\text{SiO}_3)_3$	Mono.
583. Syntagmatite . . . . .	$\text{Ca}(\text{MgFe})_3(\text{SiO}_3)_4 \cdot \text{CaMg}_2\text{Al}_2(\text{SiO}_3)_3$	Mono.
584. Bergamaskite . . . . .	$\text{Ca}(\text{MgFe})_3(\text{SiO}_3)_4 \cdot \text{CaMg}_2\text{Al}_2(\text{SiO}_3)_3$ (—Mg)	Mono.
585. Kaersutite . . . . .	Like amphibole + Ti	Mono.
586. Hastingsite . . . . .	Contains much Na	Mono.
587. Glaucophane . . . . .	$\text{NaAl}(\text{SiO}_3)_2 \cdot (\text{Fe}, \text{Mg})\text{SiO}_3$	Mono.
588. Gastaldite . . . . .	$\text{NaAl}(\text{SiO}_3)_2 \cdot (\text{Fe}, \text{Mg})\text{SiO}_3$	Mono.
588a. Riebeckite . . . . .	$2\text{NaFe}(\text{SiO}_3)_2 \cdot \text{FeSiO}_3$	Mono.
589. Crocidolite . . . . .	$\text{NaFe}(\text{SiO}_3)_2 \cdot \text{FeSiO}_3$	Mono.
590. Abriachanite . . . . .	$\text{NaFe}(\text{SiO}_3)_2 \cdot \text{FeSiO}_3$	Amor.
590a. Arfvedsonite . . . . .	$4\text{Na}_2\text{O} \cdot 3\text{CaO} \cdot 14\text{FeO} \cdot (\text{Al}, \text{Fe})_2\text{O}_3 \cdot 21\text{SiO}_2$	Mono.
591. Crossite . . . . .	Like arfvedsonite + Na	Mono.
592. Barkevikite . . . . .	Like arfvedsonite + Na	
593. Aenigmatite . . . . .	$\text{Na}_4\text{Fe}_2\text{AlFe}(\text{SiTi})_{12}\text{O}_{34}$	Triclinic
594. Beryl . . . . .	$\text{Be}_3\text{Al}_2\text{Si}_6\text{O}_{18}$	Hexag.
595. Emerald . . . . .	$\text{Be}_3\text{Al}_2\text{Si}_6\text{O}_{18}$	Hexag.
596. Aquamarine . . . . .	$\text{Be}_3\text{Al}_2\text{Si}_6\text{O}_{18}$	Hexag.
597. Davidsonite . . . . .	$\text{Be}_3\text{Al}_2\text{Si}_6\text{O}_{18}$	Hexag.
598. Eudialyte . . . . .	$\text{Na}_{13}(\text{Ca}, \text{Fe})_6\text{Cl}(\text{Si}, \text{Zr})_{20}\text{O}_{52}$	Hexag.
599. Euclite . . . . .	$\text{Na}_{13}(\text{Ca}, \text{Fe})_6\text{Cl}(\text{Si}, \text{Zr})_{20}\text{O}_{52}$	Hexag.
600. Elpidite . . . . .	$\text{Na}_2\text{O} \cdot \text{ZrO}_2 \cdot 6\text{SiO}_2 \cdot 3\text{H}_2\text{O}$	
601. Catapleite . . . . .	$\text{H}_4(\text{Na}_2, \text{Ca})\text{ZrSi}_5\text{O}_{17}$	Hexag.
602. Cappelinite . . . . .	$3\text{BaSiO}_3 \cdot 2\text{Y}_2(\text{SiO}_3)_3 \cdot 5\text{YBO}_3$	Hexag.
603. Melanocerite . . . . .	$12(\text{H}_2, \text{Ca})\text{SiO}_3 \cdot 3(\text{Y}, \text{Ce})\text{BO}_3 \cdot 2\text{H}_2(\text{Th}, \text{Ce})\text{O}_2 \cdot 8(\text{Ce}, \text{La}, \text{Bi})\text{OF}$	Hexag.
604. Caryocerite . . . . .	$6(\text{H}_2, \text{Ca})\text{SiO}_3 \cdot 2(\text{Ce}, \text{Da}, \text{Y})\text{BO}_3 \cdot 3\text{H}_2(\text{Ce}, \text{Th})\text{O}_2 \cdot 2\text{LaOF}$	Hexag.
605. Streenstrupine . . . . .	Ti, Th, Ce, La, Di, Al, Fe, Mn, Ca, Na, H, silicate	Hexag.
606. Tritomite . . . . .	$2(\text{H}_2, \text{Na}, \text{Ca})\text{SiO}_3 \cdot (\text{Ce}, \text{La}, \text{Di}, \text{Y})\text{BO}_3 \cdot \text{H}_2(\text{Ce}, \text{Th}, \text{Zr})\text{O}_2\text{F}$	Hexag.
607. Leucophanite . . . . .	$\text{Na}(\text{BeF})\text{Ca}(\text{SiO}_3)_2$	Ortho.
608. Meliphanite . . . . .	$\text{NaCa}_2\text{Be}_2\text{FSi}_3\text{O}_{16}$	Tetrag.
609. Iolite . . . . .	$\text{H}_2(\text{Mg}, \text{Fe})_4\text{Al}_3\text{Si}_{10}\text{O}_{37}$	Ortho.
610. Bonsdorffite . . . . .	$\text{H}_2(\text{Mg}, \text{Fe})_4\text{Al}_3\text{Si}_{10}\text{O}_{37}$ , altered	Ortho.
611. Fahllunite . . . . .	$\text{H}_2(\text{Mg}, \text{Fe})_4\text{Al}_3\text{Si}_{10}\text{O}_{37}$ , altered	Ortho.
612. Pyrargillite . . . . .	$\text{H}_2(\text{Mg}, \text{Fe})_4\text{Al}_3\text{Si}_{10}\text{O}_{37}$ , altered	Ortho.
613. Esmarkite . . . . .	$\text{H}_2(\text{Mg}, \text{Fe})_4\text{Al}_3\text{Si}_{10}\text{O}_{37}$ , altered	Ortho.
614. Raumite . . . . .	$\text{H}_2(\text{Mg}, \text{Fe})_4\text{Al}_3\text{Si}_{10}\text{O}_{37}$ , altered	Ortho.
615. Chlorophyllite . . . . .	$\text{H}_2(\text{Mg}, \text{Fe})_4\text{Al}_3\text{Si}_{10}\text{O}_{37}$ , altered	Ortho.
616. Aspasolite . . . . .	$\text{H}_2(\text{Mg}, \text{Fe})_4\text{Al}_3\text{Si}_{10}\text{O}_{37}$ , altered	Ortho.
617. Polychroillite . . . . .	$\text{H}_2(\text{Mg}, \text{Fe})_4\text{Al}_3\text{Si}_{10}\text{O}_{37}$ , altered	Ortho.
618. Barysilite . . . . .	$\text{Pb}_2\text{Si}_2\text{O}_7$	
619. Ganomalite . . . . .	$\text{Pb}_2\text{Si}_2\text{O}_7 \cdot (\text{Ca}, \text{Mn})_2\text{SiO}_4$	Tetrag.
620. Hyalotekite . . . . .	$(\text{Pb}, \text{Ba}, \text{Ca})_2\text{B}_2(\text{SiO}_3)_{12}$	
621. Barylite . . . . .	$\text{Ba}_4\text{Al}_4\text{Si}_7\text{O}_{24}$	
622. Roebbingite . . . . .	$5(\text{H}_2, \text{CaSiO}_3) \cdot 2(\text{CaPbSO}_4)$	
<b>c. Orthosilicates</b>		
623. Nephelite . . . . .	$\text{K}_2\text{Na}_6\text{Al}_6\text{Si}_6\text{O}_{24}$	Hexag.
624. Elaeolite . . . . .	$\text{K}_2\text{Na}_6\text{Al}_6\text{Si}_6\text{O}_{24}$	Hexag.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
581.	Green	5.5	3	Norway	Rock forming
582.	Deep green	5.5	3	Tunkinsk Mts.	
583.	Black	5.5	3	Vesuvius	
584.	Black	5.5	3	Italy	
585.	Brown	5	3	North Greenland	
586.	Brown	5	3	Ontario	
587.	Blue	6	3	California	
588.	Blue	6	3	Corsica	
588a.	Black	6	3	Ireland	
589.	Blue	4	3	Rhode Island	
590.	Blue	4	3	Scotland	Gems
590a.	Black	6	3	Colorado	
591.	Black	6	3	California	
592.	Black	6	3	Southern Norway	
593.	Black	.....	3	Southern Norway	
594.	Green	7.5	2.6	E. United States	
595.	Green	7.5	2.6	E. United States	
596.	Green	7.5	2.6	E. United States	
597.	Green	7.5	2.6	Scotland	
598.	Red	5	2.9	Western Greenland	
599.	Red	5	2.9	Norway	Rock forming
600.	.....	.....	2.5	South Greenland	
601.	Yellow	6	2.8	Norway	
602.	Brown	6	4.4	Norway	
603.	Brown	6	4	Norway	
604.	Brown	6	4	Norway	
605.	Brown	4	3	Greenland	
606.	Brown	5	4	Norway	
607.	Green	4	2.9	Norway	
608.	Yellow	5	3	Norway	Rock forming
609.	Blue	7	2.6	Connecticut	
610.	Blue	7	2.6	Finland	
611.	Various	7	2.6	Sweden	
612.	Various	7	2.6	Helsingfors	
613.	Various	7	2.6	Norway	
614.	Various	7	2.6	Finland	
615.	Various	7	2.6	Maine	
616.	Various	7	2.6	Kragero	
617.	Various	7	2.6	Kragero	
618.	White	3	6	Sweden	Rock forming
619.	Colorless	3	5.7	Sweden	
620.	White	5	3.8	Sweden	
621.	Colorless	7	4	Sweden	
622.	White	3	3	New Jersey	
623.	Colorless	5	2.5	Vesuvius	
624.	Brown	5	2.5	Maine	

	Composition	Form
<b>VII. SILICATES—continued</b>		
625. Gieseckite.....	$K_2Na_4Al_8Si_6O_{24} \cdot nH_2O$	Pseudo.
626. Eucryptite.....	$LiAlSiO_4$	Hexag.
627. Kaliophilite.....	$KAlSiO_4$	Hexag.
628. Cancrinite.....	$H_6Na_6Ca(NaCO_3)_2Al_8(SiO_4)_9$	Hexag.
629. Microsommite.....	$(Na,K)_{10}Ca_4Al_{12}Si_{12}O_{52}SCl_4$	Hexag.
630. Sodalite.....	$Na_4(AlCl)Al_3(SiO_4)_3$	Regular
631. Häthynite.....	$Na_2Ca_2(NaSO_4 \cdot Al)Al_2(SiO_4)_3$	Regular
632. Noselite.....	$Na_4(NaSO_4 \cdot Al)Al_2(SiO_4)_3$	Regular
633. Lazurite.....	$Na_4(NaS_3 \cdot Al)Al_2(SiO_4)_3$	Regular
634. Helvite.....	$(Mn,Fe)_2(Mn,S)Be_2SiO_4$	Regular
635. Danalite.....	$(Fe,Zn,Mn)_2[(Zn,Fe)_2S]Be_2(SiO_4)_3$	Regular
636. Eulytite.....	$Bi_4(SiO_4)_3$	Regular
637. Zunyite.....	$(Al(OH,F,Cl)_2)_6Al_2(Si_6O_4)_3$	Regular
<b>II III</b>		
638. Garnet.....	$R_3R_4(SiO_4)_3$	Regular
639. Grossularite.....	$Ca_3Al_2(SiO_4)_3$	Regular
640. Cinnamon-stone.....	$Ca_3Al_2(SiO_4)_3$	Regular
641. Hyacinth.....	$Ca_3Al_2(SiO_4)_3$	Regular
642. Succinite.....	$Ca_3Al_2(SiO_4)_3$	Regular
643. Romanzovite.....	$Ca_3Al_2(SiO_4)_3$	Regular
644. Pyrope.....	$Mg_3Al_2(SiO_4)_3$	Regular
645. Rhodolite.....	$Mg_3Al_2(SiO_4)_3$	Regular
646. Almandite.....	$Fe_3Al_2(SiO_4)_3$	Regular
647. Spessartite.....	$Mn_3Al_2(SiO_4)_3$	Regular
648. Andradite.....	$Ca_3Fe_2(SiO_4)_3$	Regular
649. Topazolite.....	$Ca_3Fe_2(SiO_4)_3$	Regular
650. Demantoid.....	$Ca_3Fe_2(SiO_4)_3$	Regular
651. Colophonite.....	$Ca_3Fe_2(SiO_4)_3$	Regular
652. Melanite.....	$Ca_3Fe_2(SiO_4)_3$	Regular
653. Pyreneite.....	$Ca_3Fe_2(SiO_4)_3$	Regular
654. Rothoffite.....	$(CaMg)_3Fe_2(SiO_4)_3$	Regular
655. Allochroite.....	$(Mg,Ca)_3Fe_2(SiO_4)_3$	Regular
656. Polyadelphite.....	$(Mg,Ca)_3Fe_2(SiO_4)_3$	Regular
657. Bredbergite.....	$(Mg,Ca)_3Fe_2(SiO_4)_3$	Regular
658. Aplome.....	$(Mg,Ca)_3Fe_2(SiO_4)_3$	Regular
659. Titaniferous garnet.....	$3CaO \cdot (Fe,Ti,Al)_2O_3 \cdot 3(Si,Ti)O_2$	Regular
660. Yttergranat.....	$3CaO \cdot (Fe,Ti,Y,Al)YO_3$	Regular
661. Uvarovite.....	$Ca_3Cr_2(SiO_4)_3$	Regular
662. Schorlomite.....	$Ca_3(FeTi)_2(SiTi)O_4$	Regular
663. Partschinite.....	$(Mn,Fe)_3Al_2Si_3O_{12}$	Mono.
664. Agricolite.....	$Bi_4Si_3O_{12}$	Mono.
665. Chrysolite.....	$(Mg,Fe)_2SiO_4$	Ortho.
666. Olivine.....	$(Mg,Fe)_2SiO_4$	Ortho.
667. Hyalosiderite.....	$(Mg,Fe)_2SiO_4 + Fe$	Ortho.
668. Iddingsite.....	$(Ca,Mg,Fe)_2SiO_4$	Ortho.
669. Monticellite.....	$CaMgSiO_4$	Ortho.
670. Forsterite.....	$Mg_2SiO_4$	Ortho.
671. Hortonolite.....	$(Fe,Mg,Mn)_2SiO_4$	Ortho.

## LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
625.	Brown	5	2.6	New York	Rock forming
626.	Colorless	5	2.6	Connecticut	
627.	Colorless	6	2	Mt. Somma	
628.	Gray	5	2	Maine	
629.	Colorless	5	2	Vesuvius	
630.	Gray	5	2	Maine	
631.	Blue	5.5	2	Vesuvius	
632.	Grayish	5.5	2	Andernach	
633.	Blue	5	2	Chile	
634.	Yellow	6	3	Virginia	Ornaments
635.	Red	5.5	3	Colorado	
636.	Brown	4.5	6	Saxony	
637.	Brown	7	2.8	Colorado	
638.	Red	6.5	3	Mountains	Rock forming
639.	Pale green	6.5	3.5	Ceylon	
640.	Brown	6.5	3.5	Ceylon	
641.	Brown	6.5	3.5	Ceylon	
642.	Yellow	6.5	3.5	Piedmont	
643.	Brown	6.5	3.5	Russia	
644.	Red	6.5	3.7	Bohemia	
645.	Red	6.5	3.7	North Carolina	
646.	Red	6.5	3.9	Pennsylvania	
647.	Red	6.5	4	Colorado	
648.	Yellow	6.5	3.8	Portugal	Rock forming and gems
649.	Green	6.5	3.8	France	
650.	Green	6.5	3.8	Mountains	
651.	Brown	6.5	3.8	Mountains	
652.	Black	6.5	3.8	Mountains	
653.	Black	6.5	3.8	Mountains	
654.	Brown	6.5	3.8	Mountains	
655.	Brown	6.5	3.8	Mountains	
656.	Yellowish brown	6.5	3.8	New Jersey	
657.	Yellowish brown	6.5	3.7	Sala	
658.	Brown	6.5	3.7	Siberia	Rock forming
659.	Black	6.5	3.7	Siberia	
660.	Black	6.5	3.7	Norway	
661.	Green	7.5	3	Canada	
662.	Black	7	3.8	Arkansas	Bismuth
663.	Yellow	6.5	4	Transylvania	
664.	Yellow	2	.....	Johanngeorgen.	
665.	Green	6.5	3	Virginia	
666.	Green	6.5	3	Virginia	Rock forming
667.	Green	6.5	3	Baden	
668.	Brown	.....	2.8	California	
669.	Gray	5	3	Arkansas	
670.	White	6	3	Vesuvius	
671.	Yellow	6	3.9	New York	

	Composition	Form
<b>VII. SILICATES—continued</b>		
672. Fayalite.....	$\text{Fe}_2\text{SiO}_4$	Ortho.
673. Knebelite.....	$(\text{Fe}, \text{Mn})_2\text{SiO}_4$	Ortho.
674. Tephroite.....	$\text{Mn}_2\text{SiO}_4$	Ortho.
675. Willemite.....	$\text{Zn}_2\text{SiO}_4$	Hexag.
676. Phenacite.....	$\text{Be}_2\text{SiO}_4$	Hexag.
677. Trimerite.....	$(\text{Mn}, \text{Ca})_3\text{SiO}_4 \cdot \text{Be}_2\text{SiO}_4$	Triclinic.
678. Dioptase.....	$\text{H}_2\text{CuSiO}_4$	Hexag.
679. Friedelite.....	$\text{H}_7(\text{MnCl})\text{Mn}_2\text{Si}_4\text{O}_{16}$	Hexag.
680. Pyrosmalite.....	$\text{H}_7[(\text{Fe}, \text{MnCl})(\text{Fe}, \text{Mn})_4\text{Si}_4\text{O}_{16}]$	Hexag.
681. Meionite.....	$\text{Ca}_2\text{Al}_6\text{Si}_6\text{O}_{28}$	Tetrag.
682. Wernerite.....	*Me, Ma <sub>2</sub>	Tetrag.
683. Passauite.....	*Me, Ma <sub>2</sub> or Ma <sub>3</sub>	Tetrag.
684. Glaucolite.....	*Me, Ma <sub>2</sub> or Ma <sub>3</sub>	Tetrag.
685. Mizzonite.....	Me, Ma <sub>3</sub>	Tetrag.
686. Dipyre.....	Me, Ma <sub>3</sub>	Tetrag.
687. Cousseranite.....	Me, Ma <sub>3</sub>	Tetrag.
688. Marialite.....	$\text{Na}_4\text{Al}_3\text{Si}_3\text{O}_{24}\text{Cl}$	Tetrag.
689. Sarcosite.....	$\text{Ca}_2\text{Na}_2\text{Al}_6\text{Si}_6\text{O}_{28}$	Tetrag.
690. Melilite.....	$\text{Na}_2(\text{Ca}, \text{Mg})_{11}(\text{Al}, \text{Fe})_4(\text{SiO}_4)_9$	Tetrag.
691. Humboldtillite.....	$\text{Na}_2(\text{Ca}, \text{Mg})_{11}(\text{Al}, \text{Fe})_4(\text{SiO}_4)_9$	Tetrag.
692. Gehlenite.....	$\text{Ca}_2\text{Al}_2\text{Si}_2\text{O}_{10}$	Tetrag.
693. Vesuvianite.....	$\text{H}_4\text{Ca}_{12}(\text{Al}, \text{Fe})_6\text{Si}_{10}\text{O}_{43}$	Tetrag.
694. Cyprine.....	$\text{H}_4\text{Ca}_{12}(\text{Al}, \text{Fe})_6\text{Si}_{10}\text{O}_{43}$	Tetrag.
695. Zircon.....	$\text{ZrSiO}_4$	Tetrag.
696. Hyacinth.....	$\text{ZrSiO}_4$	Tetrag.
697. Jargon.....	$\text{ZrSiO}_4$	Tetrag.
698. Thorite.....	$\text{ThSiO}_4$	Tetrag.
699. Auerlite.....	$\text{ThSiO}_4$	Tetrag.
700. Danburite.....	$\text{CaB}_2(\text{SiO}_4)_2$	Ortho.
701. Topaz.....	$\text{Al}_2(\text{F} \cdot \text{OH})_2\text{SiO}_4$	Ortho.
702. Physalite.....	$\text{Al}_2(\text{F} \cdot \text{OH})_2\text{SiO}_4$	Ortho.
703. Pyonite.....	$\text{Al}_2(\text{F} \cdot \text{OH})_2\text{SiO}_4$	Ortho.
704. Andalusite.....	$\text{Al}_2\text{SiO}_5$	Ortho.
705. Chiasolite.....	$\text{Al}_2\text{SiO}_5$	Ortho.
706. Sillimanite.....	$\text{Al}_2\text{SiO}_5$	Ortho.
707. Cyanite.....	$\text{Al}_2\text{SiO}_5$	Triclinic
708. Datolite.....	$\text{HCaBSiO}_5$	Mono.
709. Homilite.....	$(\text{Ca}, \text{Fe})_2\text{B}_2\text{Si}_2\text{O}_{10}$	Mono.
710. Euclase.....	$\text{HBeAlSiO}_5$	Mono.
711. Gadolinite.....	$\text{Be}_2\text{FeY}_2\text{Si}_2\text{O}_{10}$	Mono.
712. Yttrialite.....	$(\text{ThY})_2\text{O}_3 \cdot 2\text{SiO}_2$	Amorph.
713. Rowlandite.....	Y, Fe, U, Ca, silicate	Massive
714. Mackintoshite.....	T, Ce, U, silicate	Massive
715. Zoisite.....	$\text{HCa}_2\text{Al}_2\text{Si}_2\text{O}_{11}$	Ortho.
716. Thulite.....	$\text{HCa}_2\text{Al}_2\text{Si}_2\text{O}_{11}$	Ortho.
717. Epidote.....	$\text{HCa}_2(\text{Al}, \text{Fe})_2\text{Si}_2\text{O}_{11}$	Mono.
718. Scorza.....	$\text{HCa}_2(\text{Al}, \text{Fe})_2\text{Si}_2\text{O}_{11}$	Sand
719. Thallite.....	$\text{HCa}_2(\text{Al}, \text{Fe})_2\text{Si}_2\text{O}_{11}$	Mono.

\*Me = Meionite; Ma = Marialite.

## LIST OF MINERALS

No.	Color	Hardness	Gravity	Locality	Chief Constituent or Use
672.	Brown	6	4	Yellowstone Park	Rock forming
673.	Brown	6	4	Sweden	
674.	Red	5.5	4	New Jersey	
675.	White	5.5	3.8	New Jersey	Zinc
676.	Colorless	7.5	2.9	Colorado	
677.	Pink	6	3	Sweden	Gems
678.	Green	5	3	Arizona	
679.	Red	4	3	Pyrenees	Rock forming
680.	Gray	4	3	Sweden	
681.	Colorless	5.5	2.7	Vesuvius	
682.	White	5	2.6	Finland	
683.	Yellowish	5	2.6	Bavaria	
684.	Gray	5	2.6	Siberia	
685.	White	5	2.6	Vesuvius	
686.	White	5	2.6	Norway	
687.	White	5	2.6	Pyrenees	
688.	White	5	2.6	Naples	
689.	Red	6	2.5	Vesuvius	
690.	White	5	2.9	Vesuvius	
691.	Yellow	5	2.9	Vesuvius	
692.	Green	5	2.9	Tyrol	
693.	Brown	6.5	3	California	
694.	Blue	6.5	3	Norway	
695.	Yellow	7.5	4.6	New York	
696.	Red	7.5	4.6	Canada	
697.	Smoky	7.5	4.6	Ceylon	
698.	Black	4.5	5	Norway	Gems
699.	Orange	2.5	4	North Carolina	
700.	Yellow	7	2.9	Connecticut	
701.	Yellow	8	3	Urals	
702.	Yellow	8	3	Finbo	
703.	Yellow	8	3	Saxony	
704.	Red	7.5	3	E. United States	
705.	Brown	7	3	Maine	
706.	Brown	6	3	E. United States	
707.	Blue	5	3.5	E. United States	
708.	White	5	2.9	New Jersey	Rock forming
709.	Black	5	3	Norway	
710.	Colorless	7.5	3	Brazil	
711.	Black	6.5	4	Texas	
712.	Green	5	4.5	Texas	
713.	Drab green	.....	4.5	Texas	
714.	Black	.....	.....	Texas	
715.	Gray	6	3	Carinthia	
716.	Red	6	3	Carinthia	
717.	Green	6	3	Michigan	
718.	Green	6	3	Transylvania	
719.	Yellow	6	3	Bourg d'Oisans	



	Composition	Form
<b>VII. SILICATES—continued</b>		
720. Bucklandite.....	$\text{HCa}_2(\text{Al,Fe})_2\text{Si}_3\text{O}_{13}$	Mono.
721. Withamite.....	$\text{HCa}_2(\text{Al,Fe})_2\text{Si}_3\text{O}_{13}$	Mono.
722. Clinzoisite.....	$\text{HCa}_2(\text{Al,Fe,Mn})_2\text{Si}_3\text{O}_{13}$	Mono.
723. Picroepidote.....	$\text{HCa}_2(\text{Al,Fe,Mn})_2\text{Si}_3\text{O}_{13}$	Mono.
724. Piedmonite.....	$\text{HCa}_2(\text{Al,Fe,Mn})_2\text{Si}_3\text{O}_{13}$	Mono.
725. Allanite.....	$\text{Al,Fe,Mn,Ca,Na,K,Mg,Er,Y,La,Di,Ce, Th, silicate}$	Mono.
726. Bagrationite.....	$\text{Al,Fe,Mn,Ca,Na,K,Mg,Er,Y,La,Di,Ce, Th, silicate}$	Mono.
727. Axinite.....	$\text{H}_2(\text{Ca,Mn})_4(\text{BO})\text{Al}_3(\text{SiO}_3)_3$	Triclinic
728. Prehnite.....	$\text{H}_2\text{Ca}_2\text{Al}_2(\text{SiO}_3)_3$	Ortho.
729. Harstigitite.....	$\text{H}_7(\text{Ca,Mn})_{13}\text{Al}_3\text{Si}_{10}\text{O}_{40}$	Ortho.
730. Cuspidine.....	$\text{Ca}_2\text{Si}(\text{O,F})_4$	Mono.
<i>d. Subsilicates</i>		
731. Chondrodite.....	$\text{H}_2(\text{Mg,Fe})_{10}\text{Si}_2\text{O}_{14}\text{F}_4$	Mono.
732. Humite.....	$\text{H}_2(\text{Mg,Fe})_{10}\text{Si}_2\text{O}_{14}\text{F}_4$	Ortho.
733. Clinohumite.....	$\text{H}_2(\text{Mg,Fe})_{10}\text{Si}_2\text{O}_{14}\text{F}_4$	Mono.
734. Ilvaite.....	$\text{CaFe}_2(\text{FeOH})(\text{SiO}_3)_2$	Ortho.
735. Ardenite.....	$\text{H}_3\text{Mn}_2\text{Al}_2\text{VSi}_2\text{O}_{13}$	Ortho.
736. Långbanite.....	$37\text{Mn}_2\text{SiO}_7 \cdot 10\text{Fe}_3\text{Sb}_2\text{O}_8$	Hexag.
737. Kentrolite.....	$\text{Pb}_2\text{Mn}_2\text{Si}_2\text{O}_9$	Ortho.
738. Melanotekite.....	$\text{Pb}_2\text{Fe}_2\text{Si}_2\text{O}_9$	Ortho.
739. Bertrandite.....	$\text{H}_2\text{Be}_2\text{Si}_2\text{O}_9$	Ortho.
740. Calamine.....	$\text{H}_2\text{Zn}_2\text{SiO}_5$	Ortho.
741. Clinohedrite.....	$\text{H}_2\text{CaZnSiO}_5$	Mono.
742. Carpholite.....	$\text{H}_4\text{MnAl}_2\text{Si}_2\text{O}_{10}$	Mono.
743. Lawsonite.....	$\text{H}_4\text{CaAl}_2\text{Si}_2\text{O}_{10}$	Ortho.
744. Cerite.....	$\text{Ce}_2(\text{OH})_3\text{CeO} \cdot \text{CaFe}(\text{SiO}_3)_3$	Ortho.
745. Tourmaline.....	$\text{Fe}_2\text{Na}_2\text{B}_6\text{Al}_{14}\text{H}_8\text{Si}_{12}\text{O}_{63}$	Hexag.
746. Indicolite.....	$\text{Fe}_2\text{Na}_2\text{B}_6\text{Al}_{14}\text{H}_8\text{Si}_{12}\text{O}_{63}$	Hexag.
747. Aphrizite.....	$\text{Fe}_2\text{Na}_2\text{B}_6\text{Al}_{14}\text{H}_8\text{Si}_{12}\text{O}_{63}$	Hexag.
748. Achroite.....	$(\text{Fe}_2\text{Na}_2\text{B}_6\text{Al}_{14}\text{H}_8\text{Si}_{12}\text{O}_{63}), \text{etc.}$	Hexag.
749. Dumortierite.....	$4\text{Al}_2\text{O}_3 \cdot 3\text{SiO}_2$	Ortho.
750. Staurolite.....	$\text{HFeAl}_2\text{Si}_2\text{O}_{13}$	Ortho.
751. Nordmarkite.....	$\text{HFeAl}_2\text{Si}_2\text{O}_{13} + \text{Mg}$	Ortho.
752. Kornerupine.....	$\text{MgAl}_2\text{SiO}_6$	Ortho.
753. Sapphirine.....	$\text{Mg}_5\text{Al}_{12}\text{Si}_3\text{O}_{27}$	Mono.
<i>2. Hydrous</i>		
<i>a. Zeolites</i>		
754. Inesite.....	$2(\text{Mn,Ca})\text{SiO}_3 \cdot \text{H}_2\text{O}$	Triclinic
755. Ganophyllite.....	$\text{Mn}_2\text{Al}_2\text{Si}_2\text{O}_{10} \cdot 6\text{H}_2\text{O}$	Mono.
756. Okenite.....	$\text{H}_2\text{CaSi}_2\text{O}_6 \cdot \text{H}_2\text{O}$	Ortho.
757. Grolite.....	$\text{H}_2\text{Ca}_2(\text{SiO}_3)_3 \cdot \text{H}_2\text{O}$	Ortho.
758. Apophyllite.....	$\text{H}_2\text{KCa}_4(\text{SiO}_3)_8 \cdot 4\frac{1}{2}\text{H}_2\text{O}$	Tetrag.
759. Ptilolite.....	$(\text{CaK}_2\text{Na}_2)\text{Al}_2\text{Si}_2\text{O}_{10} \cdot 5\text{H}_2\text{O}$	Mono.
760. Mordenite.....	$(\text{CaK}_2\text{Na}_2)\text{Al}_2\text{Si}_2\text{O}_{10} \cdot 6\text{H}_2\text{O}$	Mono.

## LIST OF MINERALS

No.	Color	Hardness	Gravity	Locality	Chief Constituent or Use
720.	Black	6	3	Bourg d'Oisans	Rock forming
721.	Red	6	3	Scotland	
722.	Rose red	.....	.....	Ceylon	
723.	Yellowish	6	3	Siberia	
724.	Brown	6.5	3	Pennsylvania	
725.	Brown	5.5	3	Massachusetts	
726.	Black	6	3.8	Massachusetts	
727.	Brown	6.5	3	Maine	
728.	Green	6	2.8	Connecticut	
729.	Colorless	5.5	3	Sweden	
730.	Red	5	2.8	Mt. Somma	
731.	Yellow	6	3	Mt. Somma	Rock forming
732.	Yellow	6	3	New York	
733.	Yellow	6	3	New York	
734.	Iron black	5.5	3.9	Elba	
735.	Yellow	6.7	3.6	Belgium	
736.	Black	6.5	4.9	Sweden	
737.	Brown	5	6	Chile	
738.	Black	6.5	5.7	New Mexico	
739.	Colorless	6	2.5	Colorado	
740.	White	4.5	3	New Jersey	Zinc
741.	Colorless	5.5	3	New Jersey	
742.	Yellow	5	2.9	Hartz	
743.	Colorless	.....	3	California	
744.	Gray	5.5	4.8	Sweden	
745.	Black	7	2.9	Maine	
746.	Blue	7	2.9	Maine	
747.	Black	7	2.9	Norway	
748.	Colorless	7	2.9	Elba	
749.	Blue	7	3	Arizona	Rock forming and gems
750.	Brown	7	3.6	New Hampshire	
751.	Brown	7	3.6	Sweden	
752.	Colorless	6.5	3	Greenland	
753.	Green	7.5	3	Greenland	
754.	Red	6	3	Germany	Rock forming
755.	Brown	4	2.8	Sweden	
756.	White	4.5	2	Iceland	
757.	White	.....	.....	California	
758.	White	4.5	2	New Jersey	
759.	Colorless	.....	.....	Colorado	
760.	White	3	2	Wyoming	

	Composition	Form
<b>VII. SILICATES—continued</b>		
761. Heulandite.....	$H_4CaAl_6(SiO_3)_6 \cdot 3H_2O$	Mono.
762. Brewsterite.....	$H_4(Sr,Ba,Ca)Al_6Si_6O_{18} \cdot 3H_2O$	Mono.
763. Epistilbite.....	$H_4CaAl_6Si_6O_{18} \cdot 3H_2O$	Mono.
764. Wellsite.....	$(Ba,Ca,K_2)Al_6Si_6O_{18} \cdot 3H_2O$	Mono.
765. Phillipsite.....	$(K,Ca)Al_6Si_6O_{18} \cdot 4\frac{1}{2}H_2O$	Mono.
766. Harmotome.....	$H_2(K,Ba)Al_2Si_2O_7 \cdot 4H_2O$	Mono.
767. Stilbite.....	$H_4(Na_2,Ca)Al_6Si_6O_{18} \cdot 4H_2O$	Mono.
768. Gismondite.....	$CaAl_2Si_2O_7 \cdot 4H_2O$	Mono.
769. Laumontite.....	$H_4CaAl_2Si_2O_{14} \cdot 2H_2O$	Mono.
770. Leonhardite.....	$H_4CaAl_2Si_2O_{14} \cdot 2H_2O$	Mono.
771. Schneiderite.....	$H_4CaAl_2Si_2O_{14} \cdot 2H_2O$	Mono.
772. Laubanite.....	$Ca_2Al_2Si_2O_{15} + 6H_2O$	Mono.
773. Chabazite.....	$(Ca,Na_2)Al_2Si_2O_{13} \cdot 6H_2O$	Hexag.
774. Acadialite.....	$(Ca,Na_2)Al_2Si_2O_{13} \cdot 6H_2O$	Hexag.
775. Haydenite.....	$(Ca,Na_2)Al_2Si_2O_{13} \cdot 6H_2O$	Hexag.
776. Phacolite.....	$(Ca,Na_2)Al_2Si_2O_{13} \cdot 6H_2O$	Hexag.
777. Herschelite.....	$(Ca,Na_2)Al_2Si_2O_{13} \cdot 6H_2O$	Hexag.
778. Gmelinite.....	$(Na_2,Ca)Al_2Si_2O_{13} \cdot 6H_2O$	Hexag.
779. Levynite.....	$CaAl_2Si_2O_{10} \cdot 5H_2O$	Hexag.
780. Offretite.....	$(K,Ca)_2Al_2Si_2O_{10} \cdot 17H_2O$	Hexag.
781. Analcite.....	$NaAlSi_3O_8 \cdot H_2O$	Regular
782. Analcime.....	$NaAlSi_3O_8 \cdot H_2O$	Regular
783. Edingtonite.....	$BaAl_2Si_2O_{10} \cdot 3H_2O$	Tetrag.
784. Natrolite.....	$Na_2Al_2Si_2O_{10} \cdot 2H_2O$	Ortho.
785. Bergmannite.....	$Na_2Al_2Si_2O_{10} \cdot 2H_2O$	Ortho.
786. Scolecite.....	$CaAl_2Si_2O_{10} \cdot 3H_2O$	Mono.
787. Mesolite.....	$CaAl_2Si_2O_{10} \cdot 3H_2O + Na_2Al_2Si_2O_{10} \cdot 2H_2O$	Mono.
788. Thomsonite.....	$(Na_2,Ca)Al_2Si_2O_{10} \cdot 2\frac{1}{2}H_2O$	Ortho.
789. Ozarkite.....	$(Na_2,Ca)Al_2Si_2O_{10} \cdot 2\frac{1}{2}H_2O$	Ortho.
790. Hydronephelite.....	$HNa_2Al_2Si_2O_{10} \cdot 3H_2O$	Ortho.
<b>b. Micas</b>		
791. Muscovite.....	$H_3KAl_3Si_3O_{12}$	Mono.
792. Damourite.....	$H_3KAl_3Si_3O_{12}$	Mono.
793. Margarodite.....	$H_3KAl_3Si_3O_{12}$	Mono.
794. Gilbertite.....	$H_3KAl_3Si_3O_{12}$	Mono.
795. Sericite.....	$H_3KAl_3Si_3O_{12}$	Scaly
796. Fuchsite.....	$H_3KAl_3Si_3O_{12} + Cr$	Scaly
797. Pinite.....	$H_3KAl_3Si_3O_{12}$	Amorph.
798. Paragonite.....	$H_3NaAl_3(Si_3O_{12})_3$	Mono.
799. Lepidolite.....	$(Li,K,Na)_2(Al,Fe)OH_2(SiO_3)_3$	Mono.
800. Zinnwaldite.....	$H_3KLiFe_3Al_3Si_3O_{12}$	Mono.
801. Biotite.....	$(H,K)_2(Mg,Fe)_4(Al,Fe)_2(SiO_4)_4$	Mono.
802. Merxene.....	$(H,K)_2(Mg,Fe)_4(Al,Fe)_2(SiO_4)_4 + Fe$	Mono.
803. Anomite.....	$(H,K)_2(Mg,Fe)_4(Al,Fe)_2(SiO_4)_4 + Mn$	Mono.
804. Haughtonite.....	$(H,K)_2(Mg,Fe)_4(Al,Fe)_2(SiO_4)_4 + Mn$	Mono.
805. Manganophyllite.....	$(H,K)_2(Mg,Fe)_4(Al,Fe)_2(SiO_4)_4 + Mn$	Mono.
806. Caswellite.....	$(H,K)_2(Mg,Fe)_4(Al,Fe)_2(SiO_4)_4 + Fe$	Mono.
807. Phlogopite.....	$(H,K)_2(Mg,Fe)_4(Al,Fe)_2(SiO_4)_4 + Fe$	Mono.

## LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
761.	White	3.5	2	New Jersey	Rock forming
762.	White	5	2	Strontian	
763.	White	4	2	Nova Scotia	
764.	Colorless	4	2	North Carolina	
765.	White	4	2	Ireland	
766.	White	4.5	2	New York	
767.	White	3.5	2	New York	
768.	Colorless	4.5	2	Mt. Albano	
769.	White	3.5	2	New Jersey	
770.	White	3.5	2	Mountains	
771.	White	3.5	2	Italy	
772.	White	4.5	2	Silesia	
773.	White	4	2	New Jersey	
774.	Red	4	2	Nova Scotia	
775.	Yellow	4	2	Maryland	
776.	Colorless	4	2	Bohemia	
777.	Colorless	4	2	Sicily	
778.	White	4.5	2	New Jersey	
779.	White	4	2	Colorado	
780.	Colorless	.....	2	France	
781.	Colorless	5	2	New Jersey	Electrical purposes and rock forming
782.	Colorless	5	1.9	New Jersey	
783.	White	4	2.6	Scotland	
784.	White	5	2	New Jersey	
785.	White	5	2	Southern Norway	
786.	White	5	2	Colorado	
787.	White	5	2	Colorado	
788.	White	5	2	Colorado	
789.	White	.....	2	Arkansas	
790.	White	4.5	2	Maine	
791.	Colorless	2	2.7	Maine	
792.	Colorless	2	2.7	Maine	
793.	Pearly	2	2.7	Tyrol	
794.	Whitish	2	2.7	Cornwall	
795.	Whitish	2	2.7	Wiesbaden	
796.	Green	2	2.7	Zillerthal	
797.	Gray	2.5	2.6	Germany	
798.	Yellow	2.5	2.7	Pennsylvania	
799.	Red	2.5	2.8	Maine	
800.	Yellow	2.5	2.8	Zinnwald	
801.	Green	2.5	2.7	N. England States	
802.	Dark	2.5	2.7	Vesuvius	
803.	Dark	.....	.....	New Jersey	
804.	.....	.....	2.9	Sutherland	
805.	Red	2.5	2.7	Sweden	
806.	.....	.....	.....	New Jersey	
807.	Brown	2.5	2.7	New York	

	Composition	Form
VII. SILICATES— <i>continued</i>		
808. Lepidomelane. . . . .	$(\text{H}, \text{K})_3\text{Fe}_3(\text{Fe}, \text{Al})_4(\text{SiO}_4)_5$	Mono.
809. Alurgite. . . . .	$(\text{H}, \text{K})_2\text{Fe}_2(\text{Fe}, \text{Al})_4(\text{SiO}_4)_5 + \text{Mn}$	Mono.
810. Roscoelite. . . . .	$\text{H}_8\text{K}(\text{Mg}, \text{Fe})(\text{Al}, \text{V})_4(\text{SiO}_3)_{12}$	Mono.
811. Margarite. . . . .	$\text{H}_5\text{CaAl}_4\text{Si}_6\text{O}_{12}$	Mono.
812. Seybertite. . . . .	$\text{H}_3(\text{Mg}, \text{Ca})_5\text{Al}_2\text{Si}_2\text{O}_{16}$	Mono.
813. Zanthophyllite. . . . .	$\text{H}_8(\text{Mg}, \text{Ca})_{14}\text{Al}_{16}\text{Si}_5\text{O}_{52}$	Mono.
814. Chloritoid. . . . .	$\text{H}_2(\text{Fe}, \text{Mg})\text{Al}_2\text{SiO}_7$	Mono.
815. Sismondine. . . . .	$\text{H}_{14}\text{Fe}_7\text{Al}_{16}\text{Si}_8\text{O}_{54}$	Triclinic
816. Salmite. . . . .	$\text{H}_{14}\text{Fe}_7\text{Al}_{16}\text{Si}_8\text{O}_{54}$	Triclinic
817. Masonite. . . . .	$\text{H}_{14}\text{Fe}_7\text{Al}_{16}\text{Si}_8\text{O}_{54}$	Triclinic
818. Ottrelite. . . . .	$\text{H}_2(\text{Fe}, \text{Mn})\text{Al}_2\text{Si}_2\text{O}_9$	Mono.
819. Venasquite. . . . .	$\text{H}_2\text{FeAl}_2\text{Si}_2\text{O}_{11}$	Mono.
820. Phyllite. . . . .	$\text{H}_2\text{FeAl}_2\text{Si}_2\text{O}_{11}$	Mono.
821. Clinocllore. . . . .	$\text{H}_8(\text{Mg}, \text{Fe})_5\text{Al}_2\text{Si}_3\text{O}_{18}$	Mono.
822. Leuchtenbergite. . . . .	$\text{H}_8(\text{Mg}, \text{Fe})_5\text{Al}_2\text{Si}_3\text{O}_{18}$ , lacks Fe	Mono.
823. Kotschubeite. . . . .	$\text{H}_8(\text{Mg}, \text{Fe})_5\text{Al}_2\text{Si}_3\text{O}_{18} + \text{Cr}$	Hexag.
824. Manganchlorite. . . . .	$\text{H}_8(\text{Mg}, \text{Fe})_5\text{Al}_2\text{Si}_3\text{O}_{18} + \text{Mn}$	Hexag.
825. Penninite. . . . .	$\text{H}_8(\text{Mg}, \text{Fe})_5\text{Al}_2\text{Si}_3\text{O}_{18}$	Mono.
826. Kämmererite. . . . .	$\text{H}_8(\text{Mg}, \text{Fe})_5\text{Al}_2\text{Si}_3\text{O}_{18}$	Mono.
827. Pseudophite. . . . .	$\text{H}_8(\text{Mg}, \text{Fe})_5\text{Al}_2\text{Si}_3\text{O}_{18}$	Massive
828. Prochlorite. . . . .	$\text{H}_{40}(\text{Fe}, \text{Mg})_{23}\text{Al}_{14}\text{Si}_{13}\text{O}_{90}$	Mono.
829. Corundophilite. . . . .	$\text{H}_{30}\text{Mg}_{11}\text{Al}_6\text{Si}_6\text{O}_{45}$	Mono.
830. Amesite. . . . .	$\text{H}_4(\text{Mg}, \text{Fe})_3\text{Al}_2\text{SiO}_9$	Hexag.
831. Aphrosiderite. . . . .	$\text{H}_{10}\text{Fe}_6(\text{Fe}, \text{Al})_4\text{Si}_4\text{O}_{25}$	Hexag.
832. Diabantite. . . . .	$\text{H}_{18}(\text{Fe}, \text{Mg})_{12}\text{Al}_4\text{Si}_9\text{O}_{45}$	Hexag.
833. Delessite. . . . .	$\text{H}_{10}(\text{Mg}, \text{Fe})_4(\text{Al}, \text{Fe})_4\text{Si}_4\text{O}_{43}$	Hexag.
834. Epichlorite. . . . .	$\text{H}_7(\text{Mg}, \text{K})_7(\text{Al}, \text{Fe})_8\text{Si}_4\text{O}_{18}$	Hexag.
835. Euralite. . . . .	$\text{H}_{16}(\text{Mg}, \text{Fe}, \text{Ca})_9(\text{Al}, \text{Fe})_4\text{Si}_9\text{O}_{37}$	Amorph.
836. Chlorophaeite. . . . .	Fe, Mg, Mn, Ca, silicate	Amorph.
837. Hullite. . . . .	Fe, Mg, Mn, Ca, silicate	Massive
838. Cronstedite. . . . .	$\text{H}_6(\text{Fe}, \text{Mg})_3\text{Al}_2\text{Si}_2\text{O}_{13}$	Hexag.
839. Thuringite. . . . .	$\text{H}_{18}\text{Fe}_8(\text{Al}, \text{Fe})_8\text{Si}_4\text{O}_{41}$	Massive
840. Chamosite. . . . .	$\text{H}_6(\text{Fe}, \text{Mg})_3\text{Al}_2\text{Si}_2\text{O}_{13}$	Oolitic
841. Stilpnomelane. . . . .	$2(\text{Mg}, \text{Fe})\text{O} \cdot (\text{Fe}, \text{Al})_3\text{O}_3 \cdot 5\text{SiO}_2 \cdot 3\text{H}_2\text{O}$	Scaly
842. Strigovite. . . . .	$\text{H}_4\text{Fe}_3(\text{Al}, \text{Fe})_2\text{Si}_4\text{O}_{11}$	Hexag.
843. Rumpfite. . . . .	$\text{H}_{38}\text{Mg}_7\text{Al}_{16}\text{Si}_{10}\text{O}_{65}$	Massive
844. Vermiculite. . . . .	$\text{H}_{24}\text{Mg}_{12}(\text{Al}, \text{Fe})_4\text{Si}_9\text{O}_{46}$	Crystal.
845. Jefferisite. . . . .	$\text{H}_{12}\text{Mg}_4(\text{Al}, \text{Fe})_4\text{Si}_5\text{O}_{46}$	Crystal.
c. Serpentine, Talc		
846. Serpentine. . . . .	$\text{H}_4\text{Mg}_3\text{Si}_2\text{O}_9$	Mono.
847. Bastite. . . . .	$\text{H}_4\text{Mg}_3\text{Si}_2\text{O}_9$	Massive
848. Retinalite. . . . .	$\text{H}_4\text{Mg}_3\text{Si}_2\text{O}_9 + 3$ per cent $\text{H}_2\text{O}$	Massive
849. Bowenite. . . . .	$\text{H}_4\text{Mg}_3\text{Si}_2\text{O}_9$	Massive
850. Antigorite. . . . .	$\text{H}_4\text{Mg}_3\text{Si}_2\text{O}_9$	Massive
851. Marmolite. . . . .	$\text{H}_4\text{Mg}_3\text{Si}_2\text{O}_9$	Foliated
852. Chrysotile. . . . .	$\text{H}_4\text{Mg}_3\text{Si}_2\text{O}_9$	Fibrous
853. Picrolite. . . . .	$\text{H}_4\text{Mg}_3\text{Si}_2\text{O}_9$	Column.
854. Ophicalcite. . . . .	$\text{H}_4\text{Mg}_3\text{Si}_2\text{O}_9 + \text{MgCaCO}_3$	Massive

## LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
808.	Black	3	3	New York	} Elec. purposes
809.	Red	2	2.9	Piedmont	
810.	Brown	1.5	2.9	California	
811.	Grayish	3.5	2.9	North Carolina	
812.	Brown	4	3	New York	
813.	Bottle green	4	3	Ural Mts.	
814.	Gray	6.5	3.5	Michigan	
815.	Green	6.5	3.5	Switzerland	
816.	Green	6.5	3	Belgium	
817.	Green	6.5	3	Rhode Island	
818.	Gray	6	3	Luxembourg	
819.	Gray	5.5	3	Pyrenees Mts.	
820.	Gray	5.5	3	New England	
821.	Green	2	2.6	Pennsylvania	
822.	White	2	2.6	Ural Mts.	
823.	Red	2	2.6	Southern Urals	
824.	Reddish	.....	.....	Sweden	
825.	Green	2	2.6	Zermatt	
826.	Reddish violet	2	2.6	Pennsylvania	} Rock forming
827.	Green	2	2.6	Pennsylvania	
828.	Green	1	2.7	North Carolina	
829.	Green	2.5	2.9	Massachusetts	
830.	Green	2.5	2.7	Italy	
831.	Green	.....	2.8	Colorado	
832.	Green	2	2.7	Connecticut	
833.	Blackish green	2.5	2.8	Nova Scotia	
834.	Green	2	2.7	Hartz	
835.	Green	2.5	2.6	Finland	
836.	Green	1.5	2	Scotland	
837.	Velvet black	2	2	Ireland	
838.	Black	3	3	Bohemia	
839.	Green	2.5	3	Arkansas	
840.	Gray	3	3	Chamoson	
841.	Black	3	2.7	Silesia	
842.	Green	1	3	Silesia	
843.	White	1.5	2.6	Upper Styria	
844.	Brown	1.5	2.7	Massachusetts	
845.	Brown	1.5	2.3	Massachusetts	
846.	Green	2.5	2.5	Maine	} Rock forming
847.	.....	.....	.....	.....	
848.	Yellow	3.5	2.4	Tyrol	} Rock forming
849.	Green	5.5	2.5	Rhode Island	
850.	Green	2.5	2.6	Piedmont	
851.	White	4.5	2.4	New Jersey	} Cloth
852.	White	.....	2.2	Canada	
853.	Green	2	2.5	Maryland	} Rock forming
854.	Green	3	2.5	Pennsylvania	

	Composition	Form
<b>VII. SILICATES—continued</b>		
855. Deweylite.....	$4\text{MgO} \cdot 3\text{SiO}_2 \cdot 6\text{H}_2\text{O}$	Amorph.
856. Genthite.....	$2\text{NiO} \cdot 2\text{MgO} \cdot 3\text{SiO}_2 \cdot 6\text{H}_2\text{O}$	Amorph.
857. Garnierite.....	$\text{H}_2(\text{Ni}, \text{Mg})\text{SiO}_4 \cdot \text{H}_2\text{O}$	Amorph.
858. Talc.....	$\text{H}_2\text{Mg}_3(\text{SiO}_3)_4$	Ortho.
859. Steatite.....	$\text{H}_2\text{Mg}_3(\text{SiO}_3)_4$	Ortho.
860. French chalk.....	$\text{H}_2\text{Mg}_3(\text{SiO}_3)_4$	Ortho.
861. Rensselaerite.....	$\text{H}_2\text{Mg}_3(\text{SiO}_3)_4$	Ortho.
862. Sepiolite.....	$\text{H}_4\text{Mg}_2\text{Si}_2\text{O}_{10}$	Earthy
863. Connarite.....	$\text{H}_4\text{Ni}_2\text{Si}_2\text{O}_{10}$	Hexag.
864. Spadaite.....	$5\text{MgO} \cdot 6\text{SiO}_2 \cdot 4\text{H}_2\text{O}$	Amorph.
865. Saponite.....	Mg, Al, H, silicate	Amorph.
866. Celadonite.....	Fe, Mg, K, silicate	Earthy
867. Glauconite.....	Fe, K, H, silicate	Amorph.
868. Pholidolite.....	$5\text{H}_2\text{O} \cdot \text{K}_2\text{O} \cdot 12(\text{Fe}, \text{Mg})\text{O} \cdot \text{Al}_2\text{O}_3 \cdot 13\text{SiO}_2$	Amorph.
<i>d. Kaolins</i>		
869. Kaolinite.....	$\text{H}_4\text{Al}_2\text{Si}_2\text{O}_9$	Mono.
870. Lithomarge.....	$\text{H}_4\text{Al}_2\text{Si}_2\text{O}_9$	Compact
871. Pholerite.....	$\text{H}_4\text{Al}_2\text{Si}_2\text{O}_9$	Compact
872. Halloysite.....	$(\text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2)$	Massive
873. Pseudosteatite.....	$(\text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2)$	Massive
874. Indianaitite.....	$(\text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2)$	Massive
875. Smectite.....	$(\text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2)$	Massive
876. Bole.....	$(\text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2)$	Massive
877. Bergseife.....	$(\text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2)$	Massive
878. Newtonite.....	$\text{H}_4\text{Al}_2\text{Si}_2\text{O}_{11} \cdot n\text{H}_2\text{O}$	Hexag.
879. Cimolite.....	$2\text{Al}_2\text{O}_3 \cdot 9\text{SiO}_2 \cdot 6\text{H}_2\text{O}$	Amorph.
880. Montmorillonite.....	$\text{H}_4\text{Al}_2\text{Si}_2\text{O}_{11} \cdot n\text{H}_2\text{O}$	Massive
881. Stolpenite.....	$\text{H}_4\text{Al}_2\text{Si}_2\text{O}_{11} \cdot n\text{H}_2\text{O}$	Massive
882. Pyrophyllite.....	$\text{H}_2\text{Al}_2(\text{SiO}_3)_4$	Massive
883. Allophane.....	$\text{Al}_2\text{SiO}_5 \cdot 5\text{H}_2\text{O}$	Amorph.
884. Collyrite.....	$2\text{Al}_2\text{O}_3 \cdot \text{SiO}_2 \cdot 9\text{H}_2\text{O}$	Massive
885. Schrotterite.....	$8\text{Al}_2\text{O}_3 \cdot 3\text{SiO}_2 \cdot 30\text{H}_2\text{O}$	Massive
886. Cenosite.....	$\text{H}, \text{Ca}, (\text{Y}, \text{Er}), \text{CSi}_4\text{O}_{17}$	Ortho.
887. Thaumassite.....	$\text{CaSiO}_3 \cdot \text{CaCO}_3 \cdot \text{CaSO}_4 \cdot 15\text{H}_2\text{O}$	Tetrag.
888. Uranophane.....	$\text{CaO} \cdot 2\text{UO}_3 \cdot 2\text{SiO}_2 \cdot 6\text{H}_2\text{O}$	Ortho.
889. Chrysocolla.....	$\text{CuSiO}_3 \cdot 2\text{H}_2\text{O}$	Amorph.
890. Chloropal.....	$\text{H}_2\text{Fe}_2\text{Si}_2\text{O}_{11} \cdot 2\text{H}_2\text{O}$	Massive
891. Nontronite.....	$\text{H}_2\text{Fe}_2\text{Si}_2\text{O}_{11} \cdot 2\text{H}_2\text{O}$	Massive
892. Pinguite.....	$\text{H}_2\text{Fe}_2\text{Si}_2\text{O}_{11} \cdot 2\text{H}_2\text{O}$	Massive
893. Graminite.....	$\text{H}_2\text{Fe}_2\text{Si}_2\text{O}_{11} \cdot 2\text{H}_2\text{O}$	Massive
894. Hoferite.....	$2\text{Fe}_2\text{O}_3 \cdot 4\text{SiO}_2 \cdot 7\text{H}_2\text{O}$	Amorph.
895. Hisingerite.....	Fe, H, silicate	Amorph.
896. Bementite.....	$2\text{MnSiO}_3 \cdot \text{H}_2\text{O}$	Foliated
897. Caryophilite.....	$4\text{MnO} \cdot 3\text{SiO}_2 \cdot 3\text{H}_2\text{O}$	Massive
898. Neotocite.....	FeMgH, silicate	Amorph.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
855.	Whitish	2	2	Maryland	Rock forming
856.	Green	3	2	Texas	
857.	Green	.....	2.3	North Carolina	Nickel
858.	Green	1	2.7	Vermont	Lubricants
859.	Gray	1	2.5	Virginia	Soapstone
860.	White	1	2.5	Virginia	Lubricants
861.	White	1	2.5	New York	
862.	White	2	2	Asia Minor	Nickel
863.	Green	2.5	2	Saxony	
864.	Reddish	2.5	.....	Italy	Magnesium
865.	White	1	2	Scotland	Rock forming
866.	Green	1	.....	Verona	
867.	Green	2	2	New Jersey	Fertilizer
868.	Yellow	2	2	Sweden	Rock forming
869.	White	2	2.6	Delaware	Brick, fire clay, pottery, and rock forming
870.	White	2	2	Germany	
871.	White	2	2	France	
872.	White	1	2	Illinois	
873.	Green	2	2	Illinois	
874.	White	2	2	Indiana	
875.	Greenish	2	2	France	
876.	Brown	2	2	Illinois	
877.	Brown	2	2	California	
878.	White	1	2	Arkansas	
879.	White	1	2	Argentina	
880.	White	1	2	St. Jean Ode-Cole	
881.	White	1	2	France	
882.	White	1	2.8	North Carolina	
883.	Blue	3	1.8	Pennsylvania	
884.	White	1	2	Pyrenees	
885.	Green	3	1.9	Alabama	
886.	Brown	5	3	Norway	
887.	White	3.5	1.8	New Jersey	
888.	Yellow	3.5	1.8	North Carolina	
889.	Green	2	2	New Jersey	
890.	Yellow	2.5	1.7	Pennsylvania	
891.	Yellow	2.5	1.7	France	
892.	Green	1	1.7	Saxony	
893.	Green	1	1.7	Menzenberg	
894.	Green	1	2.3	Bohemia	
895.	Black	3	2.5	Sweden	
896.	Yellow	1	2.9	New Jersey	
897.	Brown	3	2.8	Sweden	
898.	Black	3	2.6	Sweden	



	Composition	Form
<b>VIIa. TITANATES</b>		
899. Titanite.....	$\text{CaTiSiO}_5$	Mono.
900. Sphene.....	$\text{CaTiSiO}_5$	Mono.
901. Ligurite.....	$\text{CaTiSiO}_5$	Mono.
902. Spinthere.....	$\text{CaTiSiO}_5$	Mono.
903. Lederite.....	$\text{CaTiSiO}_5$	Mono.
904. Titanomorphite.....	$\text{CaTiSiO}_5$	Mono.
905. Greenovite.....	$\text{CaTiSiO}_5$	Mono.
906. Grothite.....	$\text{CaTiSiO}_5$	Mono.
907. Keilhauite.....	$15\text{CaSiTiO}_5 \cdot (\text{Al, Fe, Y})_2(\text{Si, Ti})\text{O}_2$	Mono.
908. Guarinite.....	$\text{CaTiSiO}_5$	Ortho.
909. Tscheffkinite.....	$15\text{CaSiTiO}_5 \cdot (\text{Al, Fe, Y})_2(\text{Si, Ti})\text{O}_2$	Massive
910. Astrophyllite.....	$(\text{Na, K})_4(\text{Fe, Mn})_4\text{Ti}(\text{SiO}_4)_4$	Ortho.
911. Johnstrupite.....	$\text{Ce, Ca, Na, Ti, Fe, silicate}$	Mono.
912. Mosandrite.....	$\text{Ce, Ca, Na, Ti, Fe, silicate}$	Prism.
913. Rinkite.....	$\text{Ce, Ca, Na, Ti, Fe, silicate}$	Mono.
914. Neptunite.....	$\text{Ce, Ca, Na, Ti, Fe, silicate}$	Mono.
915. Perovskite.....	$\text{CaTiO}_3$	Regular
916. Knopite.....	$\text{CaTiO}_3$ , much Ce	Regular
917. Dysanallyte.....	$6(\text{Ca, Fe})\text{TiO}_3 \cdot (\text{Ca, Fe})\text{Nb}_2\text{O}_6$	Regular
918. Geikielite.....	$\text{MgTiO}_3$	Massive

LIST OF MINERALS

No	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
899.	Brown	5	3	Massachusetts	Rock forming
900.	Brown	5	3	Massachusetts	
901.	Yellow	5	3	Massachusetts	
902.	Green	5	3	Massachusetts	
903.	Brown	5	3	Massachusetts	
904.	White	5	3	Massachusetts	
905.	Red	5	3	Massachusetts	
906.	Brown	6	3	Dresden	
907.	Black	6.5	3.5	Norway	
908.	Yellow	6	3	Mt. Somma	
909.	Black	5	4.5	Ilmen Mts.	
910.	Yellow	3	3	Colorado	
911.	Green	.....	3	Norway	
912.	Brown	4	2.9	Norway	
913.	Brown	5	3	Greenland	
914.	Black	5	3	South Greenland	
915.	Yellow	5.5	4	New York	
916.	Black	.....	.....	Sweden	
917.	Black	5	4	Baden	
918.	Black	6	4	Ceylon	

	Composition	Form
<b>VIII. NIOBATES, TANTALATES</b>		
919. Pyrochlore.....	(G,Nb,Ti,Th,Ce,Ca,Fe,U,Mg,NaF).O	Regular
920. Hatchettolite.....	(G,Nb,Ti,Th,Ce,Ca,Fe,U,Mg,NaF).O	Regular
921. Microlite.....	Ca <sub>2</sub> Ta <sub>2</sub> O <sub>7</sub>	Regular
922. Pyrrhite.....	Ca <sub>2</sub> Ta <sub>2</sub> O <sub>7</sub> +Nb,Ti,Ce,Na	Regular
923. Fergusonite.....	(Y,Er,Ce)(Nb,Ta)O <sub>4</sub>	Tetrag.
924. Sipylite.....	Er Nb O <sub>6</sub>	Tetrag.
925. Columbite-tantalite..	(Fe,Mn)(Nb,Ta) <sub>2</sub> O <sub>6</sub>	Ortho.
926. Tapiolite.....	Fe(Ta,Nb) <sub>2</sub> O <sub>6</sub>	Regular
927. Yttrotantalite.....	W,Sn,Y,Er,Ce,U,Fe,Ca,H,Nb, tantalate	Ortho.
928. Samarskite.....	G,Sn,W,U,Ce,Di,La,Y,Er,Fe,Mn,Ca, H,Nb, tantalate	Ortho.
929. Annerodite.....	Pyroniobate of U,Y	Ortho.
930. Hielmite.....	Y,Fe,Mn,Ca,Sn,Nb, tantalate	Ortho.
931. Aeschynite.....	Ce,Th,Fe,Ca,Nb, titanate	Ortho.
932. Polymignite.....	Ce,La,Di,Fe,Ca,Nb,Zn,Sn,Th, titanate	Ortho.
933. Euxenite.....	Y,Er,Ce,U,H,Nb, titnate	Ortho.
934. Polycrase.....	G,Nb,Y,Er,Ce,U,Fe,Ta,H <sub>2</sub> O, titanate	Ortho.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
919.	Brown	5	4	Norway	Rock forming
920.	Brown	.....	4.7	North Carolina	
921.	Yellow	5.5	5	Massachusetts	
922.	Yellow	.....	.....	Urals	
923.	Black	5.5	5.8	Carolinas	
924.	Black	6	4.8	Virginia	
925.	Iron black	6	5	N. England states	
926.	Black	6	7	Finland	
927.	Black	5	5.5	Sweden	
928.	Black	5	5.6	North Carolina	
929.	Black	6	5.7	Norway	
930.	Black	5	5.8	Sweden	
931.	Black	5	4.9	Ilmen Mts.	
932.	Black	6	4.7	Norway	
933.	Black	6.5	4.9	Norway	
934.	Black	5	4.9	South Carolina	

	Composition	Form
<b>IX. PHOSPHATES, ARSENATES, ETC.</b>		
<b>1. Anhydrous</b>		
935. Xenotime .....	$\text{YPO}_4$	Tetrag.
936. Monazite .....	$(\text{Ce}, \text{La}, \text{Di})\text{PO}_4$	Mono.
937. Berzeliite .....	$(\text{CaMgMn})_3\text{As}_2\text{O}_8$	Regular
938. Monimolite .....	$(\text{Pb}, \text{Fe}, \text{Ca})_3\text{Sb}_2\text{O}_8$	Regular
939. Carminite .....	$\text{Pb}_3\text{As}_2\text{O}_8 \cdot 10\text{FeAsO}_4$	Ortho.
940. Pucherite .....	$\text{BiVO}_4$	Ortho.
941. Triphylite .....	$\text{LiFePO}_4$	Ortho.
942. Lithiophilite .....	$\text{LiMnPO}_4$	Ortho.
943. Natrophilite .....	$\text{NaMnPO}_4$	.....
944. Beryllonite .....	$\text{NaBePO}_4$	Ortho.
945. Apatite .....	$(\text{CaF}, \text{Cl})\text{Ca}_5(\text{PO}_4)_3$	Hexag.
946. Moroxite .....	$(\text{CaF}, \text{Cl})\text{Ca}_5(\text{PO}_4)_3$	Hexag.
947. Lasurapatite .....	$(\text{CaF}, \text{Cl})\text{Ca}_5(\text{PO}_4)_3$	Hexag.
948. Francolite .....	$(\text{CaF}, \text{Cl})\text{Ca}_5(\text{PO}_4)_3$	Hexag.
949. Manganapatite .....	$(\text{CaF}, \text{Cl})\text{Ca}_5(\text{PO}_4)_3 + \text{Mn}$	Hexag.
950. Phosphorite .....	$(\text{CaF}, \text{Cl})\text{Ca}_5(\text{PO}_4)_3$	Concret.
951. Eupyrchroite .....	$(\text{CaF}, \text{Cl})\text{Ca}_5(\text{PO}_4)_3$	Concret.
952. Staffelite .....	$(\text{CaF}, \text{Cl})\text{Ca}_5(\text{PO}_4)_3$	Concret.
953. Earthy apatite; osteo- lite .....	Altered apatite	Earthy
954. Pyromorphite .....	$(\text{PbCl})\text{Pb}_4(\text{PO}_4)_3$	Hexag.
955. Polysphaerite .....	$(\text{PbCl})\text{Pb}_4(\text{CO}_3)_3 + \text{Ca}$	Hexag.
956. Miesite .....	$(\text{PbCl})\text{Pb}_4(\text{PO}_4)_3 + \text{Ca}$	Hexag.
957. Nussierite .....	Impure polysphaerite	.....
958. Mimeteite .....	$(\text{PbCl})\text{Pb}_4(\text{AsO}_4)_3$	Hexag.
959. Campylite .....	$(\text{PbCl})\text{Pb}_4(\text{AsO}_4)_3 + \text{P}$	Hexag.
960. Endlichite .....	$\text{Pb}_3\text{Cl}(\text{As}, \text{VO}_4)_3$	Hexag.
961. Vanadinite .....	$(\text{PbCl})\text{Pb}_4(\text{VO}_4)_3$	Hexag.
962. Hedyphane .....	$(\text{Pb}, \text{Ca}, \text{Ce})_4(\text{AsO}_4)_3$	Mono.
963. Svabite .....	$\text{Ca}(\text{F}, \text{Cl}, \text{OH})\text{Ca}_4(\text{AsO}_4)_3$	Hexag.
964. Wagnerite .....	$(\text{MgF})\text{MgPO}_4$	Mono.
965. Spodiosite .....	$(\text{CaF})\text{CaPO}_4$	Mono.
966. Triplite .....	$(\text{Fe}, \text{Mn})\text{PO}_4$	Mono.
967. Talktriplite .....	$(\text{Fe}, \text{Mn}, \text{Ca}, \text{Mg})\text{PO}_4$	Mono.
968. Triploidite .....	$(\text{Fe}_3, \text{Mn}, \text{OH})\text{PO}_4$	Mono.
969. Adelite .....	$(\text{MgOH})\text{CaAsO}_4$	Mono.
970. Tilasite .....	$(\text{Mg}, \text{FOH})\text{CaAsO}_4$	Mono.
971. Sarkinite .....	$(\text{MnOH})\text{MnAsO}_4$	Mono.
972. Herderite .....	$(\text{CaF})\text{BePO}_4$	Mono.
973. Hamlinite .....	$\text{Al}_3\text{Sr}(\text{OH})_2\text{P}_2\text{O}_7$	Hexag.
974. Durangite .....	$\text{Na}(\text{AlF})\text{AsO}_4$	Mono.
975. Amblygonite .....	$\text{Li}(\text{AlF})\text{PO}_4$	Triclinic
<b>2. Basic</b>		
976. Olivenite .....	$\text{Cu}_3\text{As}_2\text{O}_8 \cdot \text{Cu}(\text{OH})_2$	Ortho.
977. Libethenite .....	$\text{Cu}_3\text{P}_2\text{O}_8 \cdot \text{Cu}(\text{OH})_2$	Ortho.

## LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
935.	Brown	4	4	Georgia	Rock forming
936.	Red	5	4.9	Connecticut	
937.	Yellow	5	4	Sweden	
938.	Brown	.....	6.5	Pajsberg	
939.	Red	2.5	4	Nassau	
940.	Brown	4	6	Saxony	
941.	Gray	4.5	3	Massachusetts	
942.	Yellow	4.5	3	Connecticut	
943.	Wine yellow	4.5	3	Connecticut	
944.	Colorless	5.5	2.8	Maine	
945.	Green	5	3	Maine	Phosphorus
946.	Blue	5	3	Arendal	
947.	Sky blue	5	3	Siberia	
948.	Grayish green	5	3	England	
949.	Green	5	3	Delaware	
950.	Green	5	3	Spain	
951.	Gray	4.5	3	New York	
952.	Yellow	4	3	Staffel	
953.	Green	3.5	6.5	New York	
954.	Green	3.5	6.5	Pennsylvania	Lead
955.	Brown	.....	5.8	Cornwall	
956.	Brown	.....	5	Bohemia	
957.	Yellow	.....	5	France	
958.	Yellow	3.5	7	Pennsylvania	
959.	Brown	.....	7	Cumberland	
960.	Brown	2.7	6.6	Arizona	
961.	Red	2.7	6.6	Arizona	
962.	White	4	5	Sweden	
963.	Colorless	5	3.5	Sweden	Rock forming
964.	White	5	3	Austria	
965.	Ash gray	5	2.9	Sweden	
966.	Gray	4	3	Connecticut	
967.	Gray	4	3	Horsjoberg	
968.	Brown	4.5	3.6	Connecticut	
969.	Yellow	5	3.7	Sweden	
970.	Yellow	5	3.7	Langban	
971.	Red	4	4	Sweden	
972.	White	5	2.9	Maine	Manganese
973.	Colorless	4.5	3	Maine	
974.	Orange red	.....	3.9	Mexico	
975.	White	6	3	Maine	
976.	Green	3	4	Utah	Copper
977.	Green	4	3.6	Cornwall	

	Composition	Form
<b>IX. PHOSPHATES, ARSENATES—<i>continued</i></b>		
978. Tarbuttite . . . . .	$\text{Zn}_3\text{P}_2\text{O}_8 \cdot \text{Zn}(\text{OH})_2$	Triclinic
979. Adamite . . . . .	$\text{Zn}_3\text{As}_2\text{O}_8 \cdot \text{Zn}(\text{OH})_2$	Ortho.
980. Descloizite . . . . .	$(\text{Pb}, \text{Zn})_2(\text{OH})\text{VO}_4$	Ortho.
981. Eusynchite . . . . .	$\text{PbZnCuV}_2\text{O}_8$	Massive
982. Dechenite . . . . .	$\text{PbV}_2\text{O}_6$	Massive
983. Calciovolborthite . . . . .	$(\text{Cu}, \text{Ca})_2\text{V}_2\text{O}_8 \cdot (\text{Cu}, \text{Ca})(\text{OH})_2$	Ortho.
984. Brackebuschite . . . . .	$(\text{Pb}, \text{Fe}, \text{Mn})_2\text{V}_2\text{O}_8 \cdot \text{H}_2\text{O}$	Mono.
985. Psittacinite . . . . .	$(\text{Pb}, \text{Cu})_2(\text{OH})_2\text{V}_2\text{O}_8 \cdot \text{H}_2\text{O}$	Coatings
986. Mottramite . . . . .	$(\text{Pb}, \text{Cu})_2(\text{OH})_2\text{V}_2\text{O}_8 \cdot \text{H}_2\text{O}$	Coatings
987. Clinoclase . . . . .	$\text{Cu}_3\text{As}_2\text{O}_8 \cdot 3\text{Cu}(\text{OH})_2$	Mono.
988. Erinite . . . . .	$\text{Cu}_3\text{As}_2\text{O}_8 \cdot 2\text{Cu}(\text{OH})_2$	Concent.
989. Dihydrate . . . . .	$\text{Cu}_3\text{P}_2\text{O}_8 \cdot 2\text{Cu}(\text{OH})_2$	Mono.
990. Pseudomalachite . . . . .	$\text{Cu}_3\text{P}_2\text{O}_8 \cdot 3\text{Cu}(\text{OH})_2$	Massive
991. Chondarsenite . . . . .	$\text{Mn}_3\text{As}_2\text{O}_8 \cdot 3\text{Mn}(\text{OH})_2$	Mono.
992. Xantharsenite . . . . .	$\text{Mn}_3\text{As}_2\text{O}_8 \cdot 3\text{Mn}(\text{OH})_2 + \text{H}_2\text{O}$	Mono.
993. Dufrenite . . . . .	$\text{FePO}_4 \cdot \text{Fe}(\text{OH})_3$	Ortho.
994. Lazulite . . . . .	$(\text{Fe}, \text{Mn})\text{Al}_2(\text{OH})\text{PO}_4$	Mono.
995. Tavistockite . . . . .	$\text{Ca}_3\text{P}_2\text{O}_8 \cdot 2\text{Al}(\text{OH})_3$	Mono.
996. Cirrolite . . . . .	$\text{Ca}_3\text{Al}(\text{PO}_4)_3 \cdot \text{Al}(\text{OH})_3$	Mono.
997. Arseniosiderite . . . . .	$\text{Ca}_3\text{Fe}(\text{AsO}_4)_3 \cdot 3\text{Fe}(\text{OH})_2$	Tetrag.
998. Allacite . . . . .	$\text{Mn}_3\text{As}_2\text{O}_8 \cdot 4\text{Mn}(\text{OH})_2$	Mono.
999. Synadelphite . . . . .	$2(\text{Al}, \text{Mn})\text{AsO}_4 \cdot 5\text{Mn}(\text{OH})_2$	Mono.
1000. Flinkite . . . . .	$\text{MnAsO}_4 \cdot 2\text{Mn}(\text{OH})_2$	Ortho.
1001. Hematolite . . . . .	$(\text{Al}, \text{Mn})\text{AsO}_4 \cdot 4\text{Mn}(\text{OH})_2$	Hexag.
1002. Retzian . . . . .	$\text{Mn}, \text{Ca}, \text{Ce}, \text{Li}, \text{Ca}, \text{Mg}, \text{arsenate}$	Ortho.
1003. Arsenioleite . . . . .	$\text{Sb}, \text{Fe}, \text{Mn}, \text{Pb}, \text{Ca}, \text{Mg}, \text{HSP}, \text{arsenate}$	Hexag.
1004. Manganostibiite . . . . .	$10\text{MnO} \cdot (\text{Sb}, \text{As})_2\text{O}_3$	Ortho.
1005. Atelestite . . . . .	$\text{H}_2\text{Bi}_2\text{AsO}_8$	Mono.
<b>3. Normal Hydrous</b>		
1006. Struvite . . . . .	$(\text{NH}_4)\text{MgPO}_4 \cdot 6\text{H}_2\text{O}$	Ortho.
1007. Collophanite . . . . .	$\text{Ca}_3\text{P}_2\text{O}_8 \cdot \text{H}_2\text{O}$	Amor.
1008. Pyrophosphorite . . . . .	$\text{Mg}_2\text{P}_2\text{O}_7 \cdot 4(\text{Ca}_3\text{P}_2\text{O}_8 + \text{Ca}_3\text{P}_2\text{O}_7)$	Massive
1009. Hopeite . . . . .	$\text{Zn}_3\text{P}_2\text{O}_8 \cdot \text{H}_2\text{O}$	Ortho.
1010. Dickinsonite . . . . .	$(\text{Mn}, \text{Ca}, \text{Fe}, \text{Na})_3(\text{PO}_4)_2 \cdot \frac{1}{2}\text{H}_2\text{O}$	Hexag.
1011. Fillowite . . . . .	$\text{Fe}, \text{Mn}, \text{Ca}, \text{Na}, \text{Li}, \text{hydrous phosphate}$	Mono.
1012. Roselite . . . . .	$(\text{Ca}, \text{Co}, \text{Mg})_2\text{As}_2\text{O}_8 \cdot 2\text{H}_2\text{O}$	Triclinic
1013. Brandite . . . . .	$\text{Ca}_2\text{MnAs}_2\text{O}_8 \cdot 2\text{H}_2\text{O}$	Triclinic
1014. Fairfieldite . . . . .	$\text{Ca}_2\text{MnP}_2\text{O}_8 \cdot 2\text{H}_2\text{O}$	Triclinic
1015. Messelite . . . . .	$(\text{Ca}, \text{Fe})_3\text{P}_2\text{O}_8 \cdot 2\frac{1}{2}\text{H}_2\text{O}$	Triclinic
1016. Reddingite . . . . .	$\text{Mn}_3\text{P}_2\text{O}_8 \cdot 3\text{H}_2\text{O}$	Ortho.
1017. Picropharmacolite . . . . .	$(\text{Ca}, \text{Mg})_2\text{As}_2\text{O}_8 \cdot 6\text{H}_2\text{O}$	Ortho.
1018. Trichalcite . . . . .	$\text{Cu}_3\text{As}_2\text{O}_8 \cdot 5\text{H}_2\text{O}$	Ortho.
1019. Vivianite . . . . .	$\text{Fe}_3\text{P}_2\text{O}_8 \cdot 8\text{H}_2\text{O}$	Mono.
1020. Symplesite . . . . .	$\text{Fe}_2\text{As}_2\text{O}_8 \cdot 8\text{H}_2\text{O}$	Mono.
1021. Bobierite . . . . .	$\text{Mg}_3\text{P}_2\text{O}_8 \cdot 8\text{H}_2\text{O}$	Mono.
1022. Hoernesite . . . . .	$\text{Mg}_3\text{As}_2\text{O}_8 \cdot 8\text{H}_2\text{O}$	Mono.
1023. Erythrite . . . . .	$\text{Co}_3\text{As}_2\text{O}_8 \cdot 8\text{H}_2\text{O}$	Mono.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
978.	Brown	3.5	4	Rhodesia	Zinc
979.	Yellow	3.5	4	Chile	
980.	Red	3.5	5.9	Arizona	Lead
981.	Red	3	5.5	Baden	
982.	Red	3	5.6	Bavaria	Vanadium
983.	Green	3	3.8	Thuringa	
984.	Black	.....	.....	Argentina	
985.	Green	.....	.....	Montana	
986.	Black	.....	.....	England	Copper
987.	Green	2.5	4	Cornwall	
988.	Green	4	4	Cornwall	
989.	Green	4.5	4	Urals	
990.	Dark green	4	3	Rheinbreitenbach	Manganese
991.	Yellow	3	.....	Sweden	
992.	Yellow	.....	.....	Sweden	Phosphorus
993.	Green	3.5	3	New York	
994.	Blue	5	3	North Carolina	Gems
995.	White	.....	.....	Devonshire	
996.	Yellow	5	3	Sweden	Phosphorus
997.	Brown	1	3	France	
998.	Red	4	3	Sweden	Arsenic
999.	Black	4	3	Sweden	
1000.	Brown	4	3.8	Sweden	Manganese
1001.	Red	3.5	3	Nordmark	
1002.	Brown	4	4	Nordmark	Arsenic
1003.	Red	4	.....	Sweden	
1004.	Black	.....	.....	Sweden	Manganese
1005.	Yellow	3	6	Saxony	
1006.	White	2	1.6	Victoria	Lead
1007.	Colorless	2	2	Sombrero Islands	
1008.	White	3	2	West Indies	Phosphorus
1009.	White	2.5	2.7	Altenberg	
1010.	Green	3.5	3	Connecticut	Zinc
1011.	Yellow	4	3	Connecticut	
1012.	Red	3	3.5	Saxony	Phosphorus
1013.	Colorless	5	3.6	Sweden	
1014.	White	3.5	3	Connecticut	Arsenic
1015.	Colorless	3.5	.....	Hesse	
1016.	White	3	3	Connecticut	Phosphorus
1017.	White	.....	.....	Missouri	
1018.	Green	2	.....	Turginsk	Arsenic
1019.	Colorless	1.5	2.5	New Jersey	
1020.	Indigo	2.5	2.9	Carinthia	Phosphorus
1021.	Colorless	.....	.....	Norway	
1022.	White	1	2.4	Hungary	Arsenic
1023.	Red	1.5	2.9	California	



	Composition	Form
<b>IX. PHOSPHATES, ARSE- NATES—<i>continued</i></b>		
1024. Annabergite.....	$\text{Ni}_3\text{As}_2\text{O}_8 \cdot 8\text{H}_2\text{O}$	Mono.
1025. Cabrerite.....	$(\text{Ni}, \text{Mg})_3\text{As}_2\text{O}_8 \cdot 8\text{H}_2\text{O}$	Mono.
1026. Kottigite.....	$\text{Zn}_3\text{As}_2\text{O}_8 \cdot 8\text{H}_2\text{O}$	Mono.
1027. Rhabdophanite.....	$(\text{Y}, \text{Er})_2\text{O}_3 \cdot (\text{La}, \text{Di})_2\text{O}_3 \cdot \text{P}_2\text{O}_5 \cdot \text{H}_2\text{O}$	Mono.
1028. Churchite.....	$\text{Ce}_2\text{O}_3 \cdot \text{CaO} \cdot \text{PD}_2\text{O}_5 \cdot \text{H}_2\text{O}$	Mono.
1029. Scorodite.....	$\text{FeAsO}_4 \cdot 2\text{H}_2\text{O}$	Ortho.
1030. Strengite.....	$\text{FePO}_4 \cdot 2\text{H}_2\text{O}$	Ortho.
1031. Phosphosiderite.....	$2\text{FePO}_4 \cdot 3\frac{1}{2}\text{H}_2\text{O}$	Ortho.
1032. Barrandite.....	$(\text{Al}, \text{Fe})\text{PO}_4 \cdot 2\text{H}_2\text{O}$	Ortho.
1033. Variscite.....	$\text{AlPO}_4 \cdot 2\text{H}_2\text{O}$	Ortho.
1034. Callanite.....	$\text{AlPO}_4 \cdot 2\frac{1}{2}\text{H}_2\text{O}$	Ortho.
1035. Zepharovichite.....	$\text{AlPO}_4 \cdot 3\text{H}_2\text{O}$	Ortho.
1036. Koninckite.....	$\text{FePO}_4 \cdot 3\text{H}_2\text{O}$	Ortho.
<b>4. Acid Hydrous</b>		
1037. Pharmacolite.....	$\text{HCaAsO}_4 \cdot 2\text{H}_2\text{O}$	Mono.
1038. Haidingerite.....	$\text{HCaAsO}_4 \cdot \text{H}_2\text{O}$	Ortho.
1039. Wapplerite.....	$\text{HCaAsO}_4 \cdot 3\frac{1}{2}\text{H}_2\text{O}$	Mono.
1040. Brushite.....	$\text{HCaPO}_4 \cdot 2\text{H}_2\text{O}$	Mono.
1041. Martinite.....	$\text{H}_2\text{Ca}_3(\text{PO}_4)_4 \cdot \frac{1}{2}\text{H}_2\text{O}$	Hexag.
1042. Newberyite.....	$\text{HMgPO}_4 \cdot 3\text{H}_2\text{O}$	Ortho.
1042½. Stercorite.....	$\text{HNa}(\text{NH}_4)\text{PO}_4 \cdot 4\text{H}_2\text{O}$	Mono.
1043. Hureaulite.....	$\text{H}_2\text{Mn}_5(\text{PO}_4)_4 \cdot 4\text{H}_2\text{O}$	Mono.
1044. Forbesite.....	$\text{H}_2(\text{Ni}, \text{Co})_3\text{As}_2\text{O}_8 \cdot 8\text{H}_2\text{O}$	Mono.
<b>5. Basic Hydrous</b>		
1045. Isoclasite.....	$\text{Ca}_3\text{P}_2\text{O}_8 \cdot \text{Ca}(\text{OH})_2 \cdot 4\text{H}_2\text{O}$	Mono.
1046. Hemaifibrite.....	$\text{Mn}_3\text{As}_2\text{O}_8 \cdot 3\text{Mn}(\text{OH})_2 \cdot 2\text{H}_2\text{O}$	Ortho.
1047. Euchroite.....	$\text{Cu}_3\text{As}_2\text{O}_8 \cdot \text{Cu}(\text{OH})_2 \cdot 6\text{H}_2\text{O}$	Ortho.
1048. Conichalcite.....	$(\text{Cu}, \text{Ca})_3\text{As}_2\text{O}_8 \cdot (\text{Cu}, \text{Ca})(\text{OH})_2 \cdot \frac{1}{2}\text{H}_2\text{O}$	Ortho.
1049. Bayldonite.....	$(\text{Pb}, \text{Cu})_3\text{As}_2\text{O}_8 \cdot (\text{Pb}, \text{Cu})(\text{OH})_2 \cdot \text{H}_2\text{O}$	Mono.
1050. Tagilite.....	$\text{Cu}_3\text{P}_2\text{O}_8 \cdot \text{Cu}(\text{OH})_2 \cdot 2\text{H}_2\text{O}$	Mono.
1051. Leucochalcite.....	$\text{Cu}_3\text{As}_2\text{O}_8 \cdot \text{Cu}(\text{OH})_2 \cdot 2\text{H}_2\text{O}$	Mono.
1052. Volborthite.....	$(\text{Cu}, \text{Ca}, \text{Ba})_3(\text{OH})_2\text{VO}_4 \cdot 6\text{H}_2\text{O}$	Ortho.
1053. Cornwallite.....	$\text{Cu}_3\text{As}_2\text{O}_8 \cdot 2\text{Cu}(\text{OH})_2 \cdot \text{H}_2\text{O}$	Ortho.
1054. Tyrolite.....	$\text{Cu}_3\text{As}_2\text{O}_8 \cdot 2\text{Cu}(\text{OH})_2 \cdot 7\text{H}_2\text{O}$	Ortho.
1055. Chalcophyllite.....	$7\text{CuO} \cdot \text{As}_2\text{O}_5 \cdot 14\text{H}_2\text{O}$	Hexag.
1056. Veszelyite.....	$7(\text{Zn}, \text{Cu}_2)(\text{P}, \text{As})_2\text{O}_5 \cdot 9\text{H}_2\text{O}$	Mono.
1057. Wavellite.....	$4\text{AlPO}_4 \cdot 2\text{Al}(\text{OH})_3 \cdot 9\text{H}_2\text{O}$	Ortho.
1058. Fischerite.....	$\text{AlPO}_4 \cdot \text{Al}(\text{OH})_3 \cdot 2\frac{1}{2}\text{H}_2\text{O}$	Ortho.
1059. Peganite.....	$\text{Al}(\text{PO}_4) \cdot \text{Al}(\text{OH})_3 \cdot 1\frac{1}{2}\text{H}_2\text{O}$	Ortho.
1060. Turquoise.....	$\text{AlPO}_4 \cdot \text{Al}(\text{OH})_3 \cdot \text{H}_2\text{O}$	Amorph.
1061. Wardite.....	$2\text{Al}_2\text{O}_3 \cdot \text{P}_2\text{O}_5 \cdot 4\text{H}_2\text{O}$	Crusting
1062. Sphaerite.....	$4\text{AlPO}_4 \cdot 6\text{Al}(\text{OH})_3$	Ortho.
1063. Liskeardite.....	$(\text{Al}, \text{Fe})\text{AsO}_4 \cdot 2(\text{Al}, \text{Fe})(\text{OH})_3 \cdot 5\text{H}_2\text{O}$	Ortho.
1064. Evansite.....	$2\text{AlPO}_4 \cdot 4\text{Al}(\text{OH})_3 \cdot 12\text{H}_2\text{O}$	Ortho.
1065. Coeruleolactite.....	$3\text{Al}_2\text{O}_3 \cdot 2\text{P}_2\text{O}_5 \cdot 10\text{H}_2\text{O}$	.....
1066. Augelite.....	$2\text{Al}_2\text{O}_3 \cdot \text{P}_2\text{O}_5 \cdot 3\text{H}_2\text{O}$	Mono.

## LIST OF MINERALS

No.	Color	Hardness	Gravity	Locality	Chief Constituent or Use
1024.	Green	2	.....	Nevada	Arsenic
1025.	Green	2	2.9	Spain	
1026.	Red	2.5	3	Schneeberg	Rare elements
1027.	Brown	3.5	3.9	Cornwall	
1028.	Smoke gray	3.5	3	Cornwall	Cerium
1029.	Green	3.5	3	Utah	Arsenic
1030.	Red	3	2.8	Virginia	
1031.	Red	3.7	2.7	Germany	Phosphorus
1032.	Gray	4.5	2.5	Bohemia	
1033.	Green	4	.....	Utah	
1034.	Green	3.5	2.5	Lockmariaquer	
1035.	White	5.5	2.3	Bohemia	
1036.	Yellow	3.5	2.3	Belgium	
1037.	White	2	2.6	Joachimsthal	Arsenic
1038.	White	1.5	2	Joachimsthal	Rock forming
1039.	Colorless	2	2	Joachimsthal	
1040.	Colorless	2	2	Caribbean Sea	Arsenic
1041.	Yellowish	2.8	.....	Western India	
1042.	White	3	2	Victoria	Phosphorus
1042½.	White	2	1.6	Peru	
1043.	Yellow	5	3	Connecticut	Arsenic
1044.	White	2.5	3	Atacama	
1045.	White	1	2.9	Joachimsthal	Calcium
1046.	Red	3	3.5	Sweden	Manganese
1047.	Green	3.5	3	Hungary	
1048.	Green	4	4	Utah	Copper
1049.	Green	4	5	Cornwall	
1050.	Green	3	4	Urals	
1051.	White	.....	.....	Germany	
1052.	Green	3	3	Urals	
1053.	Green	4	4	Cornwall	
1054.	Green	1	3	Utah	
1055.	Green	2	2	Utah	
1056.	Blue	3.5	3.5	Banat	
1057.	White	3	2	Saxony	
1058.	Green	5	2.4	Urals	Phosphorus
1059.	Green	3	2.4	Saxony	
1060.	Green	6	2.6	New Mexico	Gems
1061.	Green	5	2.7	Utah	Phosphorus
1062.	Gray	4	2.5	Bohemia	
1063.	White	.....	.....	Cornwall	Aluminum
1064.	Colorless	4	1.9	Hungary	
1065.	White	.....	.....	Pennsylvania	
1066.	Colorless	.....	2.7	Sweden	

	Composition	Form
<b>IX. PHOSPHATES, ARSENATES—continued</b>		
1067. Berlinite.....	$2\text{Al}_2\cdot\text{O}_3\cdot 2\text{P}_2\text{O}_5\cdot \text{H}_2\text{O}$	Massive
1068. Trolleite.....	$4\text{Al}_2\text{O}_3\cdot 3\text{P}_2\text{O}_5\cdot 3\text{H}_2\text{O}$	Compact
1069. Attacolite.....	$\text{P}_2\text{O}_5\cdot \text{Al}_2\text{O}_3\cdot \text{MnO}\cdot \text{CaO}\cdot \text{H}_2\text{O}$ , etc.	Massive
1070. Pharmacosiderite.....	$6\text{FeAsO}_4\cdot 2\text{Fe}(\text{OH})_3\cdot 12\text{H}_2\text{O}$	Regular
1071. Ludlamite.....	$2\text{Fe}_2\text{P}_2\text{O}_7\text{Fe}(\text{OH})_3\cdot 8\text{H}_2\text{O}$	Mono.
1072. Cacozenite.....	$\text{FePO}_4\text{Fe}(\text{OH})_3\cdot 4\frac{1}{2}\text{H}_2\text{O}$	Tufts
1073. Beraunite.....	$2\text{FePO}_4\cdot \text{Fe}(\text{OH})_3\cdot 2\frac{1}{2}\text{H}_2\text{O}$	Mono.
1074. Childrenite.....	$2\text{AlPO}_4\cdot 2\text{Fe}(\text{OH})_3\cdot 2\text{H}_2\text{O}$	Ortho.
1075. Eosphorite.....	$2\text{AlPO}_4\cdot 2\text{Fe}(\text{OH})_3\cdot 2\text{H}_2\text{O}$	Ortho.
1076. Masapilite.....	$\text{Ca}_2\text{Fe}_2(\text{AsO}_4)_4\cdot 2\text{FeO}(\text{OH})\cdot 5\text{H}_2\text{O}$	Ortho.
1077. Calcioferrite.....	$\text{Ca}_2\text{Fe}_2(\text{PO}_4)_4\cdot \text{Fe}(\text{OH})_3\cdot 8\text{H}_2\text{O}$	Mono.
1078. Borickite.....	$\text{Ca}_2\text{Fe}_2(\text{PO}_4)_4\cdot 12\text{Fe}(\text{OH})_3\cdot 6\text{H}_2\text{O}$	Mono.
1079. Richellite.....	$4\text{FeP}_2\text{O}_8\cdot \text{Fe}_2\text{OF}_3(\text{OH})_2\cdot 36\text{H}_2\text{O}$	Mono.
1080. Liroconite.....	$\text{Cu}_6\text{Al}(\text{AsO}_4)_5\cdot 3\text{CuAl}(\text{OH})_5\cdot 20\text{H}_2\text{O}$	Mono.
1081. Chenevixite.....	$\text{Cu}_2(\text{FeO})_2\text{As}_2\text{O}_8\cdot 3\text{H}_2\text{O}$	Mono.
1082. Henwoodite.....	$\text{Fe}, \text{Cu}, \text{Ca}, \text{Al}, \text{H}$ , phosphate	Mono.
1083. Chalcosiderite.....	$\text{CuO}\cdot 3\text{Fe}_2\text{O}_3\cdot 2\text{P}_2\text{O}_5\cdot 8\text{H}_2\text{O}$	Triclinic
1084. Andrewsrite.....	$5\text{Fe}_2\text{O}_3\cdot \text{P}_2\text{O}_5\cdot 5\text{H}_2\text{O}$	Triclinic
1085. Kehoeite.....	$\text{Fe}, \text{Zn}, \text{Ca}, \text{Mg}, \text{Al}, \text{H}$ , phosphate	Amorph.
1086. Goyazite.....	$\text{Ca}_2\text{Al}_2\text{P}_2\text{O}_8\cdot 9\text{H}_2\text{O}$	Tetrag.
1087. Plumbogummite.....	$\text{PbO}\cdot 2\text{Al}_2\text{O}_3\cdot \text{P}_2\text{O}_5\cdot 9\text{H}_2\text{O}$	Hexag.
1088. Torbernite.....	$\text{Cu}(\text{UO}_2)_2\text{P}_2\text{O}_8\cdot 8\text{H}_2\text{O}$	Tetrag.
1089. Zeunerite.....	$\text{Cu}(\text{UO}_2)_2\text{As}_2\text{O}_8\cdot 8\text{H}_2\text{O}$	Tetrag.
1090. Autunite.....	$\text{Ca}(\text{UO}_2)_2\text{P}_2\text{O}_8\cdot 8\text{H}_2\text{O}$	Ortho.
1091. Uranospinite.....	$\text{Ca}(\text{UO}_2)_2\text{As}_2\text{O}_8\cdot 8\text{H}_2\text{O}$	Ortho.
1092. Uranocircite.....	$\text{Ba}(\text{UO}_2)_2\text{P}_2\text{O}_8\cdot 8\text{H}_2\text{O}$	Ortho.
1093. Phosphuranylite.....	$(\text{UO}_2)_2\text{P}_2\text{O}_8\cdot 6\text{H}_2\text{O}$	Powder
1094. Trögerite.....	$(\text{UO}_2)_2\text{As}_2\text{O}_8\cdot 12\text{H}_2\text{O}$	Mono.
1095. Walpurgite.....	$\text{Bi}_{12}(\text{UO}_2)_3(\text{OH})_{24}(\text{AsO}_4)_4$	Triclinic
1096. Rhagite.....	$2\text{BiAsO}_4\cdot 3\text{Bi}(\text{OH})_3$	Triclinic
1097. Mixite.....	$20\text{CuO}\cdot \text{Bi}_2\text{O}_3\cdot 5\text{As}_2\text{O}_5\cdot 22\text{H}_2\text{O}$	.....
<b>6. Antimonates</b>		
1098. Atopite.....	$\text{Ca}_2\text{Sb}_2\text{O}_7$	Regular
1099. Bindheimite.....	$\text{Pb}_2\text{Sb}_2\text{O}_7\cdot 4\text{H}_2\text{O}$	Amorph.
1100. Romeite.....	$\text{CaSb}_2\text{O}_7$	Tetrag.
1101. Nadorite.....	$\text{PbClSbO}_4$	Ortho.
1102. Ecdemite.....	$\text{Pb}_4\text{As}_2\text{O}_7\cdot 2\text{PbCl}_2$	Tetrag.
1103. Ochrolite.....	$\text{Pb}_2\text{Sb}_2\text{O}_7\cdot 2\text{PbCl}_2$	Ortho.
1103a. Trippkeite.....	$\text{CuO}\cdot \text{AS}_2\text{O}_5$	Tetrag.
1104. Tripuhyite.....	$2\text{FeO}\cdot \text{Sb}_2\text{O}_5$	Ortho.
1105. Derbyllite.....	$\text{FeO}\cdot \text{Sb}_2\text{O}_5\cdot 5\text{FeO}\cdot \text{TiO}_2$	Ortho.
1106. Lewisite.....	$5\text{CaO}\cdot 2\text{TiO}_2\cdot 3\text{Sb}_2\text{O}_5$	Regular
1107. Mauzeilite.....	$\text{Pb}, \text{Ca}, \text{Ti}$ , antimonate	Regular
1108. Ammiolite.....	$\text{Cu}, \text{Hg}, \text{Fe}, \text{S}$ , antimonate	Earthy

## LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
1067.	Colorless	6	2.6	Germany	} Phosphorus
1068.	Green	5.5	3	Sweden	
1069.	Salmon red	5	3	.....	} Arsenic
1070.	Green	2.5	2.9	Utah	
1071.	Green	3	3	Cornwall	} Phosphorus
1072.	Yellow	3	3	Pennsylvania	
1073.	Brown	.....	.....	Bohemia	} Arsenic
1074.	White	4.5	3	Maine	
1075.	Pink	4.5	3	Connecticut	} Phosphorus
1076.	Black	4.5	3.5	Mexico	
1077.	Yellow	2	2	Bavaria	} Arsenic
1078.	Brown	3.5	2.6	Bohemia	
1079.	Yellow	2	2	Belgium	} Phosphorus
1080.	Blue	2	2.8	Cornwall	
1081.	Green	3.5	3.9	Utah	} Arsenic
1082.	Blue	4	2.6	Cornwall	
1083.	Green	4	3	Cornwall	} Phosphorus
1084.	Bluish green	4	3	Cornwall	
1085.	.....	.....	2.3	South Dakota	} Lead
1086.	White	5	3	Brazil	
1087.	Yellowish	4	4	Brittany	} Uranium
1088.	Green	2	3	Cornwall	
1089.	Green	2	3	Cornwall	} Bismuth
1090.	Yellow	2	3	North Carolina	
1091.	Green	2	3	Saxony	} Copper
1092.	Green	.....	3.5	Voigtland	
1093.	Yellow	.....	.....	North Carolina	} Antimony
1094.	Yellow	.....	3	Saxony	
1095.	Yellow	3.5	5.7	Saxony	} Antimony
1096.	Yellow	5	6	Saxony	
1097.	Green	3.5	5	Utah	} Lead
1098.	Yellow	5.5	5	Sweden	
1099.	Gray	4	4	Arkansas	} Lead
1100.	Yellow	5.5	4.7	Piedmont	
1101.	Yellow	3.5	7	Algeria	} Antimony
1102.	Yellow	2.5	7	Sweden	
1103.	Yellow	.....	.....	Chile	} Lead
1103a.	Bluish green	.....	.....	Brazil	
1104.	Greenish yellow	.....	5	Brazil	} Copper
1105.	Black	5	4	Brazil	
1106.	Yellow	.....	4	Brazil	} Antimony
1107.	Brown	6	5	Sweden	
1108.	Scarlet	.....	.....	Chile	} Titanium

	Composition	Form
<b>IX. PHOSPHATES, ARSENATES—<i>continued</i></b>		
7. <i>Mixed Phosphates</i>		
1109. Diadochite.....	$2\text{Fe}_2\text{O}_3 \cdot 2\text{SO}_3 \cdot \text{P}_2\text{O}_5 \cdot 12\text{H}_2\text{O}$	Mono.
1110. Destinezite.....	$2\text{Fe}_2\text{O}_3 \cdot 2\text{SO}_3 \cdot \text{P}_2\text{O}_5 \cdot 12\text{H}_2\text{O}$	Earthy
1111. Pitticite.....	$\text{Fe}_2\text{S}_2$ , arsenate	Massive
1112. Svanbergite.....	$\text{Ca}_2\text{Al}_2\text{S}_2$ , phosphate	Hexag.
1113. Beudantite.....	$\text{Fe}, \text{Pb}, \text{S}_2\text{AS}$ , phosphate	Hexag.
1114. Lindackerite.....	$3\text{NiO} \cdot 6\text{CuO} \cdot \text{SO}_3 \cdot 2\text{As}_2\text{O}_3 \cdot 7\text{H}_2\text{O}$	Ortho.
1115. Lünebergite.....	$3\text{MgO} \cdot \text{B}_2\text{O}_3 \cdot \text{P}_2\text{O}_5 \cdot 8\text{H}_2\text{O}$	Earthy
1116. Lossenite.....	$2\text{PbSO}_4 \cdot 3(\text{FeOH})_2\text{As}_2\text{O}_5 \cdot 12\text{H}_2\text{O}$	Ortho.
8. <i>Nitrates, etc.</i>		
1117. Soda niter.....	$\text{NaNO}_3$	Hexag.
1118. Niter.....	$\text{KNO}_3$	Ortho.
1119. Nitrocalcite.....	$\text{Ca}(\text{NO}_3)_2 \cdot n\text{H}_2\text{O}$	.....
1120. Nitromagnesite.....	$\text{Mg}(\text{NO}_3)_2 \cdot n\text{H}_2\text{O}$	.....
1121. Nitrobarite.....	$\text{Ba}(\text{NO}_3)_2$	Regular
1122. Gerhardite.....	$\text{Cu}(\text{NO}_3)_2 \cdot 3\text{Cu}(\text{OH})_2$	Ortho.
1123. Darapskite.....	$\text{NaNO}_3 \cdot \text{Na}_2\text{SO}_4 \cdot \text{H}_2\text{O}$	Tetrag.
1124. Nitroglauherite.....	$6\text{NaNO}_3 \cdot 2\text{Na}_2\text{SO}_4 \cdot 3\text{H}_2\text{O}$	Tetrag.
1125. Lautarite.....	$\text{Ca}(\text{IO}_3)_2$	Mono.
1126. Dietzeite.....	$7\text{Ca}(\text{IO}_3)_2 \cdot 8\text{CaCrO}_4$	Mono.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
1109.	Yellowish	3	2	Thuringia	Phosphorus
1110.	Yellowish	.....	.....	Belgium	} Arsenic
1111.	Brown	2	2	Saxony	
1112.	Yellow	5	3	Sweden	Phosphorus
1113.	Green	4	4	Cork	Lead
1114.	Green	2.5	.....	Joachimsthal	Copper
1115.	.....	.....	2	Hannover	Phosphorus
1116.	Brownish	.....	.....	Greece	Arsenic
1117.	White	1.5	2	Nevada	} Fertilizer
1118.	White	2	2	Egypt	
1119.	Gray	.....	.....	Kentucky	
1120.	White	.....	.....	Kentucky	
1121.	Colorless	.....	.....	Chile	} Copper
1122.	Green	2	3	Arizona	
1123.	Colorless	.....	.....	Chile	Soda
1124.	White	.....	.....	Atacama	Sodium
1125.	Colorless	.....	4.5	Atacama	} Iodine
1126.	Yellow	3	3.7	Atacama	

## COMPREHENSIVE

	Composition	Form
<b>X. BORATES, ETC.</b>		
1127. Sussexite.....	$2(\text{Mn,Zn,Mg})\text{O} \cdot \text{B}_2\text{O}_3 \cdot \text{H}_2\text{O}$	Ortho.
1128. Ludwigite.....	$3\text{MgO} \cdot \text{B}_2\text{O}_3 \cdot \text{FeO} \cdot \text{Fe}_2\text{O}_3$	Ortho.
1129. Pinakiolite.....	$3\text{MgO} \cdot \text{B}_2\text{O}_3 \cdot \text{MnO} \cdot \text{Mn}_2\text{O}_3$	Ortho.
1130. Nordenskiöldine.....	$\text{CaSn}(\text{BO}_3)_2$	Regular
1131. Jeremejevite.....	$\text{AlBO}_3$	Hexag.
1132. Hambergite.....	$\text{B}_2(\text{OH})\text{BO}_3$	Ortho.
1133. Szaibelyite.....	$2\text{Mg}_5\text{B}_4\text{O}_{11} \cdot 3\text{H}_2\text{O}$	Ortho.
1134. Boracite.....	$\text{Mg}_5\text{Cl}_2\text{B}_{16}\text{O}_{39}$	Regular
1135. Ascharite.....	$3\text{Mg}_2\text{B}_3\text{O}_5 \cdot 2\text{H}_2\text{O}$	Amorph.
1136. Rhodizite.....	$\text{K,Al,Cs,Rb,Na,Ca,Mg,Al, borate}$	Regular
1137. Warwickite.....	$6\text{MgO} \cdot \text{FeO} \cdot 2\text{TiO}_2 \cdot 3\text{B}_2\text{O}_3$	Ortho.
1138. Howlite.....	$\text{H,Ca}_2\text{B}_5\text{SiO}_{14}$	Ortho.
1139. Lagonite.....	$\text{Fe}_2\text{O}_3 \cdot 3\text{B}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$	Earthy
1140. Larderellite.....	$(\text{NH}_4)_2\text{O} \cdot 4\text{B}_2\text{O}_3 \cdot 4\text{H}_2\text{O}$	Mono.
1141. Colemanite.....	$\text{Ca}_2\text{B}_6\text{O}_{11} \cdot 5\text{H}_2\text{O}$	Mono.
1142. Pinnoite.....	$\text{MgB}_2\text{O}_4 \cdot 3\text{H}_2\text{O}$	Tetrag.
1143. Heintzite.....	$\text{K}_2\text{O} \cdot 4\text{MgO} \cdot 9\text{B}_2\text{O}_3 \cdot 16\text{H}_2\text{O}$	Mono.
1144. Borax.....	$\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$	Mono.
1145. Ulexite.....	$\text{NaCaB}_3\text{O}_6 \cdot 8\text{H}_2\text{O}$	Fibers
1146. Bechillite.....	$\text{CaB}_4\text{O}_7 \cdot 4\text{H}_2\text{O}$	Crusts
1147. Hydroboracite.....	$\text{CaMgB}_4\text{O}_{11} \cdot 6\text{H}_2\text{O}$	Mono.
1148. Sulfoborite.....	$3\text{MgSO}_4 \cdot 2\text{Mg}_3\text{B}_4\text{O}_9 \cdot 12\text{H}_2\text{O}$	Ortho.
1149. Uraninite.....	$\text{Pb,Th,G,Ce,La,Y,Ca,N,Fe,H, uranite}$	Regular
1150. Urannibite.....	$\text{Pb,Th,G,Ce,La,Y,Ca,N,Fe,H, uranite}$	Regular
1151. Bröggerite.....	$\text{Pb,Th,G,Ce,La,Y,Ca,N,Fe,H, uranite}$	Regular
1152. Cleveite.....	$\text{Pb,Th,G,Ce,La,Y,Ca,N,Fe,H, more U}$	Regular
1153. Nivenite.....	$\text{Pb,Th,G,Ce,La,Y,Ca,N,Fe,H, more U}$	Regular
1154. Pitchblende.....	$\text{Pb,Th,G,Ce,La,Y,Ca,N,Fe,H, more U}$	Regular
1155. Carnotite.....	$\text{K}_2\text{O} \cdot 2\text{U}_2\text{O}_7 \cdot \text{V}_2\text{O}_5 \cdot 3\text{H}_2\text{O}$	Earthy
1156. Gummite.....	$(\text{Pb,Ca,Ba})\text{U}_3\text{SiO}_{12} \cdot 6\text{H}_2\text{O}$	Amorph.
1157. Yttrogummite.....	$(\text{Pb,Ca,Ba})\text{U}_3\text{SiO}_{12} \cdot 6\text{H}_2\text{O} + \text{Y}$	Earthy
1158. Thorogummite.....	$(\text{Pb,Ca,Ba})\text{U}_3\text{SiO}_{12} \cdot 6\text{H}_2\text{O} + \text{Th}$	Earthy
1159. Uranosphaerite.....	$(\text{BiO})_2\text{U}_2\text{O}_7 \cdot 3\text{H}_2\text{O}$	Globular

## LIST OF MINERALS

No.	Color	Hardness	Gravity	Locality	Chief Constituent or Use
1127.	White	3	3	New Jersey	Manganese
1128.	Green	5	3.9	Hungary	Magnesium
1129.	Black	6	3.8	Sweden	Manganese
1130.	Yellow	5.5	4	Norway	Zinc
1131.	Colorless	6.5	3	Mt. Sektuj	Boron
1132.	White	7.5	2	Norway	Beryllium
1133.	White	3	3	Hungary	
1134.	White	7	2.9	France	
1135.	White	.....	1.9	Germany	
1136.	White	8	3	Urals	
1137.	Brown	3	3	New York	
1138.	White	3.5	2	Nova Scotia	
1139.	Yellow	.....	.....	Tuscany	
1140.	Yellow	.....	.....	Tuscany	Boron
1141.	Colorless	4	2	California	
1142.	Yellow	3	3	Stassfurt	
1143.	Colorless	4	2	Stassfurt	
1144.	White	2	1.6	Nevada	
1145.	White	1	1.6	Nevada	
1146.	Gray	.....	.....	Tuscany	
1147.	White	2	1.9	Caucasus	
1148.	Colorless	4	2	Germany	
1149.	Gray	5.5	9	Connecticut	
1150.	Gray	5.5	9	Norway	
1151.	Gray	5.5	9	Norway	
1152.	Gray	5.5	7	Norway	
1153.	Black	5.5	8	Texas	
1154.	Black	5.5	8	Colorado	Rare elements
1155.	Yellow	.....	.....	Utah	
1156.	Yellow	2.5	3.9	North Carolina	
1157.	Black	5	.....	Norway	
1158.	Brown	4	4	Texas	
1159.	Red	2.3	6	Saxony	



	Composition	Form
<b>XI. SULPHATES, ETC.</b>		
<b>1. Anhydrous Sulphates</b>		
1160. Mascagnite.....	$(\text{NH}_4)_2\text{SO}_4$	Ortho.
1161. Taylorite.....	$5\text{K}_2\text{SO}_4 \cdot (\text{NH}_4)_2\text{SO}_4$	Concret.
1162. Thenardite.....	$\text{Na}_2\text{SO}_4$	Ortho.
1163. Aphthitalite.....	$(\text{K}, \text{Na})_2\text{SO}_4$	Hexag.
1164. Glauberite.....	$\text{Na}_2\text{SO}_4 \cdot \text{CaSO}_4$	Mono.
1165. Langbeinite.....	$\text{K}_2\text{Mg}_2(\text{SO}_4)_3$	Hexag.
1166. Barite.....	$\text{BaSO}_4$	Ortho.
1167. Bologna stone.....	$\text{BaSO}_4$	Ortho.
1168. Cawk.....	$\text{BaSO}_4$	Ortho.
1169. Michel-levyite.....	$\text{BaSO}_4$	Ortho.
1170. Celestite.....	$\text{SrSO}_4$	Ortho.
1171. Apotome.....	$\text{SrSO}_4$	Ortho.
1172. Anglesite.....	$\text{PbSO}_4$	Ortho.
1173. Anhydrite.....	$\text{CaSO}_4$	Ortho.
1174. Vulpinite.....	$\text{CaSO}_4$	Scaly
1175. Tripstone.....	$\text{CaSO}_4$	Concret.
1176. Zinkosite.....	$\text{ZnSO}_4$	Ortho.
1177. Hydrocyanite.....	$\text{CuSO}_4$	Ortho.
1178. Crocoite.....	$\text{PbCrO}_4$	Mono.
1179. Leadhillite.....	$4\text{PbO} \cdot \text{SO}_3 \cdot 2\text{CO}_2 \cdot \text{H}_2\text{O}$	Mono.
1180. Susannite.....	$4\text{PbO} \cdot \text{SO}_3 \cdot 2\text{CO}_2 \cdot \text{H}_2\text{O}$	Mono.
1181. Sulphohalite.....	$3\text{Na}_2\text{SO}_4 \cdot 2\text{NaCl}$	Regular
1182. Caracolite.....	$\text{Pb}(\text{OH})\text{Cl} \cdot \text{Na}_2\text{SO}_4$	Ortho.
1183. Kainite.....	$\text{MgSO}_4 \cdot \text{KCl} \cdot 3\text{H}_2\text{O}$	Mono.
1184. Connellite.....	$\text{Cu}_{11}(\text{Cl}, \text{OH})_4\text{SO}_4 \cdot 15\text{H}_2\text{O}$	Hexag.
1185. Spangolite.....	$\text{Cu}_6\text{AlClSO}_{10} \cdot 9\text{H}_2\text{O}$	Hexag.
1186. Hanksite.....	$9\text{Na}_2\text{SO}_4 \cdot 2\text{Na}_2\text{CO}_3 \cdot \text{KCl}$	Hexag.
1187. Misenite.....	$\text{HKSO}_4$	Mono.
1188. Brochantite.....	$\text{CuSO}_4 \cdot 3\text{Cu}(\text{OH})_2$	Ortho.
1189. Lanarkite.....	$\text{Pb}_2\text{SO}_5$	Mono.
1190. Dolerophanite.....	$\text{Cu}_2\text{SO}_5$	Mono.
1191. Caledonite.....	$2(\text{Pb}, \text{Cu})\text{O} \cdot \text{SO}_3 \cdot \text{H}_2\text{O}$	Ortho.
1192. Linarite.....	$(\text{Pb}, \text{Cu})\text{SO}_4 \cdot (\text{Pb}, \text{Cu})(\text{OH})_2$	Mono.
1193. Antlerite.....	$3\text{CuSO}_4 \cdot 7\text{Cu}(\text{OH})_2$	Massive
1194. Alumian.....	$\text{Al}_2\text{O}_3 \cdot 2\text{SO}_3$	Hexag.
<b>2. Hydrous Sulphates</b>		
<b>a. Normal</b>		
1195. Lecontite.....	$(\text{Na}, \text{NH}_4, \text{K})_2\text{SO}_4 \cdot 2\text{H}_2\text{O}$	Ortho.
1196. Mirabilite.....	$\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$	Mono.
1197. Kieserite.....	$\text{MgSO}_4 \cdot \text{H}_2\text{O}$	Mono.
1198. Szmikite.....	$\text{MnSO}_4 \cdot \text{H}_2\text{O}$	Amorph.
1199. Gypsum.....	$\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$	Mono.
1200. Selenite.....	$\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$	Mono.
1201. Satin spar.....	$\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$	Mono.
1202. Alabaster.....	$\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$	Mono.
1203. Illesite.....	$(\text{Mn}, \text{Zn}, \text{Fe})\text{SO}_4 \cdot 4\text{H}_2\text{O}$	Mono.

## LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
1160.	Yellow	2	1.7	Etna	Sulphur
1161.	White	2	.....	Chincha Islands	
1162.	White	2	2	California	Sulphur
1163.	White	3	2	Vesuvius	Potash
1164.	Yellow	2.5	2.7	California	Potassium
1165.	Colorless	.....	2.8	Germany	Manganese
1166.	White	2.5	4	New York	Barium
1167.	Reddish	2.5	4	Bologna	
1168.	Reddish	2.5	4	Bologna	
1169.	Reddish	2.5	4	Quebec	
1170.	White	3	3.9	Texas	Strontium
1171.	White	3	3.9	Texas	
1172.	White	2.7	6	Pennsylvania	Lead
1173.	White	3	2.8	Tennessee	Calcium
1174.	White	3	2.8	Lombardy	
1175.	White	3	2.8	Lombardy	Zinc
1176.	White	.....	2.8	Spain	
1177.	Green	.....	.....	Vesuvius	Copper
1178.	Red	2.5	5.9	Arizona	Lead
1179.	Yellow	2.5	6	Scotland	
1180.	Yellow	2.5	6	Scotland	Sulphur
1181.	Yellow	3	2	California	
1182.	Colorless	4.5	.....	Atacama	Lead
1183.	White	2	2	Stassfurt	Sulphur
1184.	Blue	3	3	Cornwall	Copper
1185.	Green	2	3	Arizona	
1186.	White	3	2.5	California	Sodium
1187.	White	.....	.....	Naples	Potassium
1188.	Green	3.5	3.9	Colorado	Copper
1189.	White	2	5	Scotland	Lead
1190.	Brown	.....	.....	Vesuvius	Copper
1191.	Green	2.5	6	California	Lead
1192.	Blue	2	5	California	
1193.	Green	.....	3.9	Arizona	Copper
1194.	White	2	2.7	Spain	Sulphur
1195.	Colorless	2	.....	Central America	Sodium
1196.	White	1.5	1	Salt Lake, Utah	
1197.	White	3	2	Stassfurt	Magnesium
1198.	White	1.5	3	Hungary	Manganese
1199.	White	1.5	2	Michigan	Plaster
1200.	White	1.5	2	Michigan	
1201.	White	1.5	2	Michigan	Ornaments
1202.	White	1.5	2	Michigan	
1203.	Green	.....	.....	Colorado	Manganese

	Composition	Form
<b>XI. SULPHATES, ETC.—</b>		
<i>continued</i>		
1204. Epsomite . . . . .	$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	Ortho.
1205. Goslarite . . . . .	$\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$	Ortho.
1206. Morenosite . . . . .	$\text{NiSO}_4 \cdot 7\text{H}_2\text{O}$	Ortho.
1207. Melanterite . . . . .	$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	Mono.
1208. Mallardite . . . . .	$\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$	Mono.
1209. Pisanite . . . . .	$(\text{Fe}, \text{Cu})\text{SO}_4 \cdot 7\text{H}_2\text{O}$	Mono.
1210. Salvadorite . . . . .	$(\text{Cu}, \text{Fe})\text{SO}_4 \cdot 7\text{H}_2\text{O}$	Mono.
1211. Bieberite . . . . .	$\text{CoSO}_4 \cdot 7\text{H}_2\text{O}$	Mono.
1212. Chalcanthite . . . . .	$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$	Triclinic
1213. Syngenite . . . . .	$\text{CaSO}_4 \cdot \text{K}_2\text{SO}_4 \cdot \text{H}_2\text{O}$	Mono.
1214. Löweite . . . . .	$\text{MgSO}_4 \cdot \text{Na}_2\text{SO}_4 \cdot 2\frac{1}{2}\text{H}_2\text{O}$	Tetrag.
1215. Blödite . . . . .	$\text{MgSO}_4 \cdot \text{Na}_2\text{SO}_4 \cdot 4\text{H}_2\text{O}$	Mono.
1216. Leonite . . . . .	$\text{MgSO}_4 \cdot \text{K}_2\text{SO}_4 \cdot 4\text{H}_2\text{O}$	Mono.
1217. Boussingaultite . . . . .	$(\text{NH}_4)_2\text{SO}_4 \cdot \text{MgSO}_4 \cdot 6\text{H}_2\text{O}$	Mono.
1218. Picromerite . . . . .	$\text{MgSO}_4 \cdot \text{K}_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$	Mono.
1219. Polyhalite . . . . .	$2\text{CaSO}_4 \cdot \text{MgSO}_4 \cdot \text{K}_2\text{SO}_4 \cdot 2\text{H}_2\text{O}$	Mono.
1220. Pickeringite . . . . .	$\text{MgSO}_4 \cdot \text{Al}_2(\text{SO}_4)_3 \cdot 22\text{H}_2\text{O}$	Mono.
1221. Halotrichite . . . . .	$\text{FeSO}_4 \cdot \text{Al}_2(\text{SO}_4)_3 \cdot 24\text{H}_2\text{O}$	Mono.
1222. Apjohnite . . . . .	$\text{MnSO}_4 \cdot \text{Al}_2(\text{SO}_4)_3 \cdot 24\text{H}_2\text{O}$	Mono.
1223. Dietrichite . . . . .	$(\text{Zn}, \text{Fe}, \text{Mn})\text{SO}_4 \cdot \text{Al}_2(\text{SO}_4)_3 \cdot 22\text{H}_2\text{O}$	Mono.
1224. Masrite . . . . .	$\text{Fe}, \text{Mn}, \text{Co}, \text{Al}$ , sulphate	Fibrous
1225. Coquimbite . . . . .	$\text{Fe}_2(\text{SO}_4)_3 \cdot 9\text{H}_2\text{O}$	Hexag.
1226. Quenstedtite . . . . .	$\text{Fe}_2(\text{SO}_4)_3 \cdot 10\text{H}_2\text{O}$	Mono.
1227. Ihleite . . . . .	$\text{Fe}_2(\text{SO}_4)_3 \cdot 12\text{H}_2\text{O}$	Efflor.
1228. Alunogen . . . . .	$\text{Al}_2(\text{SO}_4)_3 \cdot 18\text{H}_2\text{O}$	Mono.
1229. Kröhnkite . . . . .	$\text{CuSO}_4 \cdot \text{Na}_2\text{SO}_4 \cdot 2\text{H}_2\text{O}$	Mono.
1230. Phillipite . . . . .	$\text{CuSO}_4 \cdot \text{Fe}_2(\text{SO}_4)_3 \cdot n\text{H}_2\text{O}$	Mono.
1231. Ferronatrite . . . . .	$3\text{Na}_2\text{SO}_4 \cdot \text{Fe}_2(\text{SO}_4)_3 \cdot 6\text{H}_2\text{O}$	Hexag.
1232. Römerite . . . . .	$\text{FeSO}_4 \cdot \text{Fe}_2(\text{SO}_4)_3 \cdot 12\text{H}_2\text{O}$	Triclinic
1233. Natrochalcite . . . . .	$\text{Na}_2\text{SO}_4 \cdot \text{Cu}_4(\text{OH})_2(\text{SO}_4)_2 \cdot 2\text{H}_2\text{O}$	Mono.
<i>b. Basic</i>		
1234. Langite . . . . .	$\text{CuSO}_4 \cdot 3\text{Cu}(\text{OH})_2 \cdot \text{H}_2\text{O}$	Ortho.
1235. Herrengrundite . . . . .	$2(\text{CuOH})_2\text{SO}_4 \cdot \text{Cu}(\text{OH})_2 \cdot 3\text{H}_2\text{O}$	Mono.
1236. Kamarezite . . . . .	$(\text{CuOH})_2\text{SO}_4 \cdot \text{Cu}(\text{OH})_2 \cdot 6\text{H}_2\text{O}$	Ortho.
1237. Cyanotrichite . . . . .	$4\text{CuO} \cdot \text{Al}_2\text{O}_3 \cdot \text{SO}_3 \cdot 8\text{H}_2\text{O}$	Ortho.
1238. Serpierite . . . . .	$\text{Cu}, \text{Zn}$ , sulphate	Ortho.
1239. Copiapite . . . . .	$2\text{Fe}_2\text{O}_3 \cdot 5\text{SO}_3 \cdot 18\text{H}_2\text{O}$	Mono.
1240. Castanite . . . . .	$\text{Fe}_2\text{O}_3 \cdot 2\text{SO}_3 \cdot 8\text{H}_2\text{O}$	Mono.
1241. Utahite . . . . .	$3\text{Fe}_2\text{O}_3 \cdot 2\text{SO}_3 \cdot 7\text{H}_2\text{O}$	Hexag.
1242. Amarantite . . . . .	$\text{Fe}_2\text{O}_3 \cdot 2\text{SO}_3 \cdot 7\text{H}_2\text{O}$	Triclinic
1243. Fibroferrite . . . . .	$\text{Fe}_2\text{O}_3 \cdot 2\text{SO}_3 \cdot 10\text{H}_2\text{O}$	Mono.
1244. Raimondite . . . . .	$2\text{Fe}_2\text{O}_3 \cdot 3\text{SO}_3 \cdot 7\text{H}_2\text{O}$	Hexag.
1245. Carphosiderite . . . . .	$3\text{Fe}_2\text{O}_3 \cdot 4\text{SO}_3 \cdot 10\text{H}_2\text{O}$	Hexag.
1246. Glockerite . . . . .	$2\text{Fe}_2\text{O}_3 \cdot \text{SO}_3 \cdot 6\text{H}_2\text{O}$	Earthy
1247. Knoxvilleite . . . . .	$\text{Cr}, \text{Fe}, \text{Al}, \text{H}$ , sulphate	Ortho.
1248. Redingtonite . . . . .	$\text{Cr}, \text{Fe}, \text{Al}, \text{H}$ , sulphate	Ortho.

## LIST OF MINERALS

No.	Color	Hardness	Gravity	Locality	Chief Constituent or Use
1204.	White	2	1.7	Kentucky	Medicine
1205.	White	2	1.9	Montana	Zinc
1206.	Green	2	2	Galicia	Nickel
1207.	Green	2	1.8	Utah	Iron
1208.	Colorless	.....	.....	Utah	Manganese
1209.	Blue	.....	.....	Turkey	} Copper
1210.	Green	.....	.....	Chile	
1211.	Red	.....	1.9	Bieber	Cobalt
1212.	Blue	2.5	2	Arizona	Copper
1213.	Colorless	2	2	Galicia	Potassium
1214.	Yellow	2	2	Austria	Sodium
1215.	Colorless	2	2	Chile	Magnesium
1216.	White	.....	.....	Germany	Potassium
1217.	White	.....	1.6	Tuscany	Magnesium
1218.	White	.....	2	Vesuvius	Potassium
1219.	Red	2	2	Austria	Calcium
1220.	White	1	.....	Colorado	} Aluminum
1221.	Yellow	.....	.....	New Mexico	
1222.	Yellow	1.5	1.7	Tennessee	
1223.	Yellow	2	.....	Hungary	} .....
1224.	.....	.....	.....	Egypt	
1225.	White	2	2	Chile	} Iron
1226.	Red	2	2	Chile	
1227.	Yellow	.....	1.8	Bohemia	} Aluminum
1228.	White	1.5	1.6	Bohemia	
1229.	Blue	2.5	1.9	Atacama	} Copper
1230.	Blue	.....	.....	Chile	
1231.	Gray	2	2	Chile	Sodium
1232.	Brown	3	2	Chile	Iron
1233.	Green	4	2	Chile	Copper
1234.	Blue	2	3	Cornwall	} Copper
1235.	Green	2	3	Hungary	
1236.	Green	3	3	Greece	
1237.	Blue	.....	.....	Utah	
1238.	Green	.....	.....	Greece	
1239.	Yellow	2.5	2	Chile	} Iron
1240.	Brown	3	2	Chile	
1241.	Yellow	.....	.....	Utah	
1242.	Red	2	2	Chile	
1243.	Yellow	2	1.8	Chile	
1244.	Yellow	3	3	Bolivia	
1245.	Yellow	4	2	Greenland	
1246.	Brown	.....	.....	Harz	
1247.	Yellow	.....	.....	California	} Chromium
1248.	Pale purple	.....	1.7	Knoxville	

	Composition	Form
<b>XI. SULPHATES, ETC.—</b>		
<i>continued</i>		
1249. Cyprusite.....	$7\text{Fe}_2\text{O}_3 \cdot \text{Al}_2\text{O}_3 \cdot 10\text{SO}_3 \cdot 14\text{H}_2\text{O}$	Hexag.
1250. Aluminite.....	$\text{Al}_2\text{O}_3 \cdot \text{SO}_3 \cdot 9\text{H}_2\text{O}$	Mono.
1251. Paraluminite.....	$2\text{Al}_2\text{O}_3 \cdot \text{SO}_3 \cdot 15\text{H}_2\text{O}$	Mono.
1252. Felsöbanyite.....	$2\text{Al}_2\text{O}_3 \cdot \text{SO}_3 \cdot 10\text{H}_2\text{O}$	Ortho.
1253. Botryogen.....	$\text{MgO} \cdot \text{FeO} \cdot \text{Fe}_2\text{O}_3 \cdot 4\text{SO}_3 \cdot 18\text{H}_2\text{O}$	Mono.
1254. Sideronatrite.....	$2\text{Na}_2\text{O} \cdot \text{Fe}_2\text{O}_3 \cdot 4\text{SO}_3 \cdot 7\text{H}_2\text{O}$	Ortho.
1255. Voltaite.....	$5(\text{K}_2, \text{Fe})\text{O} \cdot 2(\text{Al}, \text{Fe})_2\text{O}_3 \cdot 10\text{SO}_3 \cdot 15\text{H}_2\text{O}$	Regular
1256. Metavoltine.....	$5(\text{K}_2, \text{Na}_2, \text{Fe})\text{O} \cdot 3\text{Fe}_2\text{O}_3 \cdot 12\text{SO}_3 \cdot 18\text{H}_2\text{O}$	Hexag.
1257. Alunite.....	$\text{K}_2\text{O} \cdot 3\text{Al}_2\text{O}_3 \cdot 4\text{SO}_3 \cdot 6\text{H}_2\text{O}$	Hexag.
1258. Jarosite.....	$\text{K}_2\text{O} \cdot 3\text{Fe}_2\text{O}_3 \cdot 4\text{SO}_3 \cdot 6\text{H}_2\text{O}$	Hexag.
1259. Lowigite.....	$\text{K}_2\text{O} \cdot 3\text{Al}_2\text{O}_3 \cdot 4\text{SO}_3 \cdot 9\text{H}_2\text{O}$	Hexag.
1260. Ettringite.....	$6\text{CaO} \cdot \text{Al}_2\text{O}_3 \cdot 3\text{SO}_3 \cdot 33\text{H}_2\text{O}$	Hexag.
1261. Quetenite.....	$\text{MgO} \cdot \text{Fe}_2\text{O}_3 \cdot 3\text{SO}_3 \cdot 13\text{H}_2\text{O}$	Mono.
1262. Zincaluminite.....	$2\text{ZnSO}_4 \cdot 4\text{Zn}(\text{OH})_2 \cdot 6\text{Al}(\text{OH})_3 \cdot 5\text{H}_2\text{O}$	Hexag.
1263. Johannite.....	U, Cu, H, sulphate	Mono.
1264. Uranopilitite.....	$\text{CaU}_2\text{S}_2\text{O}_{11} \cdot 25\text{H}_2\text{O}$	Incrust.
<b>3. Tellurates</b>		
1265. Montanite.....	$\text{Bi}_2\text{O}_3 \cdot \text{TeO}_3 \cdot 2\text{H}_2\text{O}$	Earthy
1266. Emmonsite.....	$\text{Fe}_2\text{H}_2$ tellurate	Mono.
1267. Durdenite.....	$\text{Fe}_2(\text{TeO}_3)_3 \cdot 4\text{H}_2\text{O}$	Massive
1268. Chalcomenite.....	$\text{CuSeO}_3 \cdot 2\text{H}_2\text{O}$	Mono.
1269. Molybdomenite.....	Pb, selenite	Ortho.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
1249.	Yellow	2	.....	Cyprus	Iron
1250.	White	1	1	Halle	} Aluminum
1251.	White	.....	.....	Halle	
1252.	Snow white	1	2	Hungary	} Iron
1253.	Red	2	2	Sweden	
1254.	Yellow	2	2	Chile	} Aluminum
1255.	Green	3	2	Naples	
1256.	Yellow	2	2	Persia	} Iron
1257.	White	3.5	2.5	Colorado	
1258.	Yellow	2.5	3	Utah	Aluminum
1259.	Yellow	3	2	Upper Silesia	Iron
1260.	Colorless	2	1.7	Prussia	Aluminum
1261.	Brown	3	2	Chile	Calcium
1262.	White	2	2	Greece	Iron
1263.	Green	2	3	Joachimsthal	Zinc
1264.	Yellow	.....	3	Johanngeorgenst'dt	} Uranium
1265.	Yellow	.....	.....	Montana	
1266.	Green	5	.....	Arizona	} Bismuth
1267.	Yellow	2	.....	Honduras	
1268.	Blue	.....	3	Argentina	} Tellurium
1269.	White	.....	.....	Argentina	
					} Selenium

## COMPREHENSIVE

	Composition	Form
<b>XII. TUNGSTATES, MOLYB- DATES</b>		
1270. Wolframite.....	(Fe,Mn)WO <sub>4</sub>	Mono.
1271. Hübnerite.....	MnWO <sub>4</sub>	Mono.
1272. Scheelite.....	CaWO <sub>4</sub>	Tetrag.
1273. Cuprotungstite.....	CuWO <sub>4</sub>	Crystal.
1274. Powellite.....	CaMoWO <sub>4</sub>	Tetrag.
1275. Stolzite.....	PbWO <sub>4</sub>	Tetrag.
1276. Raspite.....	PbWO <sub>4</sub>	Mono.
1277. Wulfenite.....	PbMoO <sub>4</sub>	Tetrag.
1278. Reinite.....	FeWO <sub>4</sub>	Tetrag.
1279. Belonesite.....	MgMoO <sub>4</sub>	Tetrag.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
1270.	Black	5	7	Connecticut	Tungsten
1271.	Brown	5.5	7	Nevada	
1272.	White	4.5	5.9	North Carolina	
1273.	Green	4	.....	Chile	Molybdenum
1274.	Yellow	3.5	4	Michigan	
1275.	Green	2.7	7.8	Zinnwald	Tungsten
1276.	Yellow	2	.....	New South Wales	
1277.	Green	2.7	6.7	Arizona	Molybdenum
1278.	Brown	4	6.6	Japan	Tungsten
1279.	White	.....	.....	Vesuvius	Molybdenum



## COMPREHENSIVE

	Composition	Form
<b>XIII. ORGANIC ACIDS</b>		
<i>Oxalates, Mellates</i>		
1280. Whewellite.....	$\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}$	Mono.
1281. Oxammite.....	$(\text{NH}_4)_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$	Ortho.
1282. Humboldtine.....	$2\text{FeC}_2\text{O}_4 \cdot 3\text{H}_2\text{O}$	Capill.
1283. Mellite.....	$\text{Al}_2\text{C}_{12}\text{O}_{12} \cdot 18\text{H}_2\text{O}$	Tetrag.

LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
1280.	Colorless	2.5	.....	Saxony	Calcium
1281.	Yellowish	.....	.....	Peru	Ammonium
1282.	Yellow	2	2	Bohemia	Iron
1283.	Yellow	2	1.5	Bohemia	Mellitic acid

	Composition	Form
<b>XIV. HYDROCARBONS</b>		
1284. Scheererite.....	$C_nH_m$	Mono.
1285. Hatchettite.....	$C_nH_m$	Mono.
1286. Paraffin.....	$C_nH_m$	Amorph.
1287. Ozocerite.....	$C_nH_m$	Amorph.
1288. Zietrisikite.....	$C_nH_m$	Amorph.
1289. Chrismatite.....	$C_nH_m$	Amorph.
1290. Urpethite.....	$C_nH_m$	Amorph.
1291. Fichtelite.....	$C_{15}H_{28}$	Mono.
1292. Napalite.....	$C_3H_4$	Amorph.
1293. Amber.....	$C_nH_m$	Amorph.
1294. Succinite.....	$C_nH_m$	Amorph.
1295. Retinite.....	$C_nH_m$	Amorph.
1296. Gedanite.....	$C_nH_m$	Amorph.
1297. Glessite.....	$C_nH_m$	Amorph.
1298. Rumanite.....	$C_nH_m$	Amorph.
1299. Copalite.....	$C_nH_m$	Amorph.
1300. Bathvillite.....	$C_nH_m$	Amorph.
1301. Tasmanite.....	$C_nH_m$	Scales
1302. Dysodile.....	$C_nH_m$	Scales
1303. Geocerite.....	$C_{28}H_{56}O_2$	Waxy
1304. Leucopetrite.....	$C_{90}H_{84}O_4$	Waxy
1305. Pyrorretinite.....	$C_{20}H_{40}O_4$	Resinous
1306. Dopplerite.....	$C_nH_m$	Amorph.
1307. Idrialite.....	$C_{42}H_{14}O$	Earthy
1308. Posepnyte.....	$C_{22}H_{36}O_4$	Plates
1309. Petroleum, naphtha.....	$C_nH_{2n+2}$	Amorph.
1310. Pittasphalt.....	$C_nH_{2n+2}$	Viscid
1311. Asphaltum.....	$C_nH_{2n+2}$	Amorph.
1312. Elaterite.....	$C_nH_{2n+2}$	Amorph.
1313. Albertite.....	$C_nH_{2n+2}$	Amorph.
1314. Grahamite.....	$C_nH_{2n+2}$	Amorph.
1315. Gilsonite.....	$C_nH_{2n+2}$	Amorph.
1316. Mineral coal.....	$C_nH_{2n+2}$	Amorph.
1317. Anthracite.....	$C_nH_{2n+2}$	Amorph.
1318. Bituminous coal.....	$C_nH_{2n+2}$	Amorph.
1319. Coking coal.....	$C_nH_{2n}$	Amorph.
1320. Non-coking coal.....	$C_nH_{2n}$	Amorph.
1321. Cannel coal.....	$C_nH_{2n}$	Amorph.
1322. Torbanite.....	$C_nH_{2n}$	Amorph.
1323. Lignite.....	$C_nH_{2n}$	Amorph.
1324. Jet.....	$C_nH_{2n}$	Amorph.
1325. Peat.....	$C_nH_{2n}$	Amorph.

## LIST OF MINERALS

No.	Color	Hard- ness	Gravity	Locality	Chief Constituent or Use
1284.	Resinous	.....	1	Switzerland	Chemicals
1285.	White	1	.9	Switzerland	
1286.	Yellowish	.....	.....	England	Paraffin
1287.	Brown	.....	.9	Sicily	
1288.	Brown	.5	.9	Utah	
1289.	Yellow	.....	9	Saxony	
1290.	Brown	.....	.8	Urpeltz	Technical purposes
1291.	White	.....	.....	Bavaria	
1292.	Brown	2	.....	California	
1293.	Yellow	2	1	Baltic coast	
1294.	Yellow	2	1	Baltic coast	
1295.	Brown	.....	.....	Germany	
1296.	Brown	.....	.....	Baltic	Technical purposes
1297.	Brown	2	1	Baltic	
1298.	Brown	2	1	Roumania	Varnish
1299.	Yellow	.....	1	Tropics	
1300.	Brown	2	1	Scotland	Technical purposes
1301.	Brown	2	1	Tasmania	
1302.	Yellow	.....	1.2	Sicily	
1303.	White	.....	.....	Weissengels	Technical purposes
1304.	White	.....	1.2	Weissengels	
1305.	Yellow	2	1	Bohemia	
1306.	Black	2	1	Styria	
1307.	White	.....	.....	Idria	
1308.	Green	.....	.9	California	
1309.	Brown	.....	.6	United States	Oil
1310.	Greenish brown	.....	.....	Pennsylvania	
1311.	Black	.....	1	California	Technical purposes
1312.	Brown	.....	9	Derbyshire	
1313.	Black	1	1	Nova Scotia	
1314.	Black	2	1	West Virginia	
1315.	Black	2	1	Utah	
1316.	Black	2.5	1	United States	
1317.	Black	2	1	Pennsylvania	
1318.	Black	2.5	1	Pennsylvania	
1319.	Black	2	1.5	Virginia	Fuel
1320.	Black	2	1.5	Illinois	
1321.	Black	2	1.5	West Virginia	
1322.	Brown	2.2	1.1	Scotland	
1323.	Brown	1.5	1.1	Western states	Jewelry Fuel
1324.	Black	1.5	1.1	Wales	
1325.	Brown	1	1.1	Scotland	



## GENERAL INDEX



## GENERAL INDEX

The number preceding each mineral is the list number used in the List of Minerals; that following is the page number.

- Abbreviations, xxi
- 590. Abriachanite, 234
- 774. Acadialite, 242
- 81. Acanthite, 204
- 748. Achroite, 240
- 545. Acmite, 232
- 565. Actinolite, 232
- 979. Adamite, 254
- 969. Adelite, 252
- Admire, 44
- 493. Adularia, 134, 230
- 593. Aenigmatite, 234
- Aerolites, 43
- 931. Aeschynite, 250
- 436. Agaric Mineral, 226
- 291. Agate, 90, 218
- Agricola, 198
- 664. Agricolite, 236
- 70. Aguilarite, 204
- Ahnighito, 43
- 183. Aikinite, 210
- Airy's spiral, 88
- 90. Alabandite, 206
- 1202. Alabaster, 264
- 529. Alalite, 230
- 1313. Albertite, 274
- 505. Albite, 140, 230
- 374. Alexandrite, 222
- 55. Algodonite, 204
- 61. Alisonite, 204
- 998. Allacite, 254
- 725. Allanite, 240
- 10. Allemonite, 202
- 655. Allochroite, 236
- 139. Alloclasite, 208
- 31. Allopalladium, 202
- 883. Allophane, 246
- 646. Almandite, 159, 236
- 62. Altaite, 204
- 1194. Alumian, 264
- 1250. Aluminite, 268
- Aluminium, 45
- 1257. Alunite, 268
- 1228. Alunogen, 266
- 809. Alurgite, 244
- 21. Amalgam, 202
- 1242. Amaranthite, 266
- 502. Amazonstone, 230
- 1293. Amber, 187, 274
- 975. Amblygonite, 252
- 830. Amesite, 244
- 277. Amethyst, 88, 218
- 567. Amianthus, 232
- 1108. Ammiolite, 258
- Amorphous, 2
- 563. Amphibole, 152, 232
- 782. Analcime, 242
- 781. Analcite, 167, 242
- 704. Andalusite, 238
- 511. Andesine, 139, 230
- 152. Andorite, 210
- 648. Andradite, 159, 236
- 1084. Andrewsrite, 258
- 1172. Anglesite, 181, 264
- 1173. Anhydrite, 264
- 53. Animikite, 204
- 1024. Annabergite, 256
- 929. Annerodite, 250
- 803. Anomite, 242
- 514. Anorthite, 139, 142, 230
- 504. Anorthoclase, 138, 230
- 561. Anthophyllite, 232
- 1317. Anthracite, 192, 274
- 850. Antigorite, 244
- Antimonates, 175
- 12. Antimony, 31, 45, 202
- 1193. Antlerite, 264
- 945. Apatite, 175, 252
- 422. Aphrite, 226
- 747. Aphrizite, 240
- 831. Aphrosiderite, 244
- 1163. Apthitalite, 264
- 1222. Apjohnite, 266
- 658. Aplome, 236
- 758. Apophyllite, 240
- 1171. Apotome, 264
- 596. Aquamarine, 156, 234
- 450. Aragonite, 122, 226
- 735. Ardenite, 240
- 590a. Arfvedsonite, 234
- 421. Argentine, 226
- 66. Argentite, 51, 204
- Argon, 45
- 216. Argyrodite, 212



- Aristotle, 198  
 22. Arquerite, 202  
     Arsenates, 175  
     9. Arsenic, 31, 45, 202  
 1003. Arseniopleite, 254  
 997. Arseniosiderite, 254  
 325. Arsenolite, 220  
 134. Arsenopyrite, 67, 208  
 567. Asbestos, 155, 232  
 413. Asbolite, 224  
 1135. Ascharite, 262  
 305. Asmanite, 218  
 616. Aspasiolite, 234  
 1311. Asphaltum, 188, 274  
 910. Astrophyllite, 248  
 248. Atacamite, 214  
 1005. Atelestite, 254  
 1098. Atopite, 258  
 1069. Attacolite, 258  
 699. Auerlite, 238  
 1066. Augelite, 256  
     542. Augite, 147, 232  
     467. Aurichalcite, 228  
     365. Automolite, 222  
 1090. Autunite, 258  
     285. Aventurine, 218  
     Avicenna, 198  
     33. Awaruite, 202  
     Axial plane, 131  
 727. Axinite, 240  
 465. Azurite, 130, 228  
  
     Babbitt metal, 32  
     559. Babingtonite, 232  
     389. Baddeleyite, 222  
     726. Bagrationite, 240  
     536. Baikalite, 232  
     Bayly, W. S., ix, 200  
 1166. Barite, 179, 264  
     Barium, 45  
     592. Barkevikite, 234  
 1032. Barrandite, 256  
     Bartholinus, 112, 198  
     621. Barylite, 234  
     618. Barysilite, 234  
     457. Barytocalcite, 226  
     297. Basanite, 218  
     Base, 28  
     Basic Phosphates, 256  
     524. Bastite, 230  
     847. Bastite, 244  
     460. Bastnasite, 226  
 1300. Bathvillite, 274  
     401. Bauxite, 222  
     Baveno Twin, 134  
 1049. Bayldonite, 256  
     Bayley, W. S., ix, 200  
  
     1146. Bechilite, 262  
     203. Beegerite, 212  
     1279. Belonesite, 270  
     896. Bementite, 246  
     1073. Beraunite, 258  
     584. Bergamaskite, 234  
     785. Bergmannite, 242  
     877. Bergseife, 246  
     1067. Berlinite, 258  
     159. Berthierite, 210  
     739. Bertrandite, 240  
     594. Beryl, 156, 234  
     Beryllium, 45  
     944. Beryllonite, 252  
     71. Berzelianite, 204  
     937. Berzeliite, 252  
     Berzelius, 199  
     1113. Beudantite, 260  
     100. Beyrichite, 206  
     Biaxial, 89  
     1211. Bieberite, 266  
     1099. Bindheimite, 258  
     166. Binnite, 210  
     Biot, 170  
     801. Biotite, 170, 242  
     Bipyramid, 25  
     Birefringence, 55  
     262. Bischofite, 214  
     Bisectrix, 130  
     329. Bismite, 220  
     13. Bismuth, 31, 45, 202  
     44. Bismuthinite, 204  
     484. Bismutite, 228  
     129. Bismuto-smaltite, 208  
     458. Bismutospharite, 226  
     Bisphenoid, 29  
     1318. Bituminous coal, 274  
     381. Bixbyite, 222  
     Black Jack, 53  
     1215. Blodite, 266  
     Board of Museum Advisers, iii  
     1021. Bobierite, 254  
     412. Bog Manganesite, 224  
     397. Bog Ore, 222  
     876. Bole, 246  
     249a. Boleite, 214  
     1167. Bologna Stone, 264  
     610. Bonsdorffite, 234  
     1134. Boracite, 177, 262  
     Borates, 177  
     1144. Borax, 177, 262  
     1078. Borickite, 258  
     110. Bornite, 56, 206  
     Boron, 45  
     2. Bort, 20, 202  
     Boston Society, 95  
     Botryoidal, 89, 99

1253. Botryogen, 268  
 180. Boulangerite, 210  
     Bourg de Oisans Twin, 85  
 182. Bournonite, 210  
 1217. Boussingaultite, 266  
 849. Bowenite, 244  
     Boyle, 198  
     Brachydome, 26  
     Brachypinacoid, 27  
 984. Brackebuschite, 254  
 1013. Brandite, 254  
 380. Braunitz, 222  
     Bravais, 199  
 657. Bredbergite, 236  
 576. Breislakite, 232  
 103. Breithauptite, 206  
     Breithaupt, 199  
 441. Breunnerite, 226  
     Brewster, 199  
 762. Brewsterite, 242  
 1188. Brochantite, 264  
 1151. Broggerite, 262  
     Bromine, 45  
 454. Bromite, 226  
 230. Bromyrite, 214  
 173. Brongniardite, 210  
 522. Bronzite, 230  
 391. Brookite, 222  
     Brown, A. P., 200  
 403. Brucite, 222  
     Brush, 200  
 1040. Brushite, 256  
     Buck and Company, ix  
 720. Bucklandite, 240  
 301. Buhrstone, 218  
     Bunsen, 199  
 341. Bunsenite, 220  
 557. Bustamite, 232  
     Butler, G. M., 200  
 512. Bytownite, 139  
  
 1025. Cabrerite, 256  
 315. Cacholong, 218  
 1072. Cacozenite, 258  
     Cadmium, 45  
     Caesium, 45  
 280. Cairngorm, 88, 218  
 1320. Caking coal, 274  
 740. Calamine, 240  
 143. Calaverite, 208  
 1077. Calcioferrite, 258  
 983. Calciovolborthite, 254  
 415. Calcite, 112, 226  
     Calcium, 45  
 434. Calc-sinter, 226  
 434. Calc-Tufa, 116  
 1191. Caledonite, 264  
  
 1034. Callanite, 256  
 221. Calomel, 214  
 959. Campylite, 252  
 532. Canaanite, 232  
     Canada balsam, 129  
 628. Cancrinite, 236  
 217. Canfieldite, 212  
 1321. Cannel coal, 274  
     Canon Diablo, 44  
 602. Cappelenite, 234  
 1182. Caracolite, 264  
     Carbon, 45  
     3. Carbonado, 20, 202  
     Carbonates, 112  
     Carlsbad Twin, 134  
 939. Carminite, 252  
 260. Carnallite, 214  
 287. Carnelian, 90, 218  
 1155. Carnotite, 262  
 742. Carpholite, 240  
 1245. Carphosiderite, 266  
 114. Carrollite, 206  
 604. Caryocerite, 234  
 897. Caryopillite, 246  
 382. Cassiterite, 107, 222  
 1240. Castanite, 266  
 806. Caswellite, 242  
 601. Catapleiite, 234  
 284. Cat's-eye, 218  
 375. Cat's Eye, 222  
 1168. Cawk, 264  
 866. Celadonite, 246  
 1170. Celestite, 180, 264  
 517. Celsian, 230  
 886. Cenosite, 246  
 228. Cerargyrite, 214  
 744. Cerite, 240  
     Cerium, 45  
 456. Cerussite, 127, 226  
 333. Cervantite, 220  
 360. Ceylonite-Pleonaste, 220  
 773. Chabazite, 242  
 1212. Chalcantinite, 266  
 286. Chalcedony, 80, 218  
     77. Chalcocite, 52, 57, 204  
 1268. Chalcomenite, 268  
 409. Chalcophanite, 224  
 1055. Chalcophyllite, 256  
 115. Chalcopyrite, 57, 206  
 1083. Chalcosiderite, 258  
 157. Chalcostibite, 210  
 336. Chalcotrichite, 220  
 429. Chalk, 226  
 840. Chamosite, 244  
 1081. Chenevixite, 258  
 296. Chert, 90  
 466. Chessylite, 228

- Chester, 124  
 503. Chesterlite, 230  
 705. Chiasolite, 238  
 1074. Childrenite, 258  
 57. Chilenite, 204  
 246. Chiolite, 214  
 148. Chiviatite, 210  
 521. Chladnite, 230  
 239. Chloralluminite, 214  
 Chlorine, 45  
 814. Chloritoid, 244  
 237. Chloromagnesite, 214  
 549. Chloromelanite, 232  
 800. Chloropal, 246  
 836. Chlorophaeite, 244  
 615. Chlorophyllite, 234  
 361. Chlorospinel, 222  
 991. Chondrasenite, 254  
 731. Chondrodite, 240  
 1289. Chrismatite, 274  
 372. Chromite, 104, 107, 222  
 Chromium, 45  
 373. Chrysoberyl, 222  
 889. Chrysocolla, 246  
 665. Chrysolite, 236  
 288. Chrysoprase, 90, 218  
 852. Chrysotile, 244  
 1028. Churchite, 256  
 879. Cimolite, 246  
 94. Cinnabar, 55, 206  
 640. Cinnamon-Stone, 236  
 996. Cirrolite, 254  
 279. Citrine, 88, 218  
 209. Clarite, 212  
 326. Claudetite, 220  
 63. Clausthalite, 204  
 354. Clay Iron-stone, 220  
 398. Clay-ironstone, 222  
 Cleavage, 16  
 508. Cleavelandite, 230  
 1152. Cleveite, 262  
 4. Cliftonite, 202  
 821. Clinocllore, 244  
 987. Clinoclasite, 254  
 Clinodomes, 129  
 741. Clinohedrite, 240  
 733. Clinohumite, 240  
 Clinopinacoid, 129  
 722. Clinzoisite, 240  
 Cloisonné work, 33  
 1316. Coal, 190  
 Cobalt, 45  
 119. Cobaltite, 206  
 537. Coccolite, 232  
 130. Cockscomb pyrite, 65  
 1065. Coeruleolactite, 256  
 39. Cohenite, 202  
 1319. Coking Coal, 274  
 1141. Colemanite, 177, 262  
 Colloidal Silica, 90  
 1007. Collophanite, 254  
 884. Collyrite, 246  
 651. Colophonite, 236  
 89. Coloradoite, 206  
 925. Columbite-Tantalite, 250  
 Columbium, 45  
 352. Columnar hematite, 220  
 Comprehensive list, 201  
 1048. Conicalcrite, 256  
 863. Connarite, 246  
 1184. Connellite, 264  
 1209. Copalite, 274  
 1239. Copiapite, 266  
 18. Copper, 38, 45, 202  
 1225. Coquimbite, 266  
 1052. Cornwallite, 256  
 829. Corundophilite, 244  
 346. Corundum, 94, 220  
 121. Corynite, 206  
 171. Cosalite, 210  
 243. Cotunnite, 214  
 687. Couseranite, 238  
 95. Covellite, 206  
 Crayon, 173  
 378. Crednerite, 222  
 306. Cristobalite, 218  
 589. Crocidolite, 234  
 1178. Crocoite, 264  
 Cronstedt, 198  
 838. Cronstedite, 244  
 75. Crookesite, 204  
 591. Crossite, 234  
 245. Cryolite, 80, 214  
 Crystal form, 2  
 113. Cubanite, 206  
 Cube, 33  
 Cullinan diamond, 22  
 249b. Cumengite, 214  
 572. Cumingtonite, 232  
 335. Cuprite, 92, 220  
 149. Cuprobismutite, 210  
 233. Cuproiodargyrite, 214  
 60. Cuproplumbite, 204  
 1273. Cuprotungstite, 270  
 730. Cuspidine, 240  
 707. Cyanite, 163, 238  
 1237. Cyanotrichite, 266  
 516. Cyclopote, 230  
 219. Cyindrite, 212  
 694. Cyprine, 238  
 1249. Cyprusite, 268

792. Damourite, 242  
 Dana, ix, 199  
 135. Danaite, 208  
 635. Danalite, 236  
 700. Danburite, 238  
 573. Dannemorite, 232  
 1123. Darapskite, 260  
 708. Datolite, 238  
 Daubree, 109, 199  
 259. Daubreeite, 214  
 112. Daubreelite, 206  
 597. Davidsonite, 234  
 255. Daviesite, 214  
 470. Dawsonite, 228  
 982. Dechenite, 254  
 833. Delessite, 244  
 650. Demantoid, 236  
 1105. Derbyllite, 258  
 Des Cloizeaux, 199  
 980. Descloizite, 254  
 1110. Destinezite, 260  
 855. Deweyllite, 246  
 832. Diabantite, 244  
 1109. Diadochite, 260  
 538. Diallage, 232  
 1. Diamond, 5, 202  
 177. Diaphorite, 210  
 393. Diaspore, 222  
 Dichroscope, 96  
 1010. Dickinsonite, 254  
 1223. Dietrichite, 266  
 1126. Dietzeite, 260  
 989. Dihydrate, 254  
 Dimorphism, 123  
 527. Diopside, 147, 230  
 678. Dioptase, 238  
 Dioscorides, 61  
 Dioxides, 107  
 Diploid, 62  
 686. Dipyre, 238  
 Dispersion, 19  
 Disulphides, 206  
 Ditetragonal Bipyramid, 58  
 Ditetragonal Prism, 58  
 Ditrigonal polar, 166  
 Doctor of Medicine, 3  
 Dodecahedron, 72  
 416. Dog-tooth Spar, 226  
 1190. Dolerophanite, 264  
 439. Dolomite, 117, 226  
 Dome, 26  
 54. Domeykite, 204  
 1306. Dopplerite, 274  
 Double refraction, 55  
 261. Douglasite, 214  
 Drills, 20  
 993. Dufrenite, 254  
 169. Dufrenoyssite, 210  
 749. Dumortierite, 240  
 974. Durangite, 252  
 1267. Durdenite, 268  
 917. Dysanalyte, 248  
 50. Dyscrasite, 204  
 366. Dysluite, 222  
 1302. Dysodile, 274  
 Dysprosium, 45  
 953. Earthy Apatite; Osteolite, 252  
 1102. Ecdemite, 258  
 578. Edenite, 232  
 783. Edingtonite, 242  
 624. Elaeolite, 234  
 Elasticity coefficient, 119  
 1312. Elaterite, 274  
 16. Electrum, 202  
 Elements, 4  
 Elements, List of, 5, 45  
 600. Elpidite, 234  
 229. Embolite, 214  
 181. Embrithite, 210  
 595. Emerald, 156, 234  
 349. Emery, 94, 220  
 1266. Emmonsite, 268  
 156. Emplectite, 210  
 208. Enargite, 212  
 960. Endlichite, 252  
 520. Enstatite, 146, 230  
 Entantimorphous, 86  
 1075. Eosphorite, 258  
 213. Epiboulangerite, 212  
 834. Epichlorite, 244  
 491. Epididymite, 230  
 717. Epidote, 238  
 214. Epigenite, 212  
 763. Epistilbite, 242  
 1204. Epsomite, 266  
 Erbium, 45  
 988. Erinite, 254  
 Erni, 200  
 Erubiscite, 56  
 1023. Erythrite, 254  
 264. Erythrosiderite, 216  
 98. Erythrozincite, 206  
 613. Esmarkite, 234  
 1260. Ettringite, 268  
 73. Eucairite, 204  
 1047. Euchroite, 256  
 599. Eucolite, 234  
 710. Euclase, 238  
 626. Eucryptite, 236  
 598. Eudialyte, 234  
 636. Eulytite, 236  
 951. Eupyrchroite, 252  
 835. Euralite, 244

- Europium, 45  
 981. Eusynchite, 254  
 933. Euxenite, 250  
 1064. Evansite, 256  
   Extraordinary ray, 55
195. Fahlerz, 74  
 611. Fahlunite, 234  
 1014. Fairfieldite, 254  
 192. Falkenhaynite, 212  
 211. Famantinite, 212  
   Farrington, O. C., ix, 200  
 544. Fassaite, 232  
 672. Fayalite, 238  
 1252. Felsobanyite, 268  
 923. Fergusonite, 250  
   Ferrites, 104  
 1231. Ferronatrite, 266  
 1243. Fibroferrite, 266  
 1291. Fichtelite, 274  
 253. Fiedlerite, 214  
 1011. Fillowite, 254  
   Finds, 43  
 310. Fire Opal, 218  
 1058. Fischerite, 256  
 1000. Flinkite, 254  
 295. Flint, 218  
 322. Float stone, 220  
 451. Flosferri, 226  
 266. Fluellite, 216  
   Fluorescence, 79  
   Fluorine, 45  
 257. Fluocerite, 214  
 235. Fluorite, 78, 214  
 419. Fontainebleau Limestone, 226  
   Foote, W. M., ix  
 273. Footeite, 216  
 1044. Forbesite, 256  
   Ford, 199, 200  
   Forest, 3  
 670. Forsterite, 236  
   Fossil resins, 187  
   Fouque, 199  
 558. Fowlerite, 232  
   Fracture, 16  
 218. Franckeite, 212  
 948. Francolite, 252  
 369. Franklinite, 106, 222  
 196. Freibergite, 212  
 176. Freieslebenite, 210  
 860. French chalk, 246  
 679. Friedelite, 238  
   80. Friesite, 204  
 796. Fuchsite, 242  
 869. Fuller's Earth, 174  
   Fulton, 200
711. Gadolinite, 238  
   Gadolinite, 45  
 364. Gahnite, 222  
   59. Galena, 48, 204  
 158. Galenobismutite, 210  
   Gallium, 45  
 619. Ganomalite, 234  
 755. Ganophyllite, 240  
 638. Garnet, 158, 236  
 857. Garnierite, 246  
 588. Gastaldite, 234  
 475. Gay Lussite, 228  
 270. Gearsutite, 216  
 1296. Gedanite, 274  
 562. Gedrite, 232  
 692. Gehlenite, 238  
   Geikielite, 248  
 918. Geikielite, 248  
   *General Guide*, v  
 856. Genthite, 246  
 1303. Geocerite, 274  
 202. Geocronite, 212  
 1122. Gerhardtite, 260  
   Germanium, 45  
 120. Gersdorffite, 206  
 133. Geyerite, 208  
 321. Geyserite, 92, 218  
   Ghost quartz, 87  
 405. Gibbsite, 222  
 625. Gieseckite, 236  
 794. Gilbertite, 242  
 1315. Gilsonite, 274  
 311. Girasol, 218  
 768. Gismondite, 242  
 1164. Glauberite, 264  
 138. Glaucodot, 208  
 684. Glaucolite, 238  
 867. Glauconite, 246  
 587. Glaucophane, 234  
 326. Glaudetite, 220  
 1297. Glessite, 274  
   Glide planes, 47  
 1246. Glockerite, 266  
   Glucinum, 45  
 778. Gmelinite, 242  
   Goethe, 102  
 394. Goethite, 101, 222  
   15. Gold, 32, 45, 202  
 1205. Goslarite, 266  
 1086. Goyazite, 258  
 1314. Grahamite, 274  
 893. Graminite, 246  
   5. Graphite, 23, 202  
   Gratacap, 200  
   96. Greenockite, 206  
 905. Greenovite, 248  
 639. Grossularite, 159, 236

- Groth, ix, 200  
 906. Grothite, 248  
 107. Grunauite, 206  
 574. Grunerite, 232  
 86. Guadalcazarite, 206  
 45. Guanajuatite, 204  
 908. Guarinite, 248  
 187. Guitermanite, 210  
 1156. Gummite, 262  
 1199. Gypsum, 182, 264  
 757. Gyrolite, 240  
  
 Haidinger, 65  
 1038. Haidingerite, 256  
 224. Halite, 76, 214  
 872. Halloysite, 246  
 Haloids, 76  
 1221. Halotrichite, 266  
 1132. Hamburgite, 262  
 973. Hamlinite, 252  
 Hancock County, 87  
 1186. Hanksite, 264  
 972. Harderite, 252  
 Hardin County, 79  
 Hardness, Scale of, 17  
 766. Harmotome, 242  
 729. Harstigit, 240  
 586. Hastingsite, 234  
 1285. Hatchettite, 274  
 920. Hatchettolite, 250  
 101. Hauchecornite, 206  
 117. Hauerite, 206  
 Haly, 198  
 631. Hallynite, 236  
 804. Haughtonite, 242  
 376. Hausmannite, 222  
 775. Haydenite, 242  
 Heat conductivity, 30  
 534. Hedenbergite, 232  
 962. Hedyphane, 252  
 1143. Heintzite, 262  
 290. Heliotrope, 90  
 Helium, 45  
 634. Helvite, 236  
 1046. Hemaifibrite, 256  
 350. Hematite, 97, 220  
 1001. Hematolite, 254  
 Hemihedral, 70  
 Hemimorphic, 93  
 Hennepin, 191  
 1082. Henwoodite, 258  
 130. Hepatic Pyrite, 64  
 363. Heroynite, 222  
 1235. Herrengrundite, 266  
 777. Herschelite, 242  
 68. Hessite, 204  
 Hexagonal system, 68  
 Hexatetrahedron, 15  
 Hexoctahedron, 12  
 761. Heulandite, 242  
 547. Hiddenite, 232  
 930. Hielmite, 250  
 247. Hieratite, 214  
 560. Hiortdahlite, 232  
 895. Hisingerite, 246  
 History of Study of Minerals, 198  
 894. Hoferite, 246  
 1022. Hoernesite, 254  
 Holbrook, 44  
 Holland, J. G., 33  
 Holoaxial, 82  
 Homestead, 44  
 709. Homilite, 238  
 Hope diamond, 22  
 1009. Hopeite, 254  
 105. Horbachite, 206  
 577. Hornblende, 153, 232  
 296. Hornstone, 90, 218  
 51. Horsfordite, 204  
 671. Hortonolite, 236  
 Hours, iii  
 1138. Howlite, 262  
 225. Huantajayite, 214  
 1271. Hübnerite, 270  
 837. Hullite, 244  
 691. Humboldtite, 238  
 1282. Humboldtine, 272  
 732. Humite, 240  
 52. Huntite, 204  
 1043. Hureaulite, 256  
 Hutchinson, Charles F., iii  
 Huygens, 112  
 641. Hyacinth, 236  
 696. Hyacinth, 238  
 319. Hyalite, 218  
 500. Hyalophane, 138, 230  
 667. Hyalosiderite, 236  
 620. Hyalotekite, 234  
 Hydrated sesquioxides, 100  
 428. Hydraulic limestone, 226  
 Hydrous Sulphates, 264  
 1147. Hydroboracite, 262  
 Hydrocarbons, 187  
 469. Hydrocerussite, 228  
 1177. Hydrocyanite, 264  
 Hydrofluoric acid, 81  
 Hydrogen, 45  
 479. Hydrogiobertite, 228  
 478. Hydromagnesite, 228  
 790. Hydronephelite, 242  
 313. Hydrophene, 218  
 236. Hydrophilite, 214  
 407. Hydrotalcite, 224

- Hydrous carbonates, 228  
 Hydrous chlorides, 214  
 Hydrous oxides, 222  
 468. Hydrozincite, 228  
 523. Hypersthene, 146, 230
338. Ice, 220  
 418. Iceland Spar, 115, 226  
 Iddings, J. P., 200  
 668. Iddingsite, 236  
 1307. Idrialite, 274  
 1227. Ihleite, 266  
 1203. Ilesite, 264  
 387. Ilmenorutile, 222  
 356. Ilmenite, 220  
 734. Ilvaite, 240  
 874. Indianaite, 246  
 515. Indianite, 230  
 746. Indicolite, 240  
 Indium, 45  
 754. Inesite, 240  
 324. Infusorial Earth, 220  
 Intermediate plagioclases, 143  
 Intermediate oxides, 220  
 Iodine, 45  
 231. Iodobromite, 214  
 234. Iodyrite, 214  
 609. Iolite, 234  
 26. Iridium, 45, 202  
 27. Iridosmine, 202  
 32. Iron, 42, 45, 202  
 1045. Isoclasite, 256  
 Isomorphism, 122  
 300. Itacolumite, 218
371. Jacobsite, 222  
 Jackson, B. H., 200  
 Jade, 156  
 548. Jadeite, 150, 232  
 67. Jalpaite, 204  
 168. Jamesonite, 210  
 697. Jargon, 238  
 1258. Jarosite, 268  
 298. Jasper, 90, 218  
 845. Jefferisite, 244  
 541. Jeffersonite, 232  
 1131. Jeremejevite, 262  
 Jersey County, 40
1324. Jet, 274  
 1263. Johannite, 268  
 Jo Daviess County, 49  
 Johannsen, Albert, 200  
 911. Johnstrupite, 248  
 199. Jordanite, 212  
 47. Joseite, 204  
 34. Josephinite, 202  
 Jubilee, 21
585. Kaersutite, 234  
 1183. Kainite, 264  
 627. Kaliophilite, 236  
 124. Kallilite, 206  
 36. Kamacite, 202  
 1236. Kamarensite, 266  
 826. Kammererite, 244  
 Kaolins, 246  
 869. Kaolinite, 174, 246  
 581. Kataforite, 234  
 1085. Kehoeite, 258  
 907. Keilhauite, 248  
 737. Kentrolite, 240  
 145. Kermesite, 208  
 1197. Kieserite, 264  
 204. Kilbrickenite, 212  
 Kimberley, 5  
 Kimberlite, 21  
 Klaproth, 94  
 165. Klaprotholite, 210  
 673. Knebelite, 238  
 916. Knopite, 248  
 1247. Knoxvillite, 266  
 172. Kobellite, 210  
 Kohinoor, 21  
 579. Koksharovite, 232  
 23. Kongsbergite, 202  
 1036. Koninckite, 256  
 752. Kornerupine, 240  
 823. Kotschubeite, 244  
 1026. Kottigite, 256  
 Kraus, E. H., 200  
 367. Kreittonite, 222  
 263. Kremersite, 214  
 142. Krennerite, 208  
 1229. Krohnkite, 266  
 Kryptom, 45  
 Kunz, G. F., 200  
 582. Kupfferite, 234
512. Labradorite, 139, 230  
 Lacroix, ix  
 1139. Lagonite, 262  
 414. Lampadite, 224  
 1189. Lanarkite, 264  
 763. Langbanite, 240  
 1165. Langbeinite, 264  
 1234. Langite, 266  
 480. Lansfordite, 228  
 476. Lanthanite, 228  
 Lanthanum, 45  
 1140. Larderellite, 262  
 947. Lasurapatite, 252  
 772. Laubanite, 242  
 769. Laumontite, 242  
 252. Laurionite, 214  
 126. Laurite, 206

1125. Lautarite, 260  
 555. Lavenite, 232  
 533. Lavrovite, 232  
 242. Lawrencite, 214  
 743. Lawsonite, 240  
     Lawyer, 3  
 994. Lazulite, 254  
 633. Lazurite, 236  
     20. Lead, 45, 202  
 1179. Leadhillite, 264  
 1195. Lecontite, 264  
 903. Lederite, 248  
     72. Lehrbachite, 204  
 770. Leonbardite, 242  
 1216. Leonite, 266  
 799. Lepidolite, 172, 242  
 808. Lepidomelane, 244  
     Letter of transmittal, v  
 543. Leucaugite, 232  
 822. Leuchtenbergite, 244  
 518. Leucite, 145, 230  
 1051. Leucochalcite, 256  
 1304. Leucopetrite, 274  
 607. Leucophanite, 234  
 132. Leucopyrite, 208  
 779. Levynite, 242  
     Lewis, J. V., 200  
 1106. Lewisite, 258  
 977. Libethenite, 252  
 486. Liebigite, 228  
     Liebisch, 199  
 1323. Lignite, 192, 274  
 901. Ligurite, 248  
 186. Lillianite, 210  
 396. Limonite, 103, 222  
 1192. Linarite, 264  
 1114. Lindackerite, 260  
     111. Linnaeite, 206  
     Linnaeus, 198  
 1080. Liroconite, 258  
 1063. Liskeardite, 256  
     List of Illustrations, xiii  
 942. Lithiophilite, 252  
     Lithium, 45  
 427. Lithographic Stone, 226  
 870. Lithomarge, 246  
 147. Livingstonite, 210  
 131. Lollingite, 208  
     Long Island, 44  
 162. Lorandite, 210  
 1116. Lossenite, 260  
 1214. Loweite, 266  
 1259. Lowigite, 268  
     497. Loxoclase, 230  
 1071. Ludlamite, 258  
 1128. Ludwigite, 262  
 425. Lumachelle, 226  
 1115. Lunebergite, 260  
     Lutecium, 45  
 210. Luzonite, 212  
 714. Mackintoshite, 238  
     Macon County, 40  
     Macrodome, 27  
     Macropinacoid, 25  
 370. Magnesioferrite, 222  
 440. Magnesite, 120, 226  
     Magnesium, 45  
 368. Magnetite, 105, 222  
 464. Malachite, 128, 228  
 528. Malacolite, 230  
 1208. Mallardite, 266  
     Malus, 112  
     Manebach Twin, 134  
 949. Manganapatite, 252  
 824. Manganchlorite, 244  
     Manganese, 45  
 395. Manganite, 100, 222  
     Manganites, 104  
 805. Manganophyllite, 242  
 340. Manganosite, 220  
 1004. Manganostiibite, 254  
 130. Marcasite, 63, 208  
 811. Margarite, 244  
 793. Margarodite, 242  
 688. Marialite, 238  
     83. Marmatite, 206  
 851. Marmolite, 244  
     223. Marshite, 214  
 1041. Martinite, 256  
 355. Martite, 220  
 1160. Mascagnite, 264  
 513. Maskelynite, 230  
 817. Masonite, 244  
 1224. Masrite, 266  
 343. Massicot, 220  
 160. Matildite, 210  
     250. Matlockite, 214  
 1404. Maucherite, 208  
 1107. Mauzeilite, 258  
 1076. Mazapilite, 258  
 681. Meionite, 238  
 652. Melanite, 236  
 603. Melanocerite, 234  
 307. Melanophlogite, 218  
 738. Melanotekite, 240  
 1207. Melanterite, 266  
 690. Melillite, 238  
 608. Meliphanite, 234  
 1283. Mellite, 272  
 109. Melonite, 206  
 251. Mendipite, 214  
 200. Menenghinitite, 212  
 317. Menilite, 218



19. Mercury, 2, 40, 45, 202  
 802. Merxene, 242  
 Merrill, G. P., 200  
 442. Mesitite, 226  
 787. Mesolite, 242  
 1015. Messelite, 254  
 85. Metacinnabarite, 206  
   Metallurgist, 4  
   Metasilicates, 230  
 43. Metastibnite, 404  
 1256. Metavoltine, 268  
 35. Meteoric Iron, 202  
   Meteorites, 21, 42  
 161. Miargyrite, 210  
   Micas, 168  
 1169. Michel-levyite, 264  
 921. Microcline, 250  
 501. Microcline, 137, 230  
 629. Microsommitite, 236  
   Miers, ix  
 232. Miersite, 214  
 956. Miesite, 252  
 489. Milarite, 230  
 314. Milk Opal, 218  
 281. Milky Quartz, 218  
   Miller, W. H., 8, 199  
 99. Millerite, 206  
   Millsbaugh, Charles F., iii  
 958. Mimetite, 252  
   Miner, 4  
   Mineral, Definition of, 1  
 1316. Mineral coal, 274  
   Minerals, Abundance of, 1  
   Minerals, Numbers of, 1  
   Mineral production of Ill., 1  
   Minerals, uses of, 1, 287  
   Minister, 3  
 377. Minium, 222  
   Minnesota Mine, 39  
 1196. Mirabilite, 264  
 1187. Misenite, 264  
   Mitscherlich, 199  
 1097. Mixite, 258  
 685. Mizzonite, 238  
   Mohs, 199  
   Moissan, 199  
   Molybdates, 270  
 49. Molybdenite, 204  
   Molybdenum, 45  
 331. Molybdate, 220  
 1269. Molybdomenite, 48, 268  
 240. Molysite, 214  
 936. Monazite, 252  
 938. Monimolite, 252  
   Monoclinic axes, 30  
   Monoclinic system, 30, 129  
   Monosulphides, 204  
   Monoxides, 92  
 1265. Montanite, 268  
 669. Monticellite, 236  
 880. Montmorillonite, 246  
 492. Moonstone, 135  
 760. Mordenite, 240  
 1206. Morenosite, 266  
   Morian, 88  
 946. Moroxite, 252  
 912. Mosandrite, 248  
 986. Mottramite, 254  
   Moses, A. J., 200  
 569. Mountain Cork, 156, 232  
 568. Mountain Leather, 156, 232  
   Mukerop, 44  
 498. Murchisonite, 230  
 791. Muscovite, 168, 242  
 1101. Nadorite, 258  
 144. Nagyagite, 208  
 417. Nail-head Spar, 226  
   Names of minerals, 196  
 222. Nantokite, 214  
 1292. Napalite, 274  
 1233. Natrochalcite, 266  
 784. Natrolite, 168, 242  
 943. Natrophilite, 252  
 473. Natron, 228  
   Natural gas, 189  
   Naumann, 199  
 65. Naumannite, 204  
   Neodymium, 45  
   Neon, 45  
 808. Neotocite, 246  
 623. Nephelite, 234  
 550. Nephrite, 232  
 566. Nephrite, 156, 232  
 914. Neptunite, 248  
 472. Nesquehonite, 228  
 28. Nevyanskite, 202  
 1042. Newberyite, 256  
 878. Newtonite, 246  
 102. Niccolite, 206  
   Nickel, 45  
 128. Nickel-Skutterudite, 208  
   Nicol prism, 115, 129  
 386. Nigrine, 222  
   Niobates, Tantalates, 175  
 1118. Niter, 260  
   Nitrates, etc., 260  
 1121. Nitrobarite, 260  
 1119. Nitrocalcite, 260  
   Nitrogen, 45  
 1124. Nitroglauberite, 260  
 1120. Nitromagnesite, 260  
 1153. Nivenite, 262  
 258. Nocerite, 214

1320. Non-coking coal, 274  
 891. Nontronite, 246  
 1130. Nordenskiöldine, 262  
 751. Nordmarkite, 240  
 462. Northupite, 226  
 632. Noselite, 236  
 957. Nussierite, 252  
  
 1103. Ochrolite, 258  
     Octahedron, 5  
 390. Octahedrite, 222  
 780. Offretite, 242  
 756. Okenite, 240  
 91. Oldhamite, 206  
 509. Oligoclase, 139, 230  
 445. Oligonite, 226  
 976. Olivenite, 252  
 666. Olivine, 236  
 539. Omphacite, 232  
     88. Onofrite, 206  
 292. Onyx, 90, 218  
 430. Oolite, 226  
 308. Opal, 91, 218  
 316. Opal-agate, 218  
 854. Ophicalcite, 244  
     Ordinary ray, 55  
     Organic acid salts, 187  
     Orloff, 21  
     41. Orpiment, 204  
 492. Orthoclase, 133, 230  
     Orthodomes, 129  
     Orthopinacoid, 27  
     Orthorhombic system, 25  
     Osmium, 45  
 818. Ottrelite, 244  
     Owen, Charles L., iii  
     Oxalates, Mellates, 272  
 1281. Oxammite, 272  
     Oxides, 82  
     Oxychlorides, 214  
     Oxygen, 45  
 780. Ozarkite, 242  
 1287. Ozocerite, 274  
  
 268. Pachnolite, 216  
     30. Palladium, 45, 202  
 1286. Paraffin, 274  
 798. Paragonite, 242  
 1251. Paraluminite, 268  
 345. Pameleconite, 220  
     Parameters, 8  
     Paramorph, 123  
     Paramorphism, 123  
 580. Pargasite, 232  
 459. Parisite, 226  
     Parsons, C. L., 200  
 663. Partschinite, 236  
  
 683. Passauite, 238  
     Payne, Edward W., iii  
     Peacock ore, 56  
 206. Pearceite, 212  
 320. Pearl sinter, 218  
     Peary, R. E., 43  
 1325. Peat, 192, 274  
     525. Peckhamite, 230  
     552. Pectolite, 232  
 1059. Peganite, 256  
     Penfield, S. L., 200  
     254. Penfieldite, 214  
     825. Penninite, 244  
     Pentagonal dodecahedron, 62  
     92. Pentlandite, 206  
     Peoria County, 40  
     Percussion figure, 169  
     249. Percylite, 214  
     339. Periclase, 220  
     507. Pericline, 141, 230  
     506. Peristerite, 230  
     915. Perovskite, 248  
     499. Perthite, 230  
     488. Petalite, 230  
 1309. Petroleum. Naphtha, 188, 274  
     69. Petzite, 204  
     776. Phacolite, 242  
     Pharmacist, 3  
 1037. Pharmacolite, 256  
 1070. Pharmacosiderite, 258  
     676. Phenacite, 238  
 1230. Phillipite, 266  
     Phillips, A. H., 200  
     765. Phillipsite, 242  
     807. Phlogopite, 242  
     871. Pholerite, 246  
     868. Pholidolite, 246  
     461. Phosgenite, 226  
     Phosphates, 175, 260  
     Phosphorescence, 19  
     950. Phosphorite, 252  
     Phosphorus, 45  
 1031. Phosphosiderite, 256  
 1093. Phosphuranylite, 258  
     820. Phyllite, 244  
     702. Physalite, 238  
 1220. Pickeringite, 266  
     362. Picotite-Chrome Spinel, 222  
     723. Picroepidote, 240  
     853. Picrolite, 244  
 1218. Picromerite, 266  
 1017. Picropharmacolite, 254  
     724. Piedmonite, 240  
     Pinacoid, 27  
 1129. Pinakiolite, 262  
     892. Pinguite, 246  
     797. Pinite, 242

1142. Pinnoite, 262  
 Pirsson, L. V., 200  
 474. Pirssonite, 228  
 1209. Pisanite, 266  
 431. Pisolite, 155, 226  
 443. Pistomesite, 226  
 1154. Pitchblende, 262  
 1310. Pittasphalt, 274  
 1111. Pitticite, 260  
 163. Plagionite, 210  
 290. Plasma, 218  
 Plaster of Paris, 182  
 25. Platinum, 41, 45, 202  
 Plattner, 199  
 388. Plattnerite, 222  
 Pleistocene, 37  
 38. Plessite, 202  
 Pliny, 61  
 1087. Plumbogummite, 258  
 Polar, 68, 93  
 384. Polianite, 222  
 519. Pollucite, 230  
 656. Polyadelphite, 236  
 207. Polyargyrite, 212  
 205. Polybasite, 212  
 617. Polychrollite, 234  
 934. Polycrase, 250  
 106. Polydymite, 206  
 1219. Polyhalite, 266  
 932. Polymignite, 250  
 955. Polysphaerite, 252  
 Pope County, 79  
 1308. Posepnyite, 274  
 Potassium, 45  
 1274. Powellite, 270  
 Praseodymium, 45  
 289. Prase, 218  
 309. Precious Opal, 218  
 Preface, ix  
 728. Prehnite, 240  
 Prism, 27  
 828. Prochlorite, 244  
 267. Prosopite, 216  
 Prospector, 4  
 190. Proustite, 68, 212  
 84. Przibramite, 206  
 379. Pseudobrookite, 222  
 990. Pseudomalachite, 254  
 Pseudomorph, 123  
 827. Pseudophite, 244  
 873. Pseudosteatite, 246  
 410. Psilomelane, 224  
 985. Psittacinite, 254  
 759. Ptilolite, 240  
 940. Pucherite, 252  
 703. Pyonite, 238  
 Pyramid, 25  
 612. Pyrargillite, 234  
 189. Pyrrargyrite, 68, 210  
 653. Pyreneite, 236  
 116. Pyrite, 206  
 Pyritohedron, 62  
 408. Pyroaurite, 224  
 919. Pyrochlore, 250  
 404. Pyrochroite, 222  
 392. Pyrolusite, 110, 222  
 954. Pyromorphite, 176, 252  
 644. Pyrope, 159, 236  
 357. Pyrophanite, 220  
 1008. Pyrophosphorite, 254  
 882. Pyrophyllite, 246  
 1305. Pyroretinite, 274  
 680. Pyrosmalite, 238  
 193. Pyrostilpnite, 212  
 526. Pyroxene, 230  
 Pyroxene group, 145  
 922. Pyrrhite, 250  
 104. Pyrrhotite, 55, 206  
 275. Quartz, 82, 218  
 303. Quartzine, 218  
 299. Quartzite, 90, 218  
 1226. Quenstedtite, 226  
 1261. Quetenite, 268  
 Quincy, 124  
 Radium, 45  
 1244. Raimondite, 266  
 271. Ralstonite, 216  
 Rammelsberg, 199  
 137. Rammelsbergite, 208  
 1276. Raspite, 270  
 170. Rathite, 210  
 614. Raumite, 234  
 40. Realgar, 204  
 353. Red Ocherous hematite, 220  
 1016. Reddingite, 254  
 1248. Redingtonite, 266  
 Refraction, Index of, 18, 19  
 Regent diamond, 21  
 Regular system, 16  
 1278. Reinite, 270  
 482. Remingtonite, 228  
 René Just Haüy, 198  
 Reniform, 99  
 861. Reneselaerite, 246  
 312. Resin Opal, 218  
 Resins, 187  
 848. Retinalite, 244  
 1295. Retinite, 274  
 1002. Retzian, 254  
 150. Rezbanyite, 210  
 1027. Rhabdophanite, 256

1096. Rhagite, 258  
       Radium, 45  
 1136. Rhodizite, 262  
 447. Rhodochrosite, 121, 226  
 645. Rhodolite, 236  
 556. Rhodonite, 151, 232  
       Rhombohedron, 70, 114  
 496. Rhyacolite, 230  
 1079. Richellite, 258  
 575. Richterite, 232  
 588a. Riebackite, 234  
 913. Rinkite, 248  
 194. Rittingerite, 212  
       Rock, 2  
       Rock Island, 124  
 437. Rock-meal, 226  
       Roentgen, 47  
 643. Romanzovite, 236  
       Romé de L'Isle, 104, 198  
 622. Roebbingite, 234  
 1100. Romeite, 258  
 1232. Romerite, 266  
 810. Roscoelite, 244  
       Rose, 198  
 1012. Roselite, 254  
 553. Rosenbuschite, 232  
 278. Rose-quartz, 218  
 654. Rothofite, 236  
 713. Rowlandite, 238  
       Rubidium, 45  
 348. Ruby, 95, 220  
 359. Ruby Spinel-Magnesia Spinel, 220  
 426. Ruin-marble, 226  
 1298. Rumanite, 274  
 843. Rumpfitte, 244  
       Ruthenium, 45  
 385. Rutile, 109, 222  
  
 423. Saccharoidal limestone, 226  
 136. Safflorite, 208  
 283. Sagenite, 218  
 227. Sal Ammoniac, 214  
       Saline, 44  
 535. Salite, 232  
 816. Salmite, 244  
 1210. Salvadorite, 266  
       Salt, 76  
       Salt, Origin of, 77  
       Salt River, 44  
 928. Samarekite, 250  
       Samarium, 45  
       Sancy, 22  
 195. Sandbergite, 74  
 191. Sanguinite, 212  
 495. Sanidine, 230  
 865. Saponite, 246  
 347. Sapphire, 94, 220  
 282. Sapphire-quartz, 218  
 753. Sapphirine, 240  
 680. Sarcolite, 238  
 293. Sard, 90  
 293. Sardonyx, 90, 218  
       Sargasso Seas, 2  
 971. Sarkinite, 252  
 155. Sartorite, 210  
 406. Sassolite, 224  
 420. Satin Spar, 226  
 1201. Satin Spar, 264  
 238. Scacchite, 214  
       Scalenohehrons, 72, 114  
       Scandium, 45  
 175. Schapbachite, 210  
 1272. Scheelite, 270  
 1284. Scheererite, 274  
 540. Schefferite, 232  
 164. Schirmerite, 210  
 771. Schneiderite, 242  
 662. Schorlomite, 236  
 304. Schreibersite, 202  
 885. Schrotterite, 246  
       Schungite, 202  
 256. Schwartzembergite, 214  
 197. Schwatzite, 74, 212  
 786. Scolecite, 242  
 1020. Scorodite, 256  
 718. Scorza, 238  
       Selenium, 45  
 1200. Selenite, 182, 264  
       8. Selensulphur, 202  
 241. Sellaite, 214  
 174. Semseyite, 210  
       Senarmont, 47  
 327. Senarmontite, 220  
 862. Sepiolite, 246  
 795. Sericite, 242  
 846. Serpentine, 172, 244  
 1238. Serpierite, 266  
       Sesquioxides, 94  
 812. Seybertite, 244  
       Shah of Persia, 22  
 424. Shell-marble, 226  
       Shepardson, F. W., iii, v  
 444. Siderite, 120, 226  
       Siderolites, 43  
 1254. Sideronatrite, 268  
       Silicates, 132  
 294. Siliceous sinter, 218  
 302. Silicified wood, 218  
       Silicon, 45  
 706. Sillimanite, 238  
       17. Silver, 37, 45, 202  
       Sinter, 92  
 924. Sipylite, 250  
       29. Siserskite, 202

815. Sismondine, 244  
 127. Skutterudite, 208  
 118. Smaltite-Chloanthite, 206  
 570. Smaragdite, 232  
 875. Smectite, 246  
 448. Smithsonite, 121, 226  
 630. Sodalite, 236  
 1117. Soda Niter, 260  
     Sodium, 45  
     Sohncke, 199  
     Sorby, 199  
     South Star, 22  
 864. Spadaite, 246  
 1185. Spangolite, 264  
     Spearhead pyrite, 65  
     Specific gravity, 17  
     Specific heat, 23  
 351. Specular hematite, 220  
     Specularite, 99  
 125. Sperryllite, 206  
 647. Spessartite, 159, 236  
 1062. Sphaerite, 256  
 449. Sphaerocobaltite, 226  
     82. Sphalerite, 52, 206  
 900. Sphene, 248  
 446. Spherosiderite, 226  
 358. Spinel, 104, 220  
     Spinel Twins, 17  
 902. Spinther, 248  
 965. Spodiosite, 252  
 546. Spodumene, 232  
 952. Staffelite, 252  
 432. Stalactites, 116, 226  
 433. Stalagmite, 116, 226  
 215. Stannite, 212  
 276. Star-quartz, 218  
 750. Staurolite, 240  
 859. Steatite, 246  
     Steno, 81, 198  
 201. Stephanite, 212  
 1042a. Stercorite, 256  
     79. Sternbergite, 204  
 334. Stibiconite, 220  
     42. Stibnite, 46, 204  
 767. Stilbite, 166, 242  
 841. Stilpnomelane, 244  
 881. Stolpenite, 246  
 1275. Stolzite, 270  
 383. Stream Tin, 222  
 605. Streenstrupine, 234  
 1030. Strengite, 256  
     842. Strigovite, 244  
     78. Stromeyerite, 204  
 455. Strontianite, 126, 226  
     Strontium, 45  
 1006. Struvite, 254  
     58. Stutzite, 204  
 185. Stylotypite, 210  
     Subsilicates, 240  
     Sublimation deposits, 24  
 642. Succinite, 236  
 1294. Succinite, 274  
 1148. Sulfoborite, 262  
     Sulphantimonites, 68  
     Sulpharsenates, 212  
     Sulpharsenites, 68, 210  
     Sulphates, 179  
     Sulphides, 46  
     Sulphides of metals, 204  
 1181. Sulphohalite, 264  
     Sulpho-salts, 210  
     7. Sulphur, 24, 45, 202  
     Sulphuric acid, 56, 63  
 220. Sulvanite, 212  
     Summary, 194  
 154. Sundtite, 210  
 510. Sunstone, 230  
 1180. Susannite, 264  
 1127. Sussezite, 262  
 963. Svabite, 252  
 1112. Svanbergite, 260  
 108. Sychnodymite, 206  
 141. Sylvanite, 208  
 226. Sylvite, 76, 214  
     Symbol, 8  
     Symmetry axes, 18  
 1020. Symplectite, 254  
 999. Synadelphite, 254  
 1213. Syngeinitite, 266  
     583. Syntagmatite, 234  
 1133. Szaibelyite, 262  
 1198. Szmikite, 264  
 265. Tachhydrite, 216  
     37. Taenite, 202  
 1050. Tagilite, 256  
 858. Talc, 172, 246  
 967. Talktriplite, 252  
 272. Tallingite, 216  
 188. Tapalpite, 210  
 926. Tapiolite, 250  
     Tantalum, 45  
 978. Tarbuttite, 254  
 452. Tarnowitzite, 226  
 1301. Tasmanite, 274  
 995. Tavistockite, 254  
 1161. Taylorite, 264  
     Tellurates, 268  
 330. Tellurite, 220  
     11. Tellurium, 45, 202  
 483. Tengerite, 228  
 198. Tennantite, 212  
 344. Tenorite, 220  
 674. Tephroite, 238

- Terbium, 45  
 463. Teschemacherite, 228  
 46. Tetradymite, 204  
 195. Tetrahedrite, 72, 212  
 Tetragonal system, 57  
 Tetrahedral Twins, 13  
 Tetrahedron, 13  
 Tetrahexahedron, 34  
 719. Thallite, 238  
 Thallium, 45  
 887. Thaumassite, 246  
 1162. Thenardite, 264  
 471. Thermonatrite, 228  
 Theophrastus, 99  
 438. Thinolite, 226  
 Thin section, 129  
 269. Thomsenolite, 216  
 788. Thomsonite, 242  
 698. Thorite, 238  
 Thorium, 45  
 1158. Thorogummite, 262  
 716. Thulite, 238  
 Thulium, 45  
 839. Thuringite, 244  
 87. Tiemannite, 206  
 970. Tilasite, 252  
 337. Tile ore, 220  
 64. Tilkerodite, 204  
 24. Tin, 45, 202  
 Titanates, 248  
 659. Titaniferous garnet, 236  
 899. Titanite, 248  
 Titanium, 45  
 904. Titanomorphite, 248  
 701. Topaz, 162, 238  
 649. Topazolite, 236  
 1322. Torbanite, 274  
 1088. Torbernite, 258  
 745. Tourmaline, 164, 240  
 Trapezohedron, 9  
 530. Traversellite, 230  
 435. Travertine, 116, 226  
 564. Tremolite, 153, 232  
 Triassic, 21  
 1018. Trichalcite, 254  
 Triclinic system, 139  
 304. Tridymite, 218  
 677. Trimerite, 238  
 941. Triphylite, 252  
 966. Triplite, 252  
 968. Triploidite, 252  
 323. Tripolite, 220  
 1103a. Trippkeite, 258  
 1175. Tripstone, 264  
 1104. Tripuhyte, 258  
 Trisectahedron, 10  
 606. Tritomite, 234  
 1094. Trogerite, 258  
 93. Troilite, 206  
 1068. Trolleite, 258  
 477. Trona, 228  
 909. Tscheffkinit, 248  
 Tschermak, ix  
 1054. Tyrolite, 256  
 244. Tysonite, 214  
 Tungstates, 185  
 Tungsten, 45  
 332. Tungstite, 220  
 399. Turgite, 222  
 1060. Turquoise, 256  
 Tutton, ix  
 1145. Ulexite, 262  
 123. Ullmannite, 206  
 76. Umangite, 204  
 University of Chicago Press, ix  
 571. Uralite, 232  
 Uranates, 177  
 1149. Uraninite, 178, 262  
 Uranium, 45  
 1150. Uranniobite, 262  
 1159. Uranosphaerite, 262  
 1091. Uranospinit, 258  
 1092. Uranocircite, 258  
 888. Uranophane, 246  
 1264. Uranopilite, 268  
 485. Uranothallite, 228  
 1290. Urpethite, 274  
 Use of minerals, 197  
 1241. Utahite, 266  
 661. Uvarovite, 159, 236  
 494. Valencianite, 230  
 328. Valentinite, 220  
 961. Vanadinite, 252  
 Vanadium, 45  
 Van Horn, F. R., 200  
 1033. Variscite, 256  
 819. Venasquite, 244  
 844. Vermiculite, 244  
 693. Vesuvianite, 238  
 1056. Veszelyite, 256  
 Vicinal faces, 78  
 531. Violan, 230  
 1019. Vivianite, 254  
 487. Voglite, 228  
 1052. Volborthite, 256  
 1255. Voltaite, 268  
 146. Voltzite, 208  
 Von Lang, 146  
 1174. Vulpinite, 264  
 411. Wad, 224  
 964. Wagnerite, 252

1095. Walpurgite, 258  
 1039. Wapplerite, 256  
     Ward, Henry B., iii  
 1061. Wardite, 256  
 167. Warrenite, 210  
 1137. Warwickite, 262  
 1057. Wavellite, 256  
 153. Webnerite, 210  
     48. Wehrlite, 204  
     Welcome Nugget, 35  
 764. Wellsite, 242  
     Werner, 199  
 682. Wernerite, 238  
     Wherry, E. T., 200  
 1280. Whewellite, 272  
     56. Whitneyite, 204  
     Willamette, 44  
 675. Willemite, 238  
 122. Willyamite, 206  
     Winchell, A. M., and N. H., 200  
 721. Withamite, 240  
 453. Witherite, 125, 226  
 184. Wittichenite, 210  
 402. Wocheinite, 222  
 554. Wohlerite, 232  
 140. Wolfachite, 208  
 1270. Wolframite, 185, 270  
     Wollaston, 199  
     551. Wollastonite, 232  
 318. Wood Opal, 218  
 1277. Wulfenite, 186, 270  
     97. Wurtzite, 206  
  
 992. Xantharsenite, 254  
 813. Xanthophyllite, 244  
  
 400. Xanthosiderite, 222  
 212. Xanthosonite, 212  
 935. Xenotime, 252  
  
     Yellowstone National Park, 24  
     Ytterbium, 45  
 660. Yttergranat, 236  
 712. Yttrialite, 238  
     Yttrium, 45  
 274. Yttrocerite, 216  
 235a. Yttrofluorite, 214  
 1157. Yttrogummite, 262  
 927. Yttrantalite, 250  
  
 813. Zanthophyllite, 244  
 481. Zaratite, 228  
     Zeolites, 166  
 1035. Zepharovichite, 256  
 1089. Zeunerite, 258  
 1288. Zietrisikite, 274  
     14. Zinc, 45, 202  
 1262. Zincaluminite, 268  
     342. Zincite, 93, 220  
     151. Zinkenite, 210  
 1176. Zinkosite, 264  
 800. Zinnwaldite, 242  
 695. Zircon, 160, 238  
     Zirconium, 45  
     Zirkel, 199  
 715. Zoisite, 238  
     74. Zorgite, 204  
 637. Zunyite, 236







B-297

AUTHOR

TITLE

DATE DUE

PRINTED IN U.S.A.

